



VISHAL SIR'S
ACADEMY

JEE Main 2025 (January)

Chapter-wise Qs Bank

Mathematics

Content

Questions

1. Basic of Mathematics	4
2. Quadratic Equation	5
3. Complex Number	7
4. Sequences and Series	9
5. Permutation Combination	12
6. Binomial Theorem	14
7. Statistics	16
8. Matrices	17
9. Determinants	19
10. Probability	22
11. Sets and Relations	26
12. Functions	29
13. Limits	32
14. Continuity and Differentiability	33
15. Application of Derivatives	35
16. Indefinite Integration	36
17. Definite Integration	37
18. Area Under Curves	40
19. Differential Equations	43
20. Straight Lines	46
21. Circle	49
22. Parabola	51
23. Ellipse	53
24. Hyperbola	55
25. Trigonometric Ratios & Identities	56
26. Trigonometric Equations	57
27. Inverse Trigonometric Functions	58
28. Vector Algebra	59
29. Three Dimensional Geometry	62

Answer Keys

67

Solutions

1. Basic of Mathematics	69
2. Quadratic Equation	70
3. Complex Number	73
4. Sequences and Series	78
5. Permutation Combination	84
6. Binomial Theorem	87
7. Statistics	92
8. Matrices	94
9. Determinants	100
10. Probability	103
11. Sets and Relations	108
12. Functions	112
13. Limits	118
14. Continuity and Differentiability	121
15. Application of Derivatives	124
16. Indefinite Integration	126
17. Definite Integration	128
18. Area Under Curves	134
19. Differential Equations	142
20. Straight Lines	148
21. Circle	155
22. Parabola	159
23. Ellipse	165
24. Hyperbola	169
25. Trigonometric Ratios & Identities	172
26. Trigonometric Equations	173
27. Inverse Trigonometric Functions	174
28. Vector Algebra	176
29. Three Dimensional Geometry	183

Basic of Mathematics

JEE Main 2025 January

Q1. The product of all solutions of the equation $e^{5(\log_e x)^2+3} = x^8, x > 0$, is :

(1) $e^{8/5}$

(2) $e^{6/5}$

(3) e^2

(4) e



Quadratic Equation

JEE Main 2025 January

Q1. The number of solutions of the equation $\left(\frac{9}{x} - \frac{9}{\sqrt{x}} + 2\right)\left(\frac{2}{x} - \frac{7}{\sqrt{x}} + 3\right) = 0$ is:

- (1) 2
- (2) 3
- (3) 1
- (4) 4

Q2. The sum, of the squares of all the roots of the equation $x^2 + |2x - 3| - 4 = 0$, is

- (1) $3(3 - \sqrt{2})$
- (2) $6(3 - \sqrt{2})$
- (3) $6(2 - \sqrt{2})$
- (4) $3(2 - \sqrt{2})$

Q3. If the set of all $a \in \mathbf{R}$, for which the equation $2x^2 + (a - 5)x + 15 = 3a$ has no real root, is the interval (α, β) , and $X = \{x \in \mathbf{Z} : \alpha < x < \beta\}$, then $\sum_{x \in X} x^2$ is equal to :

- (1) 2109
- (2) 2129
- (3) 2119
- (4) 2139

Q4. If the equation $a(b - c)x^2 + b(c - a)x + c(a - b) = 0$ has equal roots, where $a + c = 15$ and $b = \frac{36}{5}$, then $a^2 + c^2$ is equal to

Q5. The product of all the rational roots of the equation $(x^2 - 9x + 11)^2 - (x - 4)(x - 5) = 3$, is equal to

- (1) 14
- (2) 21
- (3) 28
- (4) 7

Q6. Let α_θ and β_θ be the distinct roots of $2x^2 + (\cos \theta)x - 1 = 0, \theta \in (0, 2\pi)$. If m and M are the minimum and the maximum values of $\alpha_\theta^4 + \beta_\theta^4$, then $16(M + m)$ equals :

- (1) 24

Quadratic Equation

JEE Main 2025 January

(2) 25

(3) 17

(4) 27



Complex Number

JEE Main 2025 January

Q1. The number of complex numbers z , satisfying $|z| = 1$ and $\left|\frac{z}{\bar{z}} + \frac{\bar{z}}{z}\right| = 1$, is :

- (1) 4
- (2) 8
- (3) 10
- (4) 6

Q2. Let $|z_1 - 8 - 2i| \leq 1$ and $|z_2 - 2 + 6i| \leq 2$, $z_1, z_2 \in \mathbf{C}$. Then the minimum value of $|z_1 - z_2|$ is :

- (1) 13
- (2) 10
- (3) 3
- (4) 7

Q3. Let α, β be the roots of the equation $x^2 - ax - b = 0$ with $\text{Im}(\alpha) < \text{Im}(\beta)$. Let $P_n = \alpha^n - \beta^n$. If $P_3 = -5\sqrt{7}i$, $P_4 = -3\sqrt{7}i$, $P_5 = 11\sqrt{7}i$ and $P_6 = 45\sqrt{7}i$, then $|\alpha^4 + \beta^4|$ is equal to _____.

Q4. If $\alpha + i\beta$ and $\gamma + i\delta$ are the roots of $x^2 - (3 - 2i)x - (2i - 2) = 0$, $i = \sqrt{-1}$, then $\alpha\gamma + \beta\delta$ is equal to :

- (1) -2
- (2) 6
- (3) -6
- (4) 2

Q5. Let z_1, z_2 and z_3 be three complex numbers on the circle $|z| = 1$ with $\arg(z_1) = \frac{-\pi}{4}$, $\arg(z_2) = 0$ and $\arg(z_3) = \frac{\pi}{4}$. If $|z_1\bar{z}_2 + z_2\bar{z}_3 + z_3\bar{z}_1|^2 = \alpha + \beta\sqrt{2}$, $\alpha, \beta \in \mathbf{Z}$, then the value of $\alpha^2 + \beta^2$ is :

- (1) 24
- (2) 29
- (3) 41
- (4) 31

Q6. Let O be the origin, the point A be $z_1 = \sqrt{3} + 2\sqrt{2}i$, the point $B(z_2)$ be such that $\sqrt{3}|z_2| = |z_1|$ and $\arg(z_2) = \arg(z_1) + \frac{\pi}{6}$. Then

Complex Number

JEE Main 2025 January

(1) area of triangle ABO is $\frac{11}{\sqrt{3}}$

(2) ABO is an obtuse angled isosceles triangle

(3) area of triangle ABO is $\frac{11}{4}$

(4) ABO is a scalene triangle

Q7. Let the curve $z(1+i) + \bar{z}(1-i) = 4, z \in \mathbb{C}$, divide the region $|z-3| \leq 1$ into two parts of areas α and β . Then $|\alpha - \beta|$ equals :

(1) $1 + \frac{\pi}{2}$

(2) $1 + \frac{\pi}{3}$

(3) $1 + \frac{\pi}{6}$

(4) $1 + \frac{\pi}{4}$

Q8. Let $\left| \frac{\bar{z}-i}{2\bar{z}+i} \right| = \frac{1}{3}, z \in \mathbb{C}$, be the equation of a circle with center at C . If the area of the triangle, whose vertices are at the points $(0,0), C$ and $(\alpha, 0)$ is 11 square units, then α^2 equals:

(1) 50

(2) 100

(3) $\frac{81}{25}$

(4) $\frac{121}{25}$

Q9. If α and β are the roots of the equation $2z^2 - 3z - 2i = 0$, where $i = \sqrt{-1}$, then $16 \cdot \operatorname{Re}\left(\frac{\alpha^{19} + \beta^{19} + \alpha^{11} + \beta^{11}}{\alpha^{15} + \beta^{15}}\right) \cdot \operatorname{Im}\left(\frac{\alpha^{19} + \beta^{19} + \alpha^{11} + \beta^{11}}{\alpha^{15} + \beta^{15}}\right)$ is equal to

(1) 441

(2) 398

(3) 312

(4) 409

Q10. Let integers $a, b \in [-3, 3]$ be such that $a + b \neq 0$. Then the number of all possible ordered pairs

(a, b) , for which $\left| \frac{z-a}{z+b} \right| = 1$ and $\begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 1, z \in \mathbb{C}$, where ω and ω^2 are the roots of $x^2 + x + 1 = 0$, is equal to _____.

Sequences and Series

JEE Main 2025 January

- Q1.** If the first term of an A.P. is 3 and the sum of its first four terms is equal to one-fifth of the sum of the next four terms, then the sum of the first 20 terms is equal to
- (1) -1080
 - (2) -1020
 - (3) -1200
 - (4) -120
- Q2.** The roots of the quadratic equation $3x^2 - px + q = 0$ are 10^{th} and 11^{th} terms of an arithmetic progression with common difference $\frac{3}{2}$. If the sum of the first 11 terms of this arithmetic progression is 88, then $q - 2p$ is equal to _____.
- Q3.** In an arithmetic progression, if $S_{40} = 1030$ and $S_{12} = 57$, then $S_{30} - S_{10}$ is equal to :
- (1) 525
 - (2) 510
 - (3) 515
 - (4) 505
- Q4.** Consider an A. P. of positive integers, whose sum of the first three terms is 54 and the sum of the first twenty terms lies between 1600 and 1800. Then its 11^{th} term is :
- (1) 90
 - (2) 84
 - (3) 122
 - (4) 108
- Q5.** The number of 3 -digit numbers, that are divisible by 2 and 3, but not divisible by 4 and 9, is _____.
- Q6.** Let T_r be the r^{th} term of an A.P. If for some m , $T_m = \frac{1}{25}$, $T_{25} = \frac{1}{20}$, and $20 \sum_{r=1}^{25} T_r = 13$, then $5m \sum_{r=m}^{2m} T_r$ is equal to
- (1) 98
 - (2) 126
 - (3) 142
 - (4) 112

Sequences and Series

JEE Main 2025 January

- Q7.** The interior angles of a polygon with n sides, are in an A.P. with common difference 6° . If the largest interior angle of the polygon is 219° , then n is equal to
- Q8.** Let $a_1, a_2, \dots, a_{2024}$ be an Arithmetic Progression such that $a_1 + (a_5 + a_{10} + a_{15} + \dots + a_{2020}) + a_{2024} = 2233$. Then $a_1 + a_2 + a_3 + \dots + a_{2024}$ is equal to _____
- Q9.** Let $a_1, a_2, a_3, \dots$ be a G.P. of increasing positive terms. If $a_1 a_5 = 28$ and $a_2 + a_4 = 29$, then a_6 is equal to:
- (1) 628
(2) 812
(3) 526
(4) 784
- Q10.** If $7 = 5 + \frac{1}{7}(5 + \alpha) + \frac{1}{7^2}(5 + 2\alpha) + \frac{1}{7^3}(5 + 3\alpha) + \dots$, then the value of α is :
- (1) $\frac{6}{7}$
(2) 6
(3) $\frac{1}{7}$
(4) 1
- Q11.** If $\sum_{r=1}^n T_r = \frac{(2n-1)(2n+1)(2n+3)(2n+5)}{64}$, then $\lim_{n \rightarrow \infty} \sum_{r=1}^n \left(\frac{1}{T_r}\right)$ is equal to :
- (1) 0
(2) $\frac{2}{3}$
(3) 1
(4) $\frac{1}{3}$
- Q12.** For positive integers n , if $4a_n = (n^2 + 5n + 6)$ and $S_n = \sum_{k=1}^n \left(\frac{1}{a_k}\right)$, then the value of $507S_{2025}$ is :
- (1) 540
(2) 675
(3) 1350
(4) 135
- Q13.** The value of $\lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{k^3 + 6k^2 + 11k + 5}{(k+3)!} \right)$ is:

Sequences and Series

JEE Main 2025 January

(1) $4/3$

(2) 2

(3) $7/3$

(4) $5/3$

Q14. Suppose that the number of terms in an A.P. is $2k, k \in N$. If the sum of all odd terms of the A.P. is 40, the sum of all even terms is 55 and the last term of the A.P. exceeds the first term by 27, then k is equal to :

(1) 6

(2) 5

(3) 8

(4) 4

Q15. Let $S_n = \frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \dots$ upto n terms. If the sum of the first six terms of an A.P. with first term $-p$ and common difference p is $\sqrt{2026 S_{2025}}$, then the absolute difference between 20th and 15th terms of the A.P. is

(1) 20

(2) 90

(3) 45

(4) 25

Q16. Let $\langle a_n \rangle$ be a sequence such that $a_0 = 0, a_1 = \frac{1}{2}$ and $2a_{n+2} = 5a_{n+1} - 3a_n, n = 0, 1, 2, 3, \dots$. Then $\sum_{k=1}^{100} a_k$ is equal to

(1) $3a_{99} - 100$

(2) $3a_{100} - 100$

(3) $3a_{99} + 100$

(4) $3a_{100} + 100$

Permutation Combination

JEE Main 2025 January

- Q1.** If all the words with or without meaning made using all the letters of the word "KANPUR" are arranged as in a dictionary, then the word at 440th position in this arrangement, is :
- (1) PRNAUK
 - (2) PRKANU
 - (3) PRKAUN
 - (4) PRNAKU
- Q2.** Group A consists of 7 boys and 3 girls, while group B consists of 6 boys and 5 girls. The number of ways, 4 boys and 4 girls can be invited for a picnic if 5 of them must be from group A and the remaining 3 from group B, is equal to :
- (1) 8750
 - (2) 9100
 - (3) 8925
 - (4) 8575
- Q3.** The number of different 5 digit numbers greater than 50000 that can be formed using the digits 0 , 1, 2, 3, 4, 5, 6, 7 , such that the sum of their first and last digits should not be more than 8 , is
- (1) 4608
 - (2) 5720
 - (3) 5719
 - (4) 4607
- Q4.** The number of natural numbers, between 212 and 999 , such that the sum of their digits is 15 , is
- Q5.** Let P be the set of seven digit numbers with sum of their digits equal to 11 . If the numbers in P are formed by using the digits 1,2 and 3 only, then the number of elements in the set P is :
- (1) 173
 - (2) 164
 - (3) 158
 - (4) 161
- Q6.** The number of 6-letter words, with or without meaning, that can be formed using the letters of the word MATHS such that any letter that appears in the word must appear at least twice, is _____.

Permutation Combination

JEE Main 2025 January

Q7. From all the English alphabets, five letters are chosen and are arranged in alphabetical order. The total number of ways, in which the middle letter is 'M', is :

- (1) 5148
- (2) 6084
- (3) 4356
- (4) 14950


Q8. In a group of 3 girls and 4 boys, there are two boys B_1 and B_2 . The number of ways, in which these girls and boys can stand in a queue such that all the girls stand together, all the boys stand together, but B_1 and B_2 are not adjacent to each other, is :

- (1) 96
- (2) 144
- (3) 120
- (4) 72

Q9. The number of words, which can be formed using all the letters of the word "DAUGHTER", so that all the vowels never come together, is

- (1) 36000
- (2) 37000
- (3) 34000
- (4) 35000

Q10. The number of ways, 5 boys and 4 girls can sit in a row so that either all the boys sit together or no two boys sit together, is _____ -



Binomial Theorem

JEE Main 2025 January

- Q1.** The sum of all rational terms in the expansion of $\left(1 + 2^{1/2} + 3^{1/2}\right)^6$ is equal to
- Q2.** For some $n \neq 10$, let the coefficients of the 5th, 6th and 7th terms in the binomial expansion of $(1 + x)^{n+4}$ be in A.P. Then the largest coefficient in the expansion of $(1 + x)^{n+4}$ is:
- (1) 20
 - (2) 10
 - (3) 35
 - (4) 70
- Q3.** Suppose A and B are the coefficients of 30th and 12th terms respectively in the binomial expansion of $(1 + x)^{2n-1}$. If $2A = 5B$, then n is equal to :
- (1) 22
 - (2) 20
 - (3) 21
 - (4) 19
- Q4.** Let α, β, γ and δ be the coefficients of x^7, x^5, x^3 and x respectively in the expansion of $\left(x + \sqrt{x^3 - 1}\right)^5 + \left(x - \sqrt{x^3 - 1}\right)^5, x > 1$. If u and v satisfy the equations $\alpha u + \beta v = 18$ and $\gamma u + \delta v = 20$ then $u + v$ equals :
- (1) 5
 - (2) 3
 - (3) 4
 - (4) 8
- Q5.** If in the expansion of $(1 + x)^p(1 - x)^q$, the coefficients of x and x^2 are 1 and -2, respectively, then $p^2 + q^2$ is equal to :
- (1) 18
 - (2) 13
 - (3) 8
 - (4) 20

Binomial Theorem

JEE Main 2025 January

- Q6.** Let the coefficients of three consecutive terms T_r, T_{r+1} and T_{r+2} in the binomial expansion of $(a+b)^{12}$ be in a G.P. and let p be the number of all possible values of r . Let q be the sum of all rational terms in the binomial expansion of $(\sqrt[4]{3} + \sqrt[3]{4})^{12}$. Then $p + q$ is equal to :
- (1) 283
(2) 287
(3) 295
(4) 299
- Q7.** The least value of n for which the number of integral terms in the Binomial expansion of $(\sqrt[3]{7} + \sqrt[12]{11})^n$ is 183, is :
- (1) 2184
(2) 2196
(3) 2148
(4) 2172
- Q8.** The remainder, when 7^{103} is divided by 23, is equal to :
- (1) 6
(2) 17
(3) 9
(4) 14
- Q9.** If $\sum_{r=0}^5 \frac{{}^{11}C_{2r+1}}{2r+2} = \frac{m}{n}$, $\gcd(m, n) = 1$, then $m - n$ is equal to _____
- Q10.** If $\alpha = 1 + \sum_{r=1}^6 (-3)^{r-1} {}^{12}C_{2r-1}$, then the distance of the point $(12, \sqrt{3})$ from the line $\alpha x - \sqrt{3}y + 1 = 0$ is _____.
- Q11.** If $\sum_{r=1}^{30} \frac{r^2 ({}^{30}C_r)^2}{{}^{30}C_{r-1}} = \alpha \times 2^{29}$, then α is equal to _____

Statistics

JEE Main 2025 January

Q1. The variance of the numbers 8, 21, 34, 47, ..., 320 is

Q2. Marks obtained by all the students of class 12 are presented in a frequency distribution with classes of equal width.

Let the median of this grouped data be 14 with median class interval 12-18 and median class frequency 12. If the number of students whose marks are less than 12 is 18, then the total number of students is

(1) 52

(2) 48

(3) 44

(4) 40

Q3. For a statistical data $x_1, x_2, \dots, x_{10}$ of 10 values, a student obtained the mean as 5.5 and $\sum_{i=1}^{10} x_i^2 = 371$. He later found that he had noted two values in the data incorrectly as 4 and 5, instead of the correct values 6 and 8, respectively. The variance of the corrected data is

(1) 9

(2) 5

(3) 7

(4) 4

Q4. Let $x_1, x_2, \dots, x_{10}$ be ten observations such that $\sum_{i=1}^{10} (x_i - 2) = 30$, $\sum_{i=1}^{10} (x_i - \beta)^2 = 98$, $\beta > 2$, and their variance is $\frac{4}{5}$. If μ and σ^2 are respectively the mean and the variance of $2(x_1 - 1) + 4\beta, 2(x_2 - 1) + 4\beta, \dots, 2(x_{10} - 1) + 4\beta$, then $\frac{\beta\mu}{\sigma^2}$ is equal to :

(1) 100

(2) 120

(3) 110

(4) 90

Matrices

JEE Main 2025 January

Q1. Let $A = [a_{ij}]$ be 3×3 matrix such that $A \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$, $A \begin{bmatrix} 4 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ and $A \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, then a_{23} equals :

- (1) -1
- (2) 2
- (3) 1
- (4) 0

Q2. Let $A = [a_{ij}]$ be a matrix of order 3×3 , with $a_{ij} = (\sqrt{2})^{i+j}$. If the sum of all the elements in the third row of A^2 is $\alpha + \beta\sqrt{2}$, $\alpha, \beta \in \mathbf{Z}$, then $\alpha + \beta$ is equal to :

- (1) 280
- (2) 224
- (3) 210
- (4) 168

Q3. For a 3×3 matrix M , let $\text{trace}(M)$ denote the sum of all the diagonal elements of M . Let A be a 3×3 matrix such that $|A| = \frac{1}{2}$ and $\text{trace}(A) = 3$. If $B = \text{adj}(\text{adj}(2A))$, then the value of $|B| + \text{trace}(B)$ equals :

- (1) 56
- (2) 132
- (3) 174
- (4) 280

Q4. Let M denote the set of all real matrices of order 3×3 and let $S = \{-3, -2, -1, 1, 2\}$. Let

$$S_1 = \{A = [a_{ij}] \in M : A = A^T \text{ and } a_{ij} \in S, \forall i, j\}$$

$$S_2 = \{A = [a_{ij}] \in M : A = -A^T \text{ and } a_{ij} \in S, \forall i, j\}$$

$$S_3 = \{A = [a_{ij}] \in M : a_{11} + a_{22} + a_{33} = 0 \text{ and } a_{ij} \in S, \forall i, j\}$$

If $n(S_1 \cup S_2 \cup S_3) = 125\alpha$, then α equals _____

Q5. Let $S = \{m \in \mathbf{Z} : A^{m^2} + A^m = 3I - A^{-6}\}$, where $A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$. Then $n(S)$ is equal to _____.

Q6. Let A be a square matrix of order 3 such that $\det(A) = -2$ and $\det(3 \text{adj}(-6 \text{adj}(3A))) = 2^{m+n} \cdot 3^{mn}$, $m > n$. Then $4m + 2n$ is equal to _____

Matrices

JEE Main 2025 January

Q7.

Let A be a 3×3 matrix such that $X^T A X = O$ for all nonzero 3×1 matrices $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$. If

$$A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \\ -5 \end{bmatrix}, A \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 4 \\ -8 \end{bmatrix}, \text{ and } \det(\text{adj}(2(A + I))) = 2^\alpha 3^\beta 5^\gamma, \alpha, \beta, \gamma \in N, \text{ then } \alpha^2 + \beta^2 + \gamma^2$$

is _____.

Q8.

$$\text{Let } A = [a_{ij}] = \begin{bmatrix} \log_5 128 & \log_4 5 \\ \log_5 8 & \log_4 25 \end{bmatrix}.$$

If A_{ij} is the cofactor of a_{ij} , $C_{ij} = \sum_{k=1}^2 a_{ik} A_{jk}$, $1 \leq i, j \leq 2$, and $C = [C_{ij}]$, then $8|C|$ is equal to :

(1) 288

(2) 222

(3) 242

(4) 262

Q9. If A, B , and $(\text{adj}(A^{-1}) + \text{adj}(B^{-1}))$ are non-singular matrices of same order, then the inverse of

$A(\text{adj}(A^{-1}) + \text{adj}(B^{-1}))^{-1} B$, is equal to

(1) $AB^{-1} + A^{-1} B$

(2) $\text{adj}(B^{-1}) + \text{adj}(A^{-1})$

(3) $\frac{AB^{-1}}{|A|} + \frac{BA^{-1}}{|B|}$

(4) $\frac{1}{|AB|}(\text{adj}(B) + \text{adj}(A))$

Q10. Let $A = \begin{bmatrix} \frac{1}{\sqrt{2}} & -2 \\ 0 & 1 \end{bmatrix}$ and $P = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$, $\theta > 0$. If $B = PAP^T$, $C = P^T B^{10} P$ and the sum of the diagonal elements of C is $\frac{m}{n}$, where $\gcd(m, n) = 1$, then $m + n$ is :

(1) 127

(2) 258

(3) 65

(4) 2049

Determinants

JEE Main 2025 January

Q1. Let M and m respectively be the maximum and the minimum values of

$$f(x) = \begin{vmatrix} 1 + \sin^2 x & \cos^2 x & 4 \sin 4x \\ \sin^2 x & 1 + \cos^2 x & 4 \sin 4x \\ \sin^2 x & \cos^2 x & 1 + 4 \sin 4x \end{vmatrix}, x \in \mathbf{R}$$

Then $M^4 - m^4$ is equal to :

- (1) 1280
- (2) 1295
- (3) 1215
- (4) 1040

Q2. The system of equations

$$x + y + z = 6$$

$$x + 2y + 5z = 9,$$

$$x + 5y + \lambda z = \mu,$$

has no solution if

- (1) $\lambda = 15, \mu \neq 17$
- (2) $\lambda \neq 17, \mu \neq 18$
- (3) $\lambda = 17, \mu \neq 18$
- (4) $\lambda = 17, \mu = 18$

Q3. For some a, b , let $f(x) = \begin{vmatrix} a + \frac{\sin x}{x} & 1 & b \\ a & 1 + \frac{\sin x}{x} & b \\ a & 1 & b + \frac{\sin x}{x} \end{vmatrix}, x \neq 0, \lim_{x \rightarrow 0} f(x) = \lambda + \mu a + \nu b$. Then $(\lambda + \mu + \nu)^2$ is equal to :

- (1) 16
- (2) 25
- (3) 9
- (4) 36

Q4. If the system of linear equations :

$$x + y + 2z = 6$$

$$2x + 3y + az = a + 1$$

$$-x - 3y + bz = 2b$$

where $a, b \in \mathbf{R}$, has infinitely many solutions, then $7a + 3b$ is equal to :

- (1) 16

Determinants

JEE Main 2025 January

(2) 12

(3) 22

(4) 9

Q5. If the system of equations

$$(\lambda - 1)x + (\lambda - 4)y + \lambda z = 5$$

$$\lambda x + (\lambda - 1)y + (\lambda - 4)z = 7$$

$$(\lambda + 1)x + (\lambda + 2)y - (\lambda + 2)z = 9$$

has infinitely many solutions, then $\lambda^2 + \lambda$ is equal to

(1) 6

(2) 10

(3) 20

(4) 12

Q6. If the system of equations

$$2x - y + z = 4$$

$$5x + \lambda y + 3z = 12$$

$$100x - 47y + \mu z = 212$$

has infinitely many solutions, then $\mu - 2\lambda$ is equal to

(1) 57

(2) 59

(3) 55

(4) 56

Q7. If the system of equations

$$x + 2y - 3z = 2$$

$$2x + \lambda y + 5z = 5$$

$$14x + 3y + \mu z = 33$$

has infinitely many solutions, then $\lambda + \mu$ is equal to :

(1) 13

(2) 10

(3) 12

(4) 11



Determinants

JEE Main 2025 January

Q8. Let $\alpha, \beta (\alpha \neq \beta)$ be the values of m , for which the equations $x + y + z = 1$; $x + 2y + 4z = m$ and $x + 4y + 10z = m^2$ have infinitely many solutions. Then the value of $\sum_{n=1}^{10} (n^\alpha + n^\beta)$ is equal to :

- (1) 3080
- (2) 560
- (3) 3410
- (4) 440



Probability

JEE Main 2025 January

Q1. Let $A = [a_{ij}]$ be a square matrix of order 2 with entries either 0 or 1. Let E be the event that A is an invertible matrix. Then the probability $P(E)$ is :

(1) $\frac{3}{16}$

(2) $\frac{5}{8}$

(3) $\frac{3}{8}$

(4) $\frac{1}{8}$

Q2. Two number k_1 and k_2 are randomly chosen from the set of natural numbers. Then, the probability that the value of $i^{k_1} + i^{k_2}$, ($i = \sqrt{-1}$) is non-zero, equals

(1) $\frac{1}{2}$

(2) $\frac{3}{4}$

(3) $\frac{1}{4}$

(4) $\frac{2}{3}$

Q3. Let S be the set of all the words that can be formed by arranging all the letters of the word GARDEN. From the set S , one word is selected at random. The probability that the selected word will NOT have vowels in alphabetical order is :

(1) $\frac{1}{2}$

(2) $\frac{1}{4}$

(3) $\frac{2}{3}$

(4) $\frac{1}{3}$

Q4. Two balls are selected at random one by one without replacement from a bag containing 4 white and 6 black balls. If the probability that the first selected ball is black, given that the second selected ball is also black, is $\frac{m}{n}$, where $\gcd(m, n) = 1$, then $m + n$ is equal to :

(1) 4

(2) 14

(3) 13

(4) 11

Probability

JEE Main 2025 January


Q5. A coin is tossed three times. Let X denote the number of times a tail follows a head. If μ and σ^2 denote the mean and variance of X , then the value of $64(\mu + \sigma^2)$ is :

- (1) 51
- (2) 64
- (3) 32
- (4) 48

Q6. One die has two faces marked 1 , two faces marked 2 , one face marked 3 and one face marked 4 . Another die has one face marked 1 , two faces marked 2 , two faces marked 3 and one face marked 4. The probability of getting the sum of numbers to be 4 or 5 , when both the dice are thrown together, is

- (1) $\frac{2}{3}$
- (2) $\frac{1}{2}$
- (3) $\frac{4}{9}$
- (4) $\frac{3}{5}$

Q7. A board has 16 squares as shown in the figure:



Out of these 16 squares, two squares are chosen at random. The probability that they have no side in common is :

- (1) $\frac{7}{10}$
- (2) $\frac{4}{5}$
- (3) $\frac{23}{30}$
- (4) $\frac{3}{5}$

Q8. A and B alternately throw a pair of dice. A wins if he throws a sum of 5 before B throws a sum of 8 , and B wins if he throws a sum of 8 before A throws a sum of 5 . The probability, that A wins if A makes the first throw, is

- (1) $\frac{8}{17}$
- (2) $\frac{9}{19}$
- (3) $\frac{9}{17}$

Probability

JEE Main 2025 January

(4) $\frac{8}{19}$

Q9. If A and B are two events such that $P(A \cap B) = 0.1$, and $P(A | B)$ and $P(B | A)$ are the roots of the equation $12x^2 - 7x + 1 = 0$, then the value of $\frac{P(\overline{A \cup B})}{P(\overline{A \cap B})}$ is :

(1) $\frac{4}{3}$

(2) $\frac{7}{4}$

(3) $\frac{5}{3}$

(4) $\frac{9}{4}$

Q10. Bag 1 contains 4 white balls and 5 black balls, and Bag 2 contains n white balls and 3 black balls. One ball is drawn randomly from Bag 1 and transferred to Bag 2. A ball is then drawn randomly from Bag 2. If the probability, that the ball drawn is white, is $29/45$, then n is equal to :

(1) 6

(2) 3

(3) 5

(4) 4

Q11. Bag B_1 contains 6 white and 4 blue balls, Bag B_2 contains 4 white and 6 blue balls, and Bag B_3 contains 5 white and 5 blue balls. One of the bags is selected at random and a ball is drawn from it. If the ball is white, then the probability, that the ball is drawn from Bag B_2 , is :

(1) $\frac{4}{15}$

(2) $\frac{1}{3}$

(3) $\frac{2}{5}$

(4) $\frac{2}{3}$

Q12. Let $A = [a_{ij}]$ be a 2×2 matrix such that $a_{ij} \in \{0, 1\}$ for all i and j . Let the random variable X denote the possible values of the determinant of the matrix A . Then, the variance of X is :

(1) $\frac{3}{4}$

(2) $\frac{5}{8}$

(3) $\frac{3}{8}$

(4) $\frac{1}{4}$

Probability

JEE Main 2025 January

Q13. Three defective oranges are accidentally mixed with seven good ones and on looking at them, it is not possible to differentiate between them. Two oranges are drawn at random from the lot. If x denote the number of defective oranges, then the variance of x is

- (1) $28/75$
- (2) $18/25$
- (3) $26/75$
- (4) $14/25$



Sets and Relations

JEE Main 2025 January

Q1. The number of non-empty equivalence relations on the set $\{1, 2, 3\}$ is :

- (1) 6
- (2) 5
- (3) 7
- (4) 4

Q2. Let $R = \{(1, 2), (2, 3), (3, 3)\}$ be a relation defined on the set $\{1, 2, 3, 4\}$. Then the minimum number of elements, needed to be added in R so that R becomes an equivalence relation, is:

- (1) 10
- (2) 7
- (3) 8
- (4) 9

Q3. Define a relation R on the interval $\left[0, \frac{\pi}{2}\right)$ by xRy if and only if $\sec^2 x - \tan^2 y = 1$. Then R is :

- (1) both reflexive and transitive but not symmetric
- (2) an equivalence relation
- (3) reflexive but neither symmetric nor transitive
- (4) both reflexive and symmetric but not transitive

Q4. Let $A = \{1, 2, 3, \dots, 10\}$ and $B = \left\{\frac{m}{n} : m, n \in A, m < n \text{ and } \gcd(m, n) = 1\right\}$. Then $n(B)$ is equal to :

- (1) 36
- (2) 31
- (3) 37
- (4) 29

Q5. Let $A = \{(x, y) \in \mathbf{R} \times \mathbf{R} : |x + y| \geq 3\}$ and $B = \{(x, y) \in \mathbf{R} \times \mathbf{R} : |x| + |y| \leq 3\}$.

If $C = \{(x, y) \in A \cap B : x = 0 \text{ or } y = 0\}$, then $\sum_{(x,y) \in C} |x + y|$ is :

- (1) 15
- (2) 24
- (3) 18

Sets and Relations

JEE Main 2025 January

(4) 12

Q6. Let $S = \{p_1, p_2, \dots, p_{10}\}$ be the set of first ten prime numbers. Let $A = S \cup P$, where P is the set of all possible products of distinct elements of S . Then the number of all ordered pairs (x, y) , $x \in S$, $y \in A$, such that x divides y , is _____.

Q7. Let $A = \left\{x \in (0, \pi) - \left\{\frac{\pi}{2}\right\} : \log_{(2/\pi)} |\sin x| + \log_{(2/\pi)} |\cos x| = 2\right\}$ and $B = \{x \geq 0 : \sqrt{x}(\sqrt{x} - 4) - 3|\sqrt{x} - 2| + 6 = 0\}$. Then $n(A \cup B)$ is equal to :

(1) 4

(2) 8

(3) 6

(4) 2

Q8. Let $S = \mathbf{N} \cup \{0\}$. Define a relation R from S to $\mathbf{R}$ by :

$$R = \left\{(x, y) : \log_e y = x \log_e \left(\frac{2}{5}\right), x \in S, y \in \mathbf{R}\right\}$$

Then, the sum of all the elements in the range of R is equal to :

(1) $\frac{10}{9}$

(2) $\frac{3}{2}$

(3) $\frac{5}{2}$

(4) $\frac{5}{3}$

Q9. Let $A = \{1, 2, 3\}$. The number of relations on A , containing $(1, 2)$ and $(2, 3)$, which are reflexive and transitive but not symmetric, is _____ -

Q10. Let $X = \mathbf{R} \times \mathbf{R}$. Define a relation R on X as :

$$(a_1, b_1) R (a_2, b_2) \Leftrightarrow b_1 = b_2$$

Statement I : R is an equivalence relation.

Statement II : For some $(a, b) \in X$, the set $S = \{(x, y) \in X : (x, y)R(a, b)\}$ represents a line parallel to $y = x$.

In the light of the above statements, choose the correct answer from the options given below :

(1) Both Statement I and Statement II are false

(2) Statement I is true but Statement II is false

(3) Both Statement I and Statement II are true

(4) Statement I is false but Statement II is true

Sets and Relations

JEE Main 2025 January

Q11. The relation $R = \{(x, y) : x, y \in \mathbb{Z} \text{ and } x + y \text{ is even} \}$ is:

- (1) reflexive and symmetric but not transitive
- (2) an equivalence relation
- (3) symmetric and transitive but not reflexive
- (4) reflexive and transitive but not symmetric



Functions

JEE Main 2025 January

Q1. Let $f : \mathbf{R} - \{0\} \rightarrow (-\infty, 1)$ be a polynomial of degree 2, satisfying $f(x)f\left(\frac{1}{x}\right) = f(x) + f\left(\frac{1}{x}\right)$. If

$f(K) = -2K$, then the sum of squares of all possible values of K is :

- (1) 7
- (2) 6
- (3) 1
- (4) 9

Q2. Number of functions $f : \{1, 2, \dots, 100\} \rightarrow \{0, 1\}$, that assign 1 to exactly one of the positive integers less than or equal to 98, is equal to _____.

Q3. Let $[x]$ denote the greatest integer less than or equal to x . Then the domain of $f(x) = \sec^{-1}(2[x] + 1)$ is :

- (1) $(-\infty, -1] \cup [0, \infty)$
- (2) $(-\infty, -1] \cup [1, \infty)$
- (3) $(-\infty, \infty)$
- (4) $(-\infty, \infty) - \{0\}$ ★

Q4. If the domain of the function $\log_5(18x - x^2 - 77)$ is (α, β) and the domain of the function $\log_{(x-1)}\left(\frac{2x^2+3x-2}{x^2-3x-4}\right)$ is (γ, δ) , then $\alpha^2 + \beta^2 + \gamma^2$ is equal to :

- (1) 195
- (2) 179
- (3) 186
- (4) 174

Q5. Let the range of the function $f(x) = 6 + 16 \cos x \cdot \cos\left(\frac{\pi}{3} - x\right) \cdot \cos\left(\frac{\pi}{3} + x\right) \cdot \sin 3x \cdot \cos 6x$, $x \in \mathbf{R}$ be $[\alpha, \beta]$.

Then the distance of the point (α, β) from the line $3x + 4y + 12 = 0$ is :

- (1) 11
- (2) 8
- (3) 10
- (4) 9

Q6. Let $f(x) = \log_e x$ and $g(x) = \frac{x^4 - 2x^3 + 3x^2 - 2x + 2}{2x^2 - 2x + 1}$. Then the domain of $f \circ g$ is

Functions

JEE Main 2025 January

- (1) $[0, \infty)$
- (2) $[1, \infty)$
- (3) $(0, \infty)$
- (4) $\mathbb{R}$

Q7. Let $A = \{1, 2, 3, 4\}$ and $B = \{1, 4, 9, 16\}$. Then the number of many-one functions $f : A \rightarrow B$ such that $1 \in f(A)$ is equal to :

- (1) 151
- (2) 139
- (3) 163
- (4) 127

Q8. Let $f : [0, 3] \rightarrow A$ be defined by $f(x) = 2x^3 - 15x^2 + 36x + 7$ and $g : [0, \infty) \rightarrow B$ be defined by $g(x) = \frac{x^{2025}}{x^{2025} + 1}$. If both the functions are onto and $S = \{x \in \mathbb{Z} : x \in A \text{ or } x \in B\}$, then $n(S)$ is equal to :

- (1) 29
- (2) 30
- (3) 31
- (4) 36

Q9. Let $f(x) = \frac{2^{x+2} + 16}{2^{2x+1} + 2^{x+4} + 32}$. Then the value of $8 \left(f\left(\frac{1}{15}\right) + f\left(\frac{2}{15}\right) + \dots + f\left(\frac{59}{15}\right) \right)$ is equal to

- (1) 92
- (2) 118
- (3) 102
- (4) 108

Q10. The number of real solution(s) of the equation $x^2 + 3x + 2 = \min\{|x - 3|, |x + 2|\}$ is:

- (1) 1
- (2) 0
- (3) 2
- (4) 3

Functions

JEE Main 2025 January

Q11. Let $f : \mathbb{R} - \{0\} \rightarrow \mathbb{R}$ be a function such that $f(x) - 6f\left(\frac{1}{x}\right) = \frac{35}{3x} - \frac{5}{2}$.

If the $\lim_{x \rightarrow 0} \left(\frac{1}{\alpha x} + f(x) \right) = \beta$; $\alpha, \beta \in \mathbb{R}$, then $\alpha + 2\beta$ is equal to

- (1) 5
- (2) 3
- (3) 4
- (4) 6

Q12. If $f(x) = \frac{2^x}{2^x + \sqrt{2}}$, $x \in \mathbb{R}$, then $\sum_{k=1}^{81} f\left(\frac{k}{82}\right)$ is equal to

- (1) $81\sqrt{2}$
- (2) 41
- (3) 82
- (4) $\frac{81}{2}$

Q13. The function $f : (-\infty, \infty) \rightarrow (-\infty, 1)$, defined by $f(x) = \frac{2^x - 2^{-x}}{2^x + 2^{-x}}$ is:

- (1) Neither one-one nor onto
- (2) Onto but not one-one
- (3) Both one-one and onto
- (4) One-one but not onto

Q14. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by

$$f(x) = (2 + 3a)x^2 + \left(\frac{a+2}{a-1}\right)x + b, a \neq 1. \text{ If}$$

$$f(x+y) = f(x) + f(y) + 1 - \frac{2}{7}xy, \text{ then the value of } 28 \sum_{i=1}^5 |f(i)| \text{ is}$$

- (1) 545
- (2) 715
- (3) 735
- (4) 675

Limits

JEE Main 2025 January

Q1. $\lim_{x \rightarrow 0} \operatorname{cosec} x \left(\sqrt{2 \cos^2 x + 3 \cos x} - \sqrt{\cos^2 x + \sin x + 4} \right)$ is:

- (1) 0
- (2) $\frac{1}{\sqrt{15}}$
- (3) $\frac{1}{2\sqrt{5}}$
- (4) $-\frac{1}{2\sqrt{5}}$

Q2. If $\lim_{x \rightarrow \infty} \left(\left(\frac{e}{1-e} \right) \left(\frac{1}{e} - \frac{x}{1+x} \right) \right)^x = \alpha$, then the value of $\frac{\log_e \alpha}{1 + \log_e \alpha}$ equals :

- (1) e^{-1}
- (2) e^2
- (3) e^{-2}
- (4) e

Q3. If $\lim_{t \rightarrow 0} \left(\int_0^1 (3x+5)^t dx \right)^{\frac{1}{t}} = \frac{\alpha}{5e} \left(\frac{8}{5} \right)^{\frac{2}{3}}$, then α is equal to _____

Q4. Let $[t]$ be the greatest integer less than or equal to t . Then the least value of $p \in \mathbf{N}$ for which $\lim_{x \rightarrow 0^+} \left(x \left(\left[\frac{1}{x} \right] + \left[\frac{2}{x} \right] + \dots + \left[\frac{p}{x} \right] \right) - x^2 \left(\left[\frac{1}{x^2} \right] + \left[\frac{2^2}{x^2} \right] + \dots + \left[\frac{p^2}{x^2} \right] \right) \right) \geq 1$ is equal to _____.

Q5. $\lim_{x \rightarrow \infty} \frac{(2x^2 - 3x + 5)(3x - 1)^{\frac{x}{2}}}{(3x^2 + 5x + 4)\sqrt{(3x + 2)^x}}$ is equal to :

- (1) $\frac{2}{\sqrt{3e}}$
- (2) $\frac{2e}{\sqrt{3}}$
- (3) $\frac{2}{3\sqrt{e}}$
- (4) $\frac{2e}{3}$

Q6. Let $f(x) = \lim_{n \rightarrow \infty} \sum_{r=0}^n \left(\frac{\tan(x/2^{r+1}) + \tan^3(x/2^{r+1})}{1 - \tan^2(x/2^{r+1})} \right)$. Then $\lim_{x \rightarrow 0} \frac{e^x - e^{f(x)}}{(x - f(x))}$ is equal to

Continuity and Differentiability

JEE Main 2025 January

Q1. If the function

$$f(x) = \begin{cases} \frac{2}{x} \{ \sin(k_1 + 1)x + \sin(k_2 - 1)x \}, & x < 0 \\ 4, & x = 0 \\ \frac{2}{x} \log_e \left(\frac{2+k_1x}{2+k_2x} \right), & x > 0 \end{cases}$$

is continuous at $x = 0$, then $k_1^2 + k_2^2$ is equal to

- (1) 20
- (2) 5
- (3) 8
- (4) 10

Q2. Let $f(x)$ be a real differentiable function such that $f(0) = 1$ and $f(x+y) = f(x)f'(y) + f'(x)f(y)$ for all $x, y \in \mathbf{R}$. Then $\sum_{n=1}^{100} \log_e f(n)$ is equal to :

- (1) 2525
- (2) 5220
- (3) 2384
- (4) 2406

Q3. Let $[x]$ denote the greatest integer function, and let m and n respectively be the numbers of the points, where the function $f(x) = [x] + |x - 2|$, $-2 < x < 3$, is not continuous and not differentiable. Then $m + n$ is equal to :

- (1) 6
- (2) 8
- (3) 9
- (4) 7

Q4. Let $f(x) = \begin{cases} 3x, & x < 0 \\ \min\{1 + x + [x], x + 2[x]\}, & 0 \leq x \leq 2 \\ 5, & x > 2, \end{cases}$

where $[.]$ denotes greatest integer function. If α and β are the number of points, where f is not continuous and is not differentiable, respectively, then $\alpha + \beta$ equals _____

Q5. Let the function $f(x) = (x^2 + 1) |x^2 - ax + 2| + \cos |x|$ be not differentiable at the two points $x = \alpha = 2$ and $x = \beta$. Then the distance of the point (α, β) from the line $12x + 5y + 10 = 0$ is equal to :

- (1) 5

Continuity and Differentiability

JEE Main 2025 January

(2) 4

(3) 3

(4) 2

Q6. Let the function,

$$f(x) = \begin{cases} -3ax^2 - 2, & x < 1 \\ a^2 + bx, & x \geq 1 \end{cases}$$

be differentiable for all $x \in \mathbf{R}$, where $\mathbf{a} > 1, \mathbf{b} \in \mathbf{R}$. If the area of the region enclosed by $y = f(x)$ and the line $y = -20$ is $\alpha + \beta\sqrt{3}$, $\alpha, \beta \in \mathbf{Z}$, then the value of $\alpha + \beta$ is _____



Application of Derivatives

JEE Main 2025 January

Q1. A spherical chocolate ball has a layer of ice-cream of uniform thickness around it. When the thickness of the ice-cream layer is 1 cm, the ice-cream melts at the rate of $81 \text{ cm}^3/\text{min}$ and the thickness of the ice-cream layer decreases at the rate of $\frac{1}{4\pi} \text{ cm/min}$. The surface area (in cm^2) of the chocolate ball (without the ice-cream layer) is :

- (1) 196π
- (2) 256π
- (3) 225π
- (4) 128π

Q2. Let $(2, 3)$ be the largest open interval in which the function $f(x) = 2 \log_e(x - 2) - x^2 + ax + 1$ is strictly increasing and (b, c) be the largest open interval, in which the function $g(x) = (x - 1)^3(x + 2 - a)^2$ is strictly decreasing. Then $100(a + b - c)$ is equal to :

- (1) 420
- (2) 360
- (3) 160
- (4) 280

Q3. If the set of all values of a , for which the equation $5x^3 - 15x - a = 0$ has three distinct real roots, is the interval (α, β) , then $\beta - 2\alpha$ is equal to _____

Q4. The sum of all local minimum values of the function

$$f(x) = \begin{cases} 1 - 2x, & x < -1 \\ \frac{1}{3}(7 + 2|x|), & -1 \leq x \leq 2 \\ \frac{11}{18}(x - 4)(x - 5), & x > 2 \end{cases}$$

is

- (1) $\frac{157}{72}$
- (2) $\frac{131}{72}$
- (3) $\frac{171}{72}$
- (4) $\frac{167}{72}$

Indefinite Integration

JEE Main 2025 January

Q1. Let $I(x) = \int \frac{dx}{(x-11)^{\frac{11}{13}}(x+15)^{\frac{15}{13}}}$. If $I(37) - I(24) = \frac{1}{4} \left(\frac{1}{b^{\frac{1}{13}}} - \frac{1}{c^{\frac{1}{13}}} \right)$, $b, c \in \mathbb{N}$, then $3(b + c)$ is equal to

(1) 22

(2) 39

(3) 40

(4) 26

Q2. If $\int \frac{2x^2+5x+9}{\sqrt{x^2+x+1}} dx = x\sqrt{x^2+x+1} + \alpha\sqrt{x^2+x+1} + \beta \log_e \left| x + \frac{1}{2} + \sqrt{x^2+x+1} \right| + C$, where C is the constant of integration, then $\alpha + 2\beta$ is equal to _____.

Q3. If $\int e^x \left(\frac{x \sin^{-1} x}{\sqrt{1-x^2}} + \frac{\sin^{-1} x}{(1-x^2)^{3/2}} + \frac{x}{1-x^2} \right) dx = g(x) + C$, where C is the constant of integration, then $g\left(\frac{1}{2}\right)$ equals :

(1) $\frac{\pi}{4} \sqrt{\frac{e}{3}}$

(2) $\frac{\pi}{6} \sqrt{\frac{e}{3}}$

(3) $\frac{\pi}{4} \sqrt{\frac{e}{2}}$

(4) $\frac{\pi}{6} \sqrt{\frac{e}{2}}$

Q4. Let $\int x^3 \sin x dx = g(x) + C$, where C is the constant of integration. If $8 \left(g\left(\frac{\pi}{2}\right) + g'\left(\frac{\pi}{2}\right) \right) = \alpha\pi^3 + \beta\pi^2 + \gamma$, $\alpha, \beta, \gamma \in \mathbb{Z}$, then $\alpha + \beta - \gamma$ equals :

(1) 48

(2) 55

(3) 62

(4) 47

Q5. If $f(x) = \int \frac{1}{x^{1/4}(1+x^{1/4})} dx$, $f(0) = -6$, then $f(1)$ is equal to :

(1) $4(\log_e 2 - 2)$

(2) $2 - \log_{e^2} 2$

(3) $\log_e 2 + 2$

(4) $4(\log_e 2 + 2)$

Definite Integration

JEE Main 2025 January

Q1. The integral $80 \int_0^{\frac{\pi}{4}} \left(\frac{\sin \theta + \cos \theta}{9 + 16 \sin 2\theta} \right) d\theta$ is equal to :

- (1) $3 \log_e 4$
- (2) $4 \log_e 3$
- (3) $6 \log_e 4$
- (4) $2 \log_e 3$

Q2. Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be a twice differentiable function such that $f(2) = 1$. If $F(x) = xf(x)$ for all $x \in \mathbf{R}$, $\int_0^2 x F'(x) dx = 6$ and $\int_0^2 x^2 F''(x) dx = 40$, then $F'(2) + \int_0^2 F(x) dx$ is equal to :

- (1) 11
- (2) 13
- (3) 15
- (4) 9

Q3. If $24 \int_0^{\frac{\pi}{4}} \left(\sin \left| 4x - \frac{\pi}{12} \right| + [2 \sin x] \right) dx = 2\pi + \alpha$, where $[\cdot]$ denotes the greatest integer function, then α is equal to _____.

Q4. Let $f(x) = \int_0^{x^2} \frac{t^2 - 8t + 15}{e^t} dt$, $x \in \mathbf{R}$. Then the numbers of local maximum and local minimum points of f , respectively, are :

- (1) 2 and 3
- (2) 2 and 2
- (3) 3 and 2
- (4) 1 and 3

Q5. Let for some function $y = f(x)$, $\int_0^x t f(t) dt = x^2 f(x)$, $x > 0$ and $f(2) = 3$. Then $f(6)$ is equal to

- (1) 1
- (2) 3
- (3) 6
- (4) 2

Q6. Let f be a real valued continuous function defined on the positive real axis such that $g(x) = \int_0^x t f(t) dt$. If $g(x^3) = x^6 + x^7$, then value of $\sum_{r=1}^{15} f(r^3)$ is :

Definite Integration

JEE Main 2025 January

- (1) 270
- (2) 340
- (3) 320
- (4) 310

Q7. Let $f(x) = \int_0^x t(t^2 - 9t + 20) dt$, $1 \leq x \leq 5$. If the range of f is $[\alpha, \beta]$, then $4(\alpha + \beta)$ equals :

- (1) 253
- (2) 154
- (3) 125
- (4) 157

Q8. If $I(m, n) = \int_0^1 x^{m-1}(1-x)^{n-1} dx$, $m, n > 0$, then $I(9, 14) + I(10, 13)$ is

- (1) $I(19, 27)$
- (2) $I(9, 1)$
- (3) $I(1, 13)$
- (4) $I(9, 13)$

Q9. Let for $f(x) = 7 \tan^8 x + 7 \tan^6 x - 3 \tan^4 x - 3 \tan^2 x$, $I_1 = \int_0^{\pi/4} f(x) dx$ and $I_2 = \int_0^{\pi/4} x f(x) dx$. Then $7I_1 + 12I_2$ is equal to :

- (1) 2
- (2) 1
- (3) 2π
- (4) π

Q10. The value of

$$\int_{e^2}^{e^4} \frac{1}{x} \left(\frac{e^{((\log_e x)^2 + 1)^{-1}}}{e^{((\log_e x)^2 + 1)^{-1}} + e^{((6 - \log_e x)^2 + 1)^{-1}}} \right) dx \text{ is}$$

- (1) 2
- (2) $\log_e 2$
- (3) 1

Definite Integration

JEE Main 2025 January

(4) e^2

Q11. If $I = \int_0^{\frac{\pi}{2}} \frac{\sin^{\frac{3}{2}} x}{\sin^{\frac{3}{2}} x + \cos^{\frac{3}{2}} x} dx$, then $\int_0^{21} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} dx$ equals :

(1) $\frac{\pi^2}{12}$

(2) $\frac{\pi^2}{4}$

(3) $\frac{\pi^2}{16}$

(4) $\frac{\pi^2}{8}$

Q12. If $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{96x^2 \cos^2 x}{(1+e^x)} dx = \pi(\alpha\pi^2 + \beta)$, $\alpha, \beta \in \mathbb{Z}$, then $(\alpha + \beta)^2$ equals

(1) 64

(2) 196

(3) 144

(4) 100

Q13. Let $f : (0, \infty) \rightarrow \mathbf{R}$ be a twice differentiable function. If for some $a' = 0$, $\int_0^1 f(\lambda x) d\lambda = af(x)$, $f(1) = 1$ and $f(16) = \frac{1}{8}$, then $16 - f'\left(\frac{1}{16}\right)$ is equal to _____.

Area Under Curves

JEE Main 2025 January

Q1. The area of the region, inside the circle $(x - 2\sqrt{3})^2 + y^2 = 12$ and outside the parabola $y^2 = 2\sqrt{3}x$ is :

- (1) $3\pi + 8$
- (2) $6\pi - 16$
- (3) $3\pi - 8$
- (4) $6\pi - 8$

Q2. The area of the region enclosed by the curves $y = x^2 - 4x + 4$ and $y^2 = 16 - 8x$ is :

- (1) $\frac{8}{3}$
- (2) $\frac{4}{3}$
- (3) 8
- (4) 5

Q3. The area (in sq. units) of the region

$$\{(x, y) : 0 \leq y \leq 2|x| + 1, 0 \leq y \leq x^2 + 1, |x| \leq 3\}$$


- (1) $\frac{80}{3}$
- (2) $\frac{64}{3}$
- (3) $\frac{32}{3}$
- (4) $\frac{17}{3}$

Q4. The area of the region bounded by the curves $x(1 + y^2) = 1$ and $y^2 = 2x$ is:

- (1) $2\left(\frac{\pi}{2} - \frac{1}{3}\right)$
- (2) $\frac{\pi}{2} - \frac{1}{3}$
- (3) $\frac{\pi}{4} - \frac{1}{3}$
- (4) $\frac{1}{2}\left(\frac{\pi}{2} - \frac{1}{3}\right)$

Q5. The area of the region enclosed by the curves $y = e^x$, $y = |e^x - 1|$ and y -axis is:

- (1) $1 - \log_e 2$
- (2) $\log_e 2$
- (3) $1 + \log_e 2$



Area Under Curves

JEE Main 2025 January

(4) $2 \log_e 2 - 1$

Q6. If the area of the larger portion bounded between the curves $x^2 + y^2 = 25$ and $y = |x - 1|$ is $\frac{1}{4}(b\pi + c)$, $b, c \in N$, then $b + c$ is equal to

Q7. The area of the region $\{(x, y) : x^2 + 4x + 2 \leq y \leq |x + 2|\}$ is equal to

(1) 7

(2) 5

(3) $24/5$

(4) $20/3$

Q8. Let the area enclosed between the curves $|y| = 1 - x^2$ and $x^2 + y^2 = 1$ be α . If $9\alpha = \beta\pi + \gamma$; β, γ are integers, then the value of $|\beta - \gamma|$ equals.

(1) 27

(2) 33

(3) 15

(4) 18

Q9. If the area of the region $\{(x, y) : -1 \leq x \leq 1, 0 \leq y \leq a + e^{|x|} - e^{-x}, a > 0\}$ is $\frac{e^2 + 8e + 1}{e}$, then the value of a is :

(1) 8

(2) 7

(3) 5

(4) 6


Q10. Let the area of the region $\{(x, y) : 2y \leq x^2 + 3, y + |x| \leq 3, y \geq |x - 1|\}$ be A . Then $6A$ is equal to :

(1) 16

(2) 12

(3) 14

(4) 18



Area Under Curves

JEE Main 2025 January

Q11. Consider the region $R = \left\{ (x, y) : x \leq y \leq 9 - \frac{11}{3}x^2, x \geq 0 \right\}$.

The area, of the largest rectangle of sides parallel to the coordinate axes and inscribed in R , is:

(1) $\frac{730}{119}$

(2) $\frac{625}{111}$

(3) $\frac{821}{123}$

(4) $\frac{567}{121}$


Q12. Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be a twice differentiable function such that $f(x + y) = f(x)f(y)$ for all $x, y \in \mathbf{R}$. If $f'(0) = 4a$ and f satisfies $f''(x) - 3af'(x) - f(x) = 0$, $a > 0$, then the area of the region $R = \{(x, y) \mid 0 \leq y \leq f(ax), 0 \leq x \leq 2\}$ is:

(1) $e^2 - 1$

(2) $e^2 + 1$

(3) $e^4 + 1$

(4) $e^4 - 1$



Differential Equations

JEE Main 2025 January

Q1. Let $y = y(x)$ be the solution of the differential equation $2 \cos x \frac{dy}{dx} = \sin 2x - 4y \sin x$, $x \in \left(0, \frac{\pi}{2}\right)$. If $y\left(\frac{\pi}{3}\right) = 0$, then $y'\left(\frac{\pi}{4}\right) + y\left(\frac{\pi}{4}\right)$ is equal to _____.

Q2. Let $x = x(y)$ be the solution of the differential equation $y = \left(x - y \frac{dx}{dy}\right) \sin\left(\frac{x}{y}\right)$, $y > 0$ and $x(1) = \frac{\pi}{2}$. Then $\cos(x(2))$ is equal to :

(1) $1 - 2(\log_e 2)^2$

(2) $1 - 2(\log_e 2)$

(3) $2(\log_e 2) - 1$

(4) $2(\log_e 2)^2 - 1$

Q3. Let $x = x(y)$ be the solution of the differential equation $y^2 dx + \left(x - \frac{1}{y}\right) dy = 0$. If $x(1) = 1$, then $x\left(\frac{1}{2}\right)$ is :

(1) $\frac{1}{2} + e$

(2) $3 + e$

(3) $3 - e$

(4) $\frac{3}{2} + e$

Q4. If $x = f(y)$ is the solution of the differential equation

$$(1 + y^2) + \left(x - 2e^{\tan^{-1} y}\right) \frac{dy}{dx} = 0, y \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

with $f(0) = 1$, then $f\left(\frac{1}{\sqrt{3}}\right)$ is equal to :

(1) $e^{\pi/12}$

(2) $e^{\pi/4}$

(3) $e^{\pi/3}$

(4) $e^{\pi/6}$


Q5. Let a curve $y = f(x)$ pass through the points $(0, 5)$ and $(\log_e 2, k)$. If the curve satisfies the differential equation $2(3 + y)e^{2x} dx - (7 + e^{2x}) dy = 0$, then k is equal to

(1) 4

(2) 32

(3) 8

(4) 16



Differential Equations

JEE Main 2025 January

Q6. Let $y = y(x)$ be the solution of the differential equation $(xy - 5x^2\sqrt{1+x^2})dx + (1+x^2)dy = 0, y(0) = 0$.

Then $y(\sqrt{3})$ is equal to

(1) $\sqrt{\frac{15}{2}}$

(2) $\frac{5\sqrt{3}}{2}$

(3) $2\sqrt{2}$

(4) $\sqrt{\frac{14}{3}}$

Q7. Let f be a differentiable function such that $2(x+2)^2 f(x) - 3(x+2)^2 = 10 \int_0^x (t+2)f(t)dt, x \geq 0$. Then $f(2)$ is equal to _____.

Q8. Let $f : (0, \infty) \rightarrow \mathbf{R}$ be a function which is differentiable at all points of its domain and satisfies the condition $x^2 f'(x) = 2x f(x) + 3$, with $f(1) = 4$. Then $2f(2)$ is equal to :

(1) 39

(2) 19

(3) 29

(4) 23

Q9. If $y = y(x)$ is the solution of the differential equation,

$$\sqrt{4-x^2} \frac{dy}{dx} = \left(\left(\sin^{-1} \left(\frac{x}{2} \right) \right)^2 - y \right) \sin^{-1} \left(\frac{x}{2} \right), -2 \leq x \leq 2, y(2) = \frac{\pi^2 - 8}{4}, \text{ then } y^2(0) \text{ is equal to}$$

Q10. Let $y = f(x)$ be the solution of the differential equation $\frac{dy}{dx} + \frac{xy}{x^2-1} = \frac{x^6+4x}{\sqrt{1-x^2}}, -1 < x < 1$ such that $f(0) = 0$.

If $6 \int_{-1/2}^{1/2} f(x)dx = 2\pi - \alpha$ then α^2 is equal to _____.

Q11. Let $y = y(x)$ be the solution of the differential equation $\cos x (\log_e (\cos x))^2 dy + (\sin x - 3y \sin x \log_e (\cos x)) dx = 0, x \in (0, \frac{\pi}{2})$. If $y(\frac{\pi}{4}) = \frac{-1}{\log_e 2}$, then $y(\frac{\pi}{6})$ is equal to :

(1) $\frac{1}{\log_e (3) - \log_e (4)}$

(2) $\frac{2}{\log_e (3) - \log_e (4)}$

(3) $\frac{1}{\log_e (4) - \log_e (3)}$

(4) $-\frac{1}{\log_e (4)}$

Differential Equations

JEE Main 2025 January

Q12. If for the solution curve $y = f(x)$ of the differential equation $\frac{dy}{dx} + (\tan x)y = \frac{2+\sec x}{(1+2\sec x)^2}$,

$x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, $f\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}}{10}$, then $f\left(\frac{\pi}{4}\right)$ is equal to :

(1) $\frac{\sqrt{3}+1}{10(4+\sqrt{3})}$

(2) $\frac{5-\sqrt{3}}{2\sqrt{2}}$

(3) $\frac{9\sqrt{3}+3}{10(4+\sqrt{3})}$

(4) $\frac{4-\sqrt{2}}{14}$



Straight Lines

JEE Main 2025 January

Q1. Let the area of a $\triangle PQR$ with vertices $P(5, 4)$, $Q(-2, 4)$ and $R(a, b)$ be 35 square units. If its orthocenter and centroid are $O\left(2, \frac{14}{5}\right)$ and $C(c, d)$ respectively, then $c + 2d$ is equal to

- (1) $\frac{8}{3}$
- (2) $\frac{7}{3}$
- (3) 2
- (4) 3

Q2. Let the triangle PQR be the image of the triangle with vertices $(1, 3)$, $(3, 1)$ and $(2, 4)$ in the line $x + 2y = 2$. If the centroid of $\triangle PQR$ is the point (α, β) , then $15(\alpha - \beta)$ is equal to :

- (1) 19
- (2) 24
- (3) 21
- (4) 22

Q3. Let $A(6, 8)$, $B(10 \cos \alpha, -10 \sin \alpha)$ and $C(-10 \sin \alpha, 10 \cos \alpha)$, be the vertices of a triangle. If $L(a, 9)$ and $G(h, k)$ be its orthocenter and centroid respectively, then $(5a - 3h + 6k + 100 \sin 2\alpha)$ is equal to _____.

Q4. Let ABC be a triangle formed by the lines $7x - 6y + 3 = 0$, $x + 2y - 31 = 0$ and $9x - 2y - 19 = 0$. Let the point (h, k) be the image of the centroid of $\triangle ABC$ in the line $3x + 6y - 53 = 0$. Then $h^2 + k^2 + hk$ is equal to:

- (1) 47
- (2) 37
- (3) 36
- (4) 40

Q5. Let the points $\left(\frac{11}{2}, \alpha\right)$ lie on or inside the triangle with sides $x + y = 11$, $x + 2y = 16$ and $2x + 3y = 29$. Then the product of the smallest and the largest values of α is equal to :

- (1) 44
- (2) 22
- (3) 33
- (4) 55

Straight Lines

JEE Main 2025 January

Q6. Two equal sides of an isosceles triangle are along $-x + 2y = 4$ and $x + y = 4$. If m is the slope of its third side, then the sum, of all possible distinct values of m , is :

- (1) $-2\sqrt{10}$
- (2) 12
- (3) 6
- (4) -6

Q7. Let the lines $3x - 4y - \alpha = 0$, $8x - 11y - 33 = 0$, and $2x - 3y + \lambda = 0$ be concurrent. If the image of the point $(1, 2)$ in the line $2x - 3y + \lambda = 0$ is $\left(\frac{57}{13}, \frac{-40}{13}\right)$, then $|\alpha\lambda|$ is equal to

- (1) 84
- (2) 113
- (3) 91
- (4) 101

Q8. Let ${}^nC_{r-1} = 28$, ${}^nC_r = 56$ and ${}^nC_{r+1} = 70$. Let $A(4 \cos t, 4 \sin t)$, $B(2 \sin t, -2 \cos t)$ and $C(3r - n, r^2 - n - 1)$ be the vertices of a triangle ABC , where t is a parameter. If $(3x - 1)^2 + (3y)^2 = \alpha$, is the locus of the centroid of triangle ABC , then α equals

- (1) 6
- (2) 18
- (3) 8
- (4) 20

Q9. Let the line $x + y = 1$ meet the axes of x and y at A and B , respectively. A right angled triangle AMN is inscribed in the triangle OAB , where O is the origin and the points M and N lie on the lines OB and AB , respectively. If the area of the triangle AMN is $\frac{4}{9}$ of the area of the triangle OAB and $AN : NB = \lambda : 1$, then the sum of all possible value(s) of λ is :

- (1) 2
- (2) $\frac{5}{2}$
- (3) $\frac{1}{2}$
- (4) $\frac{13}{6}$

Straight Lines

JEE Main 2025 January

Q10. A rod of length eight units moves such that its ends A and B always lie on the lines $x - y + 2 = 0$ and $y + 2 = 0$, respectively. If the locus of the point P , that divides the rod AB internally in the ratio $2 : 1$ is $9(x^2 + \alpha y^2 + \beta xy + \gamma x + 28y) - 76 = 0$, then $\alpha - \beta - \gamma$ is equal to :

(1) 22

(2) 21

(3) 23

(4) 24



Circle

JEE Main 2025 January

Q1. Let the equation of the circle, which touches x -axis at the point $(a, 0)$, $a > 0$ and cuts off an intercept of length b on y -axis be $x^2 + y^2 - \alpha x + \beta y + \gamma = 0$. If the circle lies below x -axis, then the ordered pair $(2a, b^2)$ is equal to

- (1) $(\gamma, \beta^2 - 4\alpha)$
- (2) $(\alpha, \beta^2 + 4\gamma)$
- (3) $(\gamma, \beta^2 + 4\alpha)$
- (4) $(\alpha, \beta^2 - 4\gamma)$

Q2. Let the line $x + y = 1$ meet the circle $x^2 + y^2 = 4$ at the points A and B. If the line perpendicular to AB and passing through the mid point of the chord AB intersects the circle at C and D , then the area of the quadrilateral ADBC is equal to :

- (1) $\sqrt{14}$
- (2) $3\sqrt{7}$
- (3) $2\sqrt{14}$
- (4) $5\sqrt{7}$

Q3. Let a circle C pass through the points $(4, 2)$ and $(0, 2)$, and its centre lie on $3x + 2y + 2 = 0$. Then the length of the chord, of the circle C , whose mid-point is $(1, 2)$, is :

- (1) $\sqrt{3}$
- (2) $2\sqrt{2}$
- (3) $2\sqrt{3}$
- (4) $4\sqrt{2}$

Q4. A circle C of radius 2 lies in the second quadrant and touches both the coordinate axes. Let r be the radius of a circle that has centre at the point $(2, 5)$ and intersects the circle C at exactly two points. If the set of all possible values of r is the interval (α, β) , then $3\beta - 2\alpha$ is equal to :

- (1) 10
- (2) 15
- (3) 12
- (4) 14

Q5. Let circle C be the image of $x^2 + y^2 - 2x + 4y - 4 = 0$ in the line $2x - 3y + 5 = 0$ and A be the point on C such that OA is parallel to x -axis and A lies on the right hand side of the centre O of C . If $B(\alpha, \beta)$, with $\beta < 4$,

Circle

JEE Main 2025 January

lies on C such that the length of the arc AB is $(1/6)^{\text{th}}$ of the perimeter of C , then $\beta - \sqrt{3}\alpha$ is equal to

(1) $3 + \sqrt{3}$

(2) 4

(3) $4 - \sqrt{3}$

(4) 3



Parabola

JEE Main 2025 January

- Q1.** The focus of the parabola $y^2 = 4x + 16$ is the centre of the circle C of radius 5. If the values of λ , for which C passes through the point of intersection of the lines $3x - y = 0$ and $x + \lambda y = 4$, are λ_1 and λ_2 , $\lambda_1 < \lambda_2$, then $12\lambda_1 + 29\lambda_2$ is equal to _____
- Q2.** If the equation of the parabola with vertex $V\left(\frac{3}{2}, 3\right)$ and the directrix $x + 2y = 0$ is $\alpha x^2 + \beta y^2 - \gamma xy - 30x - 60y + 225 = 0$, then $\alpha + \beta + \gamma$ is equal to :
- (1) 7
(2) 9
(3) 8
(4) 6
- Q3.** Two parabolas have the same focus $(4, 3)$ and their directrices are the x -axis and the y -axis, respectively. If these parabolas intersect at the points A and B , then $(AB)^2$ is equal to :
- (1) 392
(2) 384
(3) 192
(4) 96
- Q4.** Let the parabola $y = x^2 + px - 3$, meet the coordinate axes at the points P, Q and R . If the circle C with centre at $(-1, -1)$ passes through the points P, Q and R , then the area of $\triangle PQR$ is :
- (1) 7
(2) 4
(3) 6
(4) 5
- Q5.** Let the shortest distance from $(a, 0)$, $a > 0$, to the parabola $y^2 = 4x$ be 4. Then the equation of the circle passing through the point $(a, 0)$ and the focus of the parabola, and having its centre on the axis of the parabola is :
- (1) $x^2 + y^2 - 10x + 9 = 0$
(2) $x^2 + y^2 - 6x + 5 = 0$
(3) $x^2 + y^2 - 4x + 3 = 0$
(4) $x^2 + y^2 - 8x + 7 = 0$

Parabola

JEE Main 2025 January

Q6. Let A and B be the two points of intersection of the line $y + 5 = 0$ and the mirror image of the parabola $y^2 = 4x$ with respect to the line $x + y + 4 = 0$. If d denotes the distance between A and B, and a denotes the area of $\triangle SAB$, where S is the focus of the parabola $y^2 = 4x$, then the value of $(a + d)$ is _____.

Q7. If the line $3x - 2y + 12 = 0$ intersects the parabola $4y = 3x^2$ at the points A and B, then at the vertex of the parabola, the line segment AB subtends an angle equal to

- (1) $\tan^{-1}\left(\frac{4}{5}\right)$
- (2) $\tan^{-1}\left(\frac{9}{7}\right)$
- (3) $\tan^{-1}\left(\frac{11}{9}\right)$
- (4) $\frac{\pi}{2} - \tan^{-1}\left(\frac{3}{2}\right)$


Q8. Let ABCD be a trapezium whose vertices lie on the parabola $y^2 = 4x$. Let the sides AD and BC of the trapezium be parallel to y-axis. If the diagonal AC is of length $\frac{25}{4}$ and it passes through the point (1, 0), then the area of ABCD is

- (1) $\frac{75}{4}$
- (2) $\frac{25}{2}$
- (3) $\frac{125}{8}$
- (4) $\frac{75}{8}$

Q9. Let $P(4, 4\sqrt{3})$ be a point on the parabola $y^2 = 4ax$ and PQ be a focal chord of the parabola. If M and N are the foot of perpendiculars drawn from P and Q respectively on the directrix of the parabola, then the area of the quadrilateral PQMN is equal to :

- (1) $17\sqrt{3}$
- (2) $\frac{263\sqrt{3}}{8}$
- (3) $\frac{34\sqrt{3}}{3}$
- (4) $\frac{343\sqrt{3}}{8}$

Q10. Let $y^2 = 12x$ be the parabola and S be its focus. Let PQ be a focal chord of the parabola such that $(SP)(SQ) = \frac{147}{4}$. Let C be the circle described taking PQ as a diameter. If the equation of a circle C is $64x^2 + 64y^2 - \alpha x - 64\sqrt{3}y = \beta$, then $\beta - \alpha$ is equal to _____.



Q1. Let the product of the focal distances of the point $\left(\sqrt{3}, \frac{1}{2}\right)$ on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, (a > b)$, be $\frac{7}{4}$. Then the absolute difference of the eccentricities of two such ellipses is

- (1) $\frac{1-\sqrt{3}}{\sqrt{2}}$
- (2) $\frac{3-2\sqrt{2}}{2\sqrt{3}}$
- (3) $\frac{3-2\sqrt{2}}{3\sqrt{2}}$
- (4) $\frac{1-2\sqrt{2}}{\sqrt{3}}$

Q2. Let the ellipse $E_1 : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$ and $E_2 : \frac{x^2}{A^2} + \frac{y^2}{B^2} = 1, A < B$ have same eccentricity $\frac{1}{\sqrt{3}}$. Let the product of their lengths of latus rectums be $\frac{32}{\sqrt{3}}$, and the distance between the foci of E_1 be 4. If E_1 and E_2 meet at A, B, C and D , then the area of the quadrilateral $ABCD$ equals :

- (1) $\frac{12\sqrt{6}}{5}$
- (2) $6\sqrt{6}$
- (3) $\frac{18\sqrt{6}}{5}$
- (4) $\frac{24\sqrt{6}}{5}$

Q3. The length of the chord of the ellipse $\frac{x^2}{4} + \frac{y^2}{2} = 1$, whose mid-point is $\left(1, \frac{1}{2}\right)$, is :

- (1) $\frac{5}{3}\sqrt{15}$
- (2) $\frac{1}{3}\sqrt{15}$
- (3) $\frac{2}{3}\sqrt{15}$
- (4) $\sqrt{15}$

Q4. The equation of the chord, of the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$, whose mid-point is $(3, 1)$ is :

- (1) $48x + 25y = 169$
- (2) $5x + 16y = 31$
- (3) $25x + 101y = 176$
- (4) $4x + 122y = 134$

Ellipse

JEE Main 2025 January


Q5. If the midpoint of a chord of the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$ is $(\sqrt{2}, 4/3)$, and the length of the chord is $\frac{2\sqrt{\alpha}}{3}$, then α is :

- (1) 20
- (2) 22
- (3) 18
- (4) 26

Q6. If $\alpha x + \beta y = 109$ is the equation of the chord of the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$, whose mid point is $(\frac{5}{2}, \frac{1}{2})$, then $\alpha + \beta$ is equal to :

- (1) 58
- (2) 46
- (3) 37
- (4) 72

Q7. Let $E_1 : \frac{x^2}{9} + \frac{y^2}{4} = 1$ be an ellipse. Ellipses E_i 's are constructed such that their centres and eccentricities are same as that of E_1 , and the length of minor axis of E_i is the length of major axis of E_{i+1} ($i \geq 1$). If A_i is the area of the ellipse E_i , then $\frac{5}{\pi} (\sum_{i=1}^{\infty} A_i)$, is equal to



Hyperbola

JEE Main 2025 January

- Q1.** Let the foci of a hyperbola be $(1, 14)$ and $(1, -12)$. If it passes through the point $(1, 6)$, then the length of its latus-rectum is :
- (1) $\frac{24}{5}$
(2) $\frac{25}{6}$
(3) $\frac{144}{5}$
(4) $\frac{288}{5}$
- Q2.** Let $H_1 : \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ and $H_2 : -\frac{x^2}{A^2} + \frac{y^2}{B^2} = 1$ be two hyperbolas having length of latus rectums $15\sqrt{2}$ and $12\sqrt{5}$ respectively. Let their eccentricities be $e_1 = \sqrt{\frac{5}{2}}$ and e_2 respectively. If the product of the lengths of their transverse axes is $100\sqrt{10}$, then $25e_2^2$ is equal to _____.
- Q3.** If A and B are the points of intersection of the circle $x^2 + y^2 - 8x = 0$ and the hyperbola $\frac{x^2}{9} - \frac{y^2}{4} = 1$ and a point P moves on the line $2x - 3y + 4 = 0$, then the centroid of $\triangle PAB$ lies on the line :
- (1) $x + 9y = 36$
(2) $4x - 9y = 12$
(3) $6x - 9y = 20$
(4) $9x - 9y = 32$
- Q4.** Let $E : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, $a > b$ and $H : \frac{x^2}{A^2} - \frac{y^2}{B^2} = 1$. Let the distance between the foci of E and the foci of H be $2\sqrt{3}$. If $a - A = 2$, and the ratio of the eccentricities of E and H is $\frac{1}{3}$, then the sum of the lengths of their latus rectums is equal to:
- (1) 10
(2) 9
(3) 8
(4) 7
- Q5.** Let the circle C touch the line $x - y + 1 = 0$, have the centre on the positive x -axis, and cut off a chord of length $\frac{4}{\sqrt{13}}$ along the line $-3x + 2y = 1$. Let H be the hyperbola $\frac{x^2}{\alpha^2} - \frac{y^2}{\beta^2} = 1$, whose one of the foci is the centre of C and the length of the transverse axis is the diameter of C . Then $2\alpha^2 + 3\beta^2$ is equal to _____.

Trigonometric Ratios

JEE Main 2025 January

Q1. If $\sin x + \sin^2 x = 1$, $x \in \left(0, \frac{\pi}{2}\right)$, then $(\cos^{12} x + \tan^{12} x) + 3(\cos^{10} x + \tan^{10} x + \cos^8 x + \tan^8 x) + (\cos^6 x + \tan^6 x)$ is equal to :

- (1) 4
- (2) 1
- (3) 3
- (4) 2

Q2. The value of $(\sin 70^\circ)(\cot 10^\circ \cot 70^\circ - 1)$ is


- (1) $2/3$
- (2) 1
- (3) 0
- (4) $3/2$

Q3. If $\frac{\pi}{2} \leq x \leq \frac{3\pi}{4}$, then $\cos^{-1}\left(\frac{12}{13}\cos x + \frac{5}{13}\sin x\right)$ is equal to

- (1) $x - \tan^{-1} \frac{4}{3}$
- (2) $x + \tan^{-1} \frac{4}{5}$
- (3) $x - \tan^{-1} \frac{5}{12}$
- (4) $x + \tan^{-1} \frac{5}{12}$

Q4. If $\sum_{r=1}^{13} \left\{ \frac{1}{\sin\left(\frac{\pi}{4} + (r-1)\frac{\pi}{6}\right) \sin\left(\frac{\pi}{4} + \frac{r\pi}{6}\right)} \right\} = a\sqrt{3} + b$, $a, b \in \mathbf{Z}$, then $a^2 + b^2$ is equal to :

- (1) 10
- (2) 4
- (3) 2
- (4) 8



Trigonometric Equations

JEE Main 2025 January

Q1. The sum of all values of $\theta \in [0, 2\pi]$ satisfying $2 \sin^2 \theta = \cos 2\theta$ and $2 \cos^2 \theta = 3 \sin \theta$ is

(1) 4π

(2) $\frac{5\pi}{6}$

(3) π

(4) $\frac{\pi}{2}$



Inverse Trigonometric Functions

JEE Main 2025 January

Q1. If for some $\alpha, \beta; \alpha \leq \beta, \alpha + \beta = 8$ and $\sec^2(\tan^{-1} \alpha) + \operatorname{cosec}^2(\cot^{-1} \beta) = 36$, then $\alpha^2 + \beta$ is _____.

Q2. $\cos\left(\sin^{-1} \frac{3}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{33}{65}\right)$ is equal to:

(1) 1

(2) 0

(3) $\frac{32}{65}$

(4) $\frac{33}{65}$

Q3. Using the principal values of the inverse trigonometric functions, the sum of the maximum and the minimum values of $16 \left((\sec^{-1} x)^2 + (\operatorname{cosec}^{-1} x)^2 \right)$ is :

(1) $24\pi^2$

(2) $22\pi^2$

(3) $31\pi^2$

(4) $18\pi^2$

Q4. If $\alpha > \beta > \gamma > 0$, then the expression $\cot^{-1}\left\{\beta + \frac{(1+\beta^2)}{(\alpha-\beta)}\right\} + \cot^{-1}\left\{\gamma + \frac{(1+\gamma^2)}{(\beta-\gamma)}\right\} + \cot^{-1}\left\{\alpha + \frac{(1+\alpha^2)}{(\gamma-\alpha)}\right\}$ is equal to :

(1) π

(2) 0

(3) $\frac{\pi}{2} - (\alpha + \beta + \gamma)$

(4) 3π

Q5. Let $S = \{x : \cos^{-1} x = \pi + \sin^{-1} x + \sin^{-1}(2x + 1)\}$. Then $\sum_{x \in S} (2x - 1)^2$ is equal to _____.

Vector Algebra

JEE Main 2025 January

Q1. Let $\vec{a} = 3\hat{i} - \hat{j} + 2\hat{k}$, $\vec{b} = \vec{a} \times (\hat{i} - 2\hat{k})$ and $\vec{c} = \vec{b} \times \hat{k}$. Then the projection of $\vec{c} - 2\hat{j}$ on $\vec{a}$ is :

- (1) $2\sqrt{14}$
- (2) $\sqrt{14}$
- (3) $3\sqrt{7}$
- (4) $2\sqrt{7}$

Q2. If the components of $\vec{a} = \alpha\hat{i} + \beta\hat{j} + \gamma\hat{k}$ along and perpendicular to $\vec{b} = 3\hat{i} + \hat{j} - \hat{k}$ respectively, are $\frac{16}{11}(3\hat{i} + \hat{j} - \hat{k})$ and $\frac{1}{11}(-4\hat{i} - 5\hat{j} - 17\hat{k})$, then $\alpha^2 + \beta^2 + \gamma^2$ is equal to :

- (1) 26
- (2) 18
- (3) 23
- (4) 16

Q3. Let $\vec{a}$ and $\vec{b}$ be two unit vectors such that the angle between them is $\frac{\pi}{3}$. If $\lambda\vec{a} + 2\vec{b}$ and $3\vec{a} - \lambda\vec{b}$ are perpendicular to each other, then the number of values of λ in $[-1, 3]$ is :


- (1) 2
- (2) 1
- (3) 0
- (4) 3

Q4. Let the arc AC of a circle subtend a right angle at the centre O . If the point B on the arc AC , divides the arc AC such that $\frac{\text{length of arc } AB}{\text{length of arc } BC} = \frac{1}{5}$, and $\vec{OC} = \alpha\vec{OA} + \beta\vec{OB}$, then $\alpha + \sqrt{2}(\sqrt{3} - 1)\beta$ is equal to

- (1) $2\sqrt{3}$
- (2) $2 - \sqrt{3}$
- (3) $5\sqrt{3}$
- (4) $2 + \sqrt{3}$

Q5. Let $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$, $\vec{b} = 3\hat{i} + \hat{j} - \hat{k}$ and $\vec{c}$ be three vectors such that $\vec{c}$ is coplanar with $\vec{a}$ and $\vec{b}$. If the vector $\vec{C}$ is perpendicular to $\vec{b}$ and $\vec{a} \cdot \vec{c} = 5$, then $|\vec{c}|$ is equal to

- (1) $\sqrt{\frac{11}{6}}$



Vector Algebra

JEE Main 2025 January

(2) $\frac{1}{3\sqrt{2}}$

(3) 16

(4) 18

Q6. Let A, B, C be three points in xy -plane, whose position vector are given by $\sqrt{3}\hat{i} + \hat{j}$, $\hat{i} + \sqrt{3}\hat{j}$ and $a\hat{i} + (1-a)\hat{j}$ respectively with respect to the origin O. If the distance of the point C from the line bisecting the angle between the vectors $\vec{OA}$ and $\vec{OB}$ is $\frac{9}{\sqrt{2}}$, then the sum of all the possible values of a is :

(1) 2

(2) $9/2$

(3) 1

(4) 0

Q7. Let $\vec{c}$ be the projection vector of $\vec{b} = \lambda\hat{i} + 4\hat{k}$, $\lambda > 0$, on the vector $\vec{a} = \hat{i} + 2\hat{j} + 2\hat{k}$. If $|\vec{a} + \vec{c}| = 7$, then the area of the parallelogram formed by the vectors $\vec{b}$ and $\vec{c}$ is _____

Q8. Let the position vectors of three vertices of a triangle be $4\vec{p} + \vec{q} - 3\vec{r}$, $-5\vec{p} + \vec{q} + 2\vec{r}$ and $2\vec{p} - \vec{q} + 2\vec{r}$. If the position vectors of the orthocenter and the circumcenter of the triangle are $\frac{\vec{p} + \vec{q} + \vec{r}}{4}$ and $\alpha\vec{p} + \beta\vec{q} + \gamma\vec{r}$ respectively, then $\alpha + 2\beta + 5\gamma$ is equal to :

(1) 3

(2) 4

(3) 1

(4) 6

Q9. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + 2\hat{j} + \hat{k}$ and $\vec{d} = \vec{a} \times \vec{b}$. If $\vec{c}$ is a vector such that $\vec{a} \cdot \vec{c} = |\vec{c}|$, $|\vec{c} - 2\vec{a}|^2 = 8$ and the angle between $\vec{d}$ and $\vec{c}$ is $\frac{\pi}{4}$, then $|10 - 3\vec{b} \cdot \vec{c}| + |\vec{d} \times \vec{c}|^2$ is equal to

Q10. Let $\vec{a} = 2\hat{i} - \hat{j} + 3\hat{k}$, $\vec{b} = 3\hat{i} - 5\hat{j} + \hat{k}$ and $\vec{c}$ be a vector such that $\vec{a} \times \vec{c} = \vec{c} \times \vec{b}$ and $(\vec{a} + \vec{c}) \cdot (\vec{b} + \vec{c}) = 168$. Then the maximum value of $|\vec{c}|^2$ is :

(1) 462

(2) 77

(3) 154

Q11. Let $\hat{a}$ be a unit vector perpendicular to the vectors $\vec{b} = \hat{i} - 2\hat{j} + 3\hat{k}$ and $\vec{c} = 2\hat{i} + 3\hat{j} - \hat{k}$, and makes an angle of $\cos^{-1}\left(-\frac{1}{3}\right)$ with the vector $\hat{i} + \hat{j} + \hat{k}$. If $\hat{a}$ makes an angle of $\frac{\pi}{3}$ with the vector $\hat{i} + \alpha\hat{j} + \hat{k}$, then the value of α is :

- (1) $\sqrt{6}$
- (2) $-\sqrt{6}$
- (3) $-\sqrt{3}$
- (4) $\sqrt{3}$

Q12. Let the position vectors of the vertices A, B and C of a tetrahedron $ABCD$ be $\hat{i} + 2\hat{j} + \hat{k}$, $\hat{i} + 3\hat{j} - 2\hat{k}$ and $2\hat{i} + \hat{j} - \hat{k}$ respectively. The altitude from the vertex D to the opposite face ABC meets the median line segment through A of the triangle ABC at the point E . If the length of AD is $\frac{\sqrt{110}}{3}$ and the volume of the tetrahedron is $\frac{\sqrt{805}}{6\sqrt{2}}$, then the position vector of E is

- (1) $\frac{1}{12}(7\hat{i} + 4\hat{j} + 3\hat{k})$
- (2) $\frac{1}{2}(\hat{i} + 4\hat{j} + 7\hat{k})$ ★
- (3) $\frac{1}{6}(12\hat{i} + 12\hat{j} + \hat{k})$
- (4) $\frac{1}{6}(7\hat{i} + 12\hat{j} + \hat{k})$

Three Dimensional Geometry

JEE Main 2025 January

Q1. Let $L_1 : \frac{x-1}{3} = \frac{y-1}{-1} = \frac{z+1}{0}$ and $L_2 : \frac{x-2}{2} = \frac{y}{0} = \frac{z+4}{\alpha}$, $\alpha \in \mathbf{R}$, be two lines, which intersect at the point B . If P is the foot of perpendicular from the point $A(1, 1, -1)$ on L_2 , then the value of $26\alpha(PB)^2$ is _____

Q2. The distance of the line $\frac{x-2}{2} = \frac{y-6}{3} = \frac{z-3}{4}$ from the point $(1, 4, 0)$ along the line $\frac{x}{1} = \frac{y-2}{2} = \frac{z+3}{3}$ is :

(1) $\sqrt{17}$

(2) $\sqrt{15}$

(3) $\sqrt{14}$

(4) $\sqrt{13}$

Q3. If the square of the shortest distance between the lines $\frac{x-2}{1} = \frac{y-1}{2} = \frac{z+3}{-3}$ and $\frac{x+1}{2} = \frac{y+3}{4} = \frac{z+5}{-5}$ is $\frac{m}{n}$, where m, n are coprime numbers, then $m + n$ is equal to :

(1) 21

(2) 9

(3) 14

(4) 6

Q4. If the image of the point $(4, 4, 3)$ in the line $\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-1}{3}$ is (α, β, γ) , then $\alpha + \beta + \gamma$ is equal to

(1) 9

(2) 12

(3) 7

(4) 8

Q5. Let $L_1 : \frac{x-1}{1} = \frac{y-2}{-1} = \frac{z-1}{2}$ and $L_2 : \frac{x+1}{-1} = \frac{y-2}{2} = \frac{z}{1}$ be two lines.

Let L_3 be a line passing through the point (α, β, γ) and be perpendicular to both L_1 and L_2 . If L_3 intersects L_1 , then $|5\alpha - 11\beta - 8\gamma|$ equals :

(1) 20

(2) 18

(3) 25

(4) 16

Three Dimensional Geometry

JEE Main 2025 January

Q6. Let a straight line L pass through the point $P(2, -1, 3)$ and be perpendicular to the lines $\frac{x-1}{2} = \frac{y+1}{1} = \frac{z-3}{-2}$ and $\frac{x-3}{1} = \frac{y-2}{3} = \frac{z+2}{4}$. If the line L intersects the yz -plane at the point Q , then the distance between the points P and Q is :

(1) $\sqrt{10}$

(2) $2\sqrt{3}$

(3) 2

(4) 3

Q7. Let $A(x, y, z)$ be a point in xy -plane, which is equidistant from three points $(0, 3, 2)$, $(2, 0, 3)$ and $(0, 0, 1)$.

Let $B = (1, 4, -1)$ and $C = (2, 0, -2)$. Then among the statements

(S1) : $\triangle ABC$ is an isosceles right angled triangle, and

(S2) : the area of $\triangle ABC$ is $\frac{9\sqrt{2}}{2}$,

(1) both are true

(2) only (S2) is true

(3) only (S1) is true

(4) both are false

Q8. Let $L_1 : \frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ and $L_2 : \frac{x-2}{3} = \frac{y-4}{4} = \frac{z-5}{5}$ be two lines. Then which of the following points lies on the line of the shortest distance between L_1 and L_2 ?

(1) $\left(\frac{14}{3}, -3, \frac{22}{3}\right)$

(2) $\left(-\frac{5}{3}, -7, 1\right)$

(3) $\left(2, 3, \frac{1}{3}\right)$

(4) $\left(\frac{8}{3}, -1, \frac{1}{3}\right)$

Q9. Let a line pass through two distinct points $P(-2, -1, 3)$ and Q , and be parallel to the vector $3\hat{i} + 2\hat{j} + 2\hat{k}$. If the distance of the point Q from the point $R(1, 3, 3)$ is 5, then the square of the area of $\triangle PQR$ is equal to :

(1) 148

(2) 136

(3) 144

(4) 140

Three Dimensional Geometry

JEE Main 2025 January

Q10. The perpendicular distance, of the line $\frac{x-1}{2} = \frac{y+2}{-1} = \frac{z+3}{2}$ from the point $P(2, -10, 1)$, is :

- (1) 6
- (2) $5\sqrt{2}$
- (3) $4\sqrt{3}$
- (4) $3\sqrt{5}$

Q11. Let the distance between two parallel lines be 5 units and a point P lie between the lines at a unit distance from one of them. An equilateral triangle PQR is formed such that Q lies on one of the parallel lines, while R lies on the other. Then $(QR)^2$ is equal to _____ -.

Q12. Let P be the foot of the perpendicular from the point $Q(10, -3, -1)$ on the line $\frac{x-3}{7} = \frac{y-2}{-1} = \frac{z+1}{-2}$. Then the area of the right angled triangle PQR , where R is the point $(3, -2, 1)$, is

- (1) $9\sqrt{15}$
- (2) $\sqrt{30}$
- (3) $8\sqrt{15}$
- (4) $3\sqrt{30}$

Q13. Let in a $\triangle ABC$, the length of the side AC be 6, the vertex B be $(1, 2, 3)$ and the vertices A, C lie on the line $\frac{x-6}{3} = \frac{y-7}{2} = \frac{z-7}{-2}$. Then the area (in sq. units) of $\triangle ABC$ is:

- (1) 17
- (2) 21
- (3) 56
- (4) 42

Q14. Let the line passing through the points $(-1, 2, 1)$ and parallel to the line $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z}{4}$ intersect the line $\frac{x+2}{3} = \frac{y-3}{2} = \frac{z-4}{1}$ at the point P . Then the distance of P from the point $Q(4, -5, 1)$ is

- (1) 5
- (2) $5\sqrt{5}$
- (3) $5\sqrt{6}$
- (4) 10

Three Dimensional Geometry

JEE Main 2025 January

Q15. Let P be the image of the point Q(7, -2, 5) in the line $L : \frac{x-1}{2} = \frac{y+1}{3} = \frac{z}{4}$ and R(5, p, q) be a point on L. Then the square of the area of $\triangle PQR$ is _____.

Q16. The square of the distance of the point $\left(\frac{15}{7}, \frac{32}{7}, 7\right)$ from the line $\frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7}$ in the direction of the vector $\hat{i} + 4\hat{j} + 7\hat{k}$ is :

(1) 54

(2) 44

(3) 41

(4) 66

Q17. Let $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ and $\vec{b} = 2\hat{i} + 7\hat{j} + 3\hat{k}$. Let $L_1 : \vec{r} = (-\hat{i} + 2\hat{j} + \hat{k}) + \lambda \vec{a}, \lambda \in \mathbf{R}$ and $L_2 : \vec{r} = (\hat{j} + \hat{k}) + \mu \vec{b}, \mu \in \mathbf{R}$ be two lines. If the line L_3 passes through the point of intersection of L_1 and L_2 , and is parallel to $\vec{a} + \vec{b}$, then L_3 passes through the point :

(1) (5, 17, 4)

(2) (2, 8, 5)

(3) (8, 26, 12)

(4) (-1, -1, 1)

Q18. Let the point A divide the line segment joining the points $P(-1, -1, 2)$ and $Q(5, 5, 10)$ internally in the ratio $r : 1 (r > 0)$. If O is the origin and $(\vec{OQ} \cdot \vec{OA}) - \frac{1}{5} |\vec{OP} \times \vec{OA}|^2 = 10$, then the value of r is :

(1) $\sqrt{7}$

(2) 14

(3) 3

(4) 7

Q19. Let P be the foot of the perpendicular from the point (1, 2, 2) on the line $L : \frac{x-1}{1} = \frac{y+1}{-1} = \frac{z-2}{2}$. Let the line $\vec{r} = (-\hat{i} + \hat{j} - 2\hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k}), \lambda \in \mathbf{R}$, intersect the line L at Q. Then $2(PQ)^2$ is equal to :

(1) 25

(2) 19

(3) 29

(4) 27

Three Dimensional Geometry

JEE Main 2025 January



Basic of Mathematics

1. (1)

Quadratic Equation

1. (4) 2. (3) 3. (4) 4. (117) 5. (1) 6. (2)

Complex Number

1. (2) 2. (4) 3. (31) 4. (4) 5. (2) 6. (2) 7. (1) 8. (2)
9. (1) 10. (10)

Sequences and Series

1. (1) 2. (474) 3. (3) 4. (1) 5. (125) 6. (2) 7. (20) 8. (11132)
9. (4) 10. (2) 11. (2) 12. (2) 13. (4) 14. (2) 15. (4) 16. (2)

Permutation Combination

1. (3) 2. (3) 3. (4) 4. (64) 5. (4) 6. (1405) 7. (1) 8. (2)
9. (1) 10. (17280)

Binomial Theorem

1. (612) 2. (3) 3. (3) 4. (1) 5. (2) 6. (1) 7. (1) 8. (4)
9. (2035) 10. (5) 11. (465)

Statistics

1. (8788) 2. (3) 3. (3) 4. (1)

Matrices

1. (1) 2. (2) 3. (4) 4. (1613) 5. (2) 6. (34) 7. (44) 8. (3)
9. (4) 10. (3)

Determinants

1. (1) 2. (3) 3. (1) 4. (1) 5. (4) 6. (1) 7. (3) 8. (4)

Probability

1. (3) 2. (2) 3. (1) 4. (2) 5. (4) 6. (2) 7. (2) 8. (2)
9. (4) 10. (1) 11. (1) 12. (3) 13. (1)

Sets and Relations

1. (2) 2. (2) 3. (2) 4. (2) 5. (4) 6. (5120) 7. (2) 8. (4)
9. (3) 10. (2) 11. (2)

Functions

1. (2) 2. (392) 3. (3) 4. (3) 5. (1) 6. (4) 7. (1) 8. (2)
9. (2) 10. (3) 11. (3) 12. (4) 13. (4) 14. (4)

Limits

1. (4) 2. (4) 3. (64) 4. (24) 5. (3) 6. (1)

Continuity and Differentiability

1. (4) 2. (1) 3. (2) 4. (5) 5. (3) 6. (34)

Application of Derivatives

1. (2) 2. (2) 3. (30) 4. (1)

Indefinite Integration

1. (2) 2. (16) 3. (2) 4. (2) 5. (1)

Definite Integration

1. (2) 2. (1) 3. (12) 4. (1) 5. (1) 6. (4) 7. (4) 8. (4)
9. (2) 10. (3) 11. (3) 12. (4) 13. (112)

Area Under Curves

1. (2) 2. (1) 3. (2) 4. (2) 5. (1) 6. (77) 7. (4) 8. (2)
9. (3) 10. (3) 11. (4) 12. (1)

Differential Equations

1. (1) 2. (4) 3. (3) 4. (4) 5. (3) 6. (2) 7. (19) 8. (1)
9. (4) 10. (27) 11. (1) 12. (4)

Straight Lines

1. (4) 2. (4) 3. (145) 4. (2) 5. (3) 6. (3) 7. (3) 8. (4)
9. (1) 10. (3)

Circle

1. (4) 2. (3) 3. (3) 4. (2) 5. (2)

Parabola

1. (15) 2. (2) 3. (3) 4. (3) 5. (2) 6. (14) 7. (2) 8. (1)
9. (4) 10. (1328)

Ellipse

1. (2) 2. (4) 3. (3) 4. (1) 5. (2) 6. (1) 7. (54)

Hyperbola

1. (4) 2. (55) 3. (3) 4. (3) 5. (19)

Trigonometric Ratios & Identities

1. (4) 2. (2) 3. (3) 4. (4)

Trigonometric Equations

1. (3)

Inverse Trigonometric Functions

1. (14) 2. (2) 3. (2) 4. (1) 5. (5)

Vector Algebra

1. (1) 2. (1) 3. (3) 4. (2) 5. (1) 6. (3) 7. (16) 8. (1)
9. (6) 10. (4) 11. (2) 12. (4)

Three Dimensional Geometry

1. (216) 2. (3) 3. (2) 4. (1) 5. (3) 6. (4) 7. (3) 8. (1)
9. (2) 10. (4) 11. (28) 12. (4) 13. (2) 14. (2) 15. (957) 16. (4)
17. (3) 18. (4) 19. (4)

Q1. $e^{5(\ln x)^2+3} = x^8$

(1) $\Rightarrow \ln e^{5(\ln x)^2+3} = \ln x^8$

$$\Rightarrow 5(\ln x)^2 + 3 = 8 \ln x$$

$$(\ln x = t)$$

$$\Rightarrow 5t^2 - 8t + 3 = 0$$

$$t_1 + t_2 = \frac{8}{5}$$

$$\ln x_1 x_2 = \frac{8}{5}$$

$$x_1 x_2 = e^{8/5}$$



Quadratic Equation

JEE Main 2025 January

Q1. Consider $\frac{1}{\sqrt{x}} = \alpha$ $x > 0$

(4) $\{9\alpha^2 - 9\alpha + 2\} \{2\alpha^2 - 7\alpha + 3\} = 0$

$$(3\alpha - 2)(3\alpha - 1)(\alpha - 3)(2\alpha - 1) = 0$$

$$\alpha = \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, 3$$

$$x = 9, 4, \frac{9}{4}, \frac{1}{9}$$

So, no. of solutions = 4

Q2. For $x \geq \frac{3}{2}$

(3) $x^2 + 2x - 3 - 4 = 0$

$$x^2 + 2x - 7 = 0$$

$$x = \frac{-2 \pm \sqrt{4 + 28}}{2} = -1 \pm 2\sqrt{2}$$

Only $2\sqrt{2} - 1$ is acceptable root

For $x < \frac{3}{2}$

$$x^2 - 2x + 3 - 4 = 0$$

$$x^2 - 2x - 1 = 0$$

$$x = \frac{2 \pm \sqrt{4 + 4}}{2} = 1 \pm \sqrt{2}$$

Only $1 - \sqrt{2}$ is acceptable

$$\begin{aligned} \text{Sum of the square} &= (1 - \sqrt{2})^2 + (2\sqrt{2} - 1)^2 \\ &= 6(2 - \sqrt{2}) \end{aligned}$$

Q3. $(a - 5)^2 - 8(15 - 3a) < 0$

(4) $a^2 + 14a + 25 - 120 < 0$

$$a^2 + 14a - 95 < 0$$

$$(a + 19)(a - 5) < 0$$

$$a \in (-19, 5)$$

$$\therefore -19 < x < 5$$

$$\therefore \sum_{x \in X} x^2 = (1^2 + 2^2 + \dots + 4^2) + (1^2 + 2^2 + \dots + 18^2)$$

$$= \frac{4 \times 5 \times 9}{6} + \frac{18 \times 19 \times 37}{6}$$

$$= 30 + 2109$$

$$= 2139$$

Quadratic Equation

JEE Main 2025 January

Q4. $a(b-c)x^2 + b(c-a)x + c(a-b) = 0$

(117) $x = 1$ is root $\therefore$ other root is 1

$$\alpha + \beta = -\frac{b(c-a)}{a(b-c)} = 2$$

$$\Rightarrow -bc + ab = 2ab - 2ac$$

$$\Rightarrow 2ac = ab + bc$$

$$\Rightarrow 2ac = b(a+c)$$

$$\Rightarrow 2ac = 15b \dots (1)$$

$$\Rightarrow 2ac = 15 \left(\frac{36}{5} \right) = 108$$

$$\Rightarrow ac = 54$$

$$a + c = 15$$

$$a^2 + c^2 + 2ac = 225$$

$$a^2 + c^2 = 225 - 108 = 117$$

Q5. $(x^2 - 9x + 11)^2 - (x-4)(x-5) = 3$

(1) $(x^2 - 9x + 11)^2 - (x^2 - 9x + 20) = 3$

Let $x^2 - 9x + 11 = t$

$$t^2 - (t+9) = 3$$

$$\Rightarrow t^2 - t - 12 = 0$$

$$\Rightarrow t^2 - 4t + 3t - 12 = 0$$

$$\Rightarrow t(t-4) + 3(t-4) = 0$$

$$\Rightarrow t = 4 \text{ or } -3$$

$$x^2 - 9x + 11 = 4$$

$$x^2 - 9x + 7 = 0$$

Here, we will get irrational roots

$$x^2 - 9x + 11 = -3$$

$$x^2 - 9x + 14 = 0$$

$$x^2 - 7x - 2x + 14 = 0$$

$$\Rightarrow x = 7, 2$$

$$\Rightarrow \text{Product of all rational roots} = 14$$

Q6. $2x^2 + (\cos \theta)x - 1 = 0$

(2) $\alpha_\theta + \beta_\theta = \frac{-\cos \theta}{2}$

$$\alpha_\theta \cdot \beta_\theta = \frac{-1}{2}$$

$$\alpha_\theta^2 + \beta_\theta^2 = (\alpha_\theta + \beta_\theta)^2 - 2\alpha_\theta\beta_\theta \frac{\cos^2 \theta}{4} + 1$$

$$\alpha_\theta^4 + \beta_\theta^4 = (\alpha_\theta^2 + \beta_\theta^2)^2 - 2\alpha_\theta^2\beta_\theta^2 = \left(\frac{\cos^2 \theta}{4} + 1 \right)^2 - \frac{2}{4}$$

$$= \left(\frac{\cos^2 \theta}{4} + 1 \right)^2 - \frac{1}{2}$$

Maximum when $\cos \theta = 1$

Quadratic Equation

JEE Main 2025 January

$$M = \left(\frac{1}{4} + 1\right)^2 - \frac{1}{2}$$

$$M = \frac{17}{16}$$

Minimum when $\cos \theta = 0$

$$m = 1 - \frac{1}{2} = \frac{1}{2}$$

$$16(M + m) = 16\left(\frac{17}{16} + \frac{1}{2}\right) = 25$$



Q1. $|z| = 1$

(2) $\left| \frac{z}{\bar{z}} + \frac{\bar{z}}{z} \right| = 1$

$$\Rightarrow |z^2 + (\bar{z})^2| = 1$$

Let $z = x + iy$

$$\Rightarrow |(x + iy)^2 + (x - iy)^2| = 1$$

$$\Rightarrow |2x^2 - 2y^2| = 1$$



$$\Rightarrow |x^2 - y^2| = \frac{1}{2}$$

$$\Rightarrow x^2 - y^2 = \frac{\pm 1}{2}$$

and $x^2 + y^2 = 1$

Case I: $x^2 - y^2 = \frac{1}{2}$


Case II: $x^2 - y^2 = -\frac{1}{2}$



Hence, we get 8 complex number

Q2.

(4)



$$\therefore AB = \sqrt{100} = 10$$

$$\therefore |Z_1 - Z_2|_{\min} = 10 - 2 - 1 = 7$$

Q3. $\alpha + \beta = a$ $\alpha\beta = -b$

(31) $P_6 = aP_5 + bP_4$

$$45\sqrt{7}i = a \times 11\sqrt{7}i + b(-3\sqrt{7})i$$

$$45 = 11a - 3b$$

and

Complex Number

JEE Main 2025 January

$$P_5 = aP_4 + bP_3$$

$$11\sqrt{7}i = a(-3\sqrt{7}i) + b(-5\sqrt{7}i)$$

$$11 = -3a - 5b$$

$$a = 3, b = -4$$

$$|\alpha^4 + \beta^4| = \sqrt{(\alpha^4 - \beta^4)^2 + 4\alpha^4\beta^4}$$

$$= \sqrt{-63 + 4.4^4}$$

$$= \sqrt{-63 + 1024} = \sqrt{961} = 31$$

Q4. $x^2 - (3 - 2i)x - (2i - 2) = 0$

(4) $x = \frac{(3 - 2i) \pm \sqrt{(3 - 2i)^2 - 4(1)(-(2i - 2))}}{2(1)}$

$$= \frac{(3 - 2i) \pm \sqrt{9 - 4 - 12i + 8i - 8}}{2}$$

$$= \frac{3 - 2i \pm \sqrt{-3 - 4i}}{2}$$

$$= \frac{3 - 2i \pm \sqrt{(1)^2 + (2i)^2 - 2(1)(2i)}}{2}$$

$$= \frac{3 - 2i \pm (1 - 2i)}{2}$$

$$\Rightarrow \frac{3 - 2i + 1 - 2i}{2} \text{ or } \frac{3 - 2i - 1 + 2i}{2}$$

$$\Rightarrow 2 - 2i \text{ or } 1 + 0i$$

$$\text{So } \alpha\gamma + \beta\delta = 2(1) + (-2)(0) = 2$$

Q5. $|z| = 1$

(2) $\arg(z_1) = -\frac{\pi}{4}, \arg(z_2) = 0, \arg(z_3) = \frac{\pi}{4}$

$$z_1 = |1|e^{-\frac{\pi}{4}} = \frac{1}{\sqrt{2}} - \frac{i}{\sqrt{2}}$$

$$z_2 = 1 + 0i$$

$$z_3 = \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}$$

$$z_1\bar{z}_2 = \frac{1 - i}{\sqrt{2}}$$

$$z_2\bar{z}_3 = \frac{1 - i}{\sqrt{2}}$$

$$z_3\bar{z}_1 = \frac{(1 + i)^2}{2}$$

$$z_1\bar{z}_2 + z_2\bar{z}_3 + z_3\bar{z}_1 = \sqrt{2} + i(1 - \sqrt{2})$$

$$|z_1\bar{z}_2 + z_2\bar{z}_3 + z_3\bar{z}_1|^2 = 5 - 2\sqrt{2}$$

$$\alpha = 5, \beta = -2$$


$$\alpha_2^2 + \beta^2 = 29$$

Complex Number

JEE Main 2025 January

Q6.

(2)



$$OA = |z_1| = \sqrt{3+8} = \sqrt{11}$$

$$\text{and } OB = \frac{1}{\sqrt{3}}|z_1| = \sqrt{\frac{11}{3}}$$

$$AB^2 = OA^2 + OB^2 - 2 \cdot OA \cdot OB \cos \frac{\pi}{6}$$

$$= 11 + \frac{11}{3} - 2 \cdot \frac{11}{\sqrt{3}} \cdot \frac{\sqrt{3}}{2}$$

$$\therefore AB = \sqrt{\frac{11}{3}}$$

$$\therefore \text{Area of } \triangle ABD = \frac{1}{2} \cdot OA \cdot OB \cdot \sin \frac{\pi}{6}$$


$$= \frac{11}{4\sqrt{3}} \text{ sq. units}$$

$$\text{Here } OB = AB \text{ and } \angle A = \frac{2\pi}{3}$$

$\therefore \triangle ABD$ is an obtuse angled isosceles triangle.

Q7.

(1)



$$\text{Let } z = x + iy$$

$$(x + iy)(1 + i) + (x - iy)(1 - i) = 4$$

$$x + ix + iy - y + x - ix - iy - y = 4$$

$$2x - 2y = 4$$

$$x - y = 2$$

$$|z - 3| \leq 1$$

$$(x - 3)^2 + y^2 \leq 1$$

$$\text{Area of shaded region} = \frac{\pi \cdot 1^2}{4} - \frac{1}{2} \cdot 1 \cdot 1 = \frac{\pi}{4} - \frac{1}{2}$$

Area of unshaded region inside the circle

$$= \frac{3}{4}\pi \cdot 1^2 + \frac{1}{2} \cdot 1 \cdot 1 = \frac{3\pi}{4} + \frac{1}{2}$$

$$\therefore \text{difference of area} = \left(\frac{3\pi}{4} + \frac{1}{2}\right) - \left(\frac{\pi}{4} - \frac{1}{2}\right)$$

$$= \frac{\pi}{2} + 1$$

Q8. Let $z = x + iy \Rightarrow \bar{z} = x - iy$

(2) $3|\bar{z} - i| = 1|2\bar{z} + i|$


$$= 3|(x - (y + 1)i)| = |2x + i(1 - 2y)|$$

$$= 3\sqrt{x^2 + (y + 1)^2} = \sqrt{(2x)^2 + (1 - 2y)^2}$$

$$= 9(x^2 + y^2 + 2y + 1) = 4x^2 + 4y^2 - 4y + 1$$

$$\Rightarrow 5x^2 + 5y^2 + 22y + 8 = 0$$

$$\Rightarrow \text{Centre} \equiv \left(0, -\frac{11}{5}\right)$$



Area of Δ

$$= \frac{1}{2}|\alpha| \left| \frac{-11}{5} \right| = 11$$

$$\Rightarrow |\alpha| = 10$$

$$\Rightarrow \alpha^2 = 100$$

Q9. $2z^2 - 3z - 2i = 0 \dots(i)$

(1) $2\left(z - \frac{i}{z}\right) = 3$

As α, β are roots of (i)

$$\alpha - \frac{i}{\alpha} = \frac{3}{2}$$


$$\Rightarrow \alpha^2 - \frac{1}{\alpha^2} - 2i = \frac{9}{4}$$

$$\Rightarrow \alpha^2 - \frac{1}{\alpha^2} = \frac{9}{4} + 2i$$

Squaring both sides

$$\Rightarrow \alpha^4 + \frac{1}{\alpha^4} - 2 = \frac{81}{16} - 4 + 9i$$

$$\Rightarrow \alpha^4 + \frac{1}{\alpha^4} = \frac{49}{16} + 9i$$



Complex Number

JEE Main 2025 January

Similarly, $\beta^4 + \frac{1}{\beta^4} = \frac{49}{16} + 9i$

$$\frac{\alpha^{19} + \beta^{19} + \alpha^{11} + \beta^{11}}{\alpha^{15} + \beta^{15}} = \frac{\alpha^{15} \left(\alpha^4 + \frac{1}{\alpha^4} \right) + \beta^{15} \left(\beta^4 + \frac{1}{\beta^4} \right)}{\alpha^{15} + \beta^{15}}$$

$$= \frac{49}{16} + 9i$$

$$\operatorname{Re} \left(\frac{\alpha^{19} + \beta^{19} + \alpha^{11} + \beta^{11}}{\alpha^{15} + \beta^{15}} \right) = \frac{49}{16}$$

$$\operatorname{Im} \left(\frac{\alpha^{19} + \beta^{19} + \alpha^{11} + \beta^{11}}{\alpha^{15} + \beta^{15}} \right) = 9$$

$$\Rightarrow 16 \operatorname{Re} \left(\frac{\alpha^{19} + \beta^{19} + \alpha^{11} + \beta^{11}}{\alpha^{15} + \beta^{15}} \right) \cdot \operatorname{Im} \left(\frac{\alpha^{19} + \beta^{19} + \alpha^{11} + \beta^{11}}{\alpha^{15} + \beta^{15}} \right)$$

$$= 16 \times \frac{49}{16} \times 9$$

$$= 441$$

Q10. $a, b \in I, -3 \leq a, b \leq 3, a + b \neq 0$

$$(10) \quad \begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 1$$

$$\Rightarrow \begin{vmatrix} z & z & z \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 1$$

$$\Rightarrow z \begin{vmatrix} 1 & 1 & 1 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 1$$

$$\Rightarrow z \begin{vmatrix} 1 & 0 & 0 \\ \omega & z+\omega^2-\omega & 1-\omega \\ \omega^2 & 1-\omega^2 & z+\omega-\omega^2 \end{vmatrix} = 1$$

$$\Rightarrow z^3 = 1$$

$$\Rightarrow z = \omega, \omega^2, 1$$

Now

$$|1-a| = |1+b|$$

$$\Rightarrow 10 \text{ pairs}$$

Q1. (1) $a = 3$

$$S_4 = \frac{1}{5}(S_8 - S_4)$$

$$\Rightarrow 5S_4 = S_8 - S_4$$

$$\Rightarrow 6S_4 = S_8$$

$$\Rightarrow 6 \cdot \frac{4}{2}[2 \times 3 + (4 - 1) \times d]$$

$$= \frac{8}{2}[2 \times 3 + (8 - 1)d]$$

$$\Rightarrow 12(6 + 3d) = 4(6 + 7d)$$

$$\Rightarrow 18 + 9d = 6 + 7d$$

$$\Rightarrow d = -6$$

$$S_{20} = \frac{20}{2}[2 \times 3 + (20 - 1)(-6)]$$

$$= 10[6 - 114]$$

$$= -1080$$

Q2. $S_{11} = \frac{11}{2}(2a + 10d) = 88$

(474)

$$a + 5d = 8$$

$$a = 8 - 5 \times \frac{3}{2} = \frac{1}{2}$$

Roots are

$$T_{10} = a + 9d = \frac{1}{2} + 9 \times \frac{3}{2} = 14$$

$$T_{11} = a + 10d = \frac{1}{2} + 10 \times \frac{3}{2} = \frac{31}{2}$$

$$\frac{p}{3} = T_{10} + T_{11} = 14 + \frac{31}{2} = \frac{59}{2}$$

$$p = \frac{177}{2}$$

$$\frac{q}{3} = T_{10} \times T_{11} = 7 \times 31 = 217$$

$$q = 651$$

$$q - 2p$$

$$= 651 - 177$$

$$= 474$$

Q3. (3) $S_{40} = 1030 \Rightarrow \frac{40}{2}[2a + 39d] = 1030$

$$\Rightarrow 2a + 39d = \frac{103}{2} \dots (1)$$

$$S_{12} = 57 \Rightarrow \frac{12}{2}[2a + 11d] = 57$$

$$\Rightarrow 2a + 11d = \frac{57}{6} \dots (2)$$

Equation (1) - equation (2)

Sequences and Series

JEE Main 2025 January

$$28d = \frac{103}{2} - \frac{57}{6}$$

$$28d = \frac{309 - 57}{6}$$

$$d = \frac{3}{2}$$

$$\Rightarrow a = -\frac{7}{2}$$

$$S_{30} - S_{10} = \frac{30}{2}[2a + 29d] - \frac{10}{2}[2a + 9d]$$

$$= 15[2a + 29d] - 5[2a + 9d]$$

$$= 5[6a + 87d - 2a - 9d]$$

$$= 5[4a + 78d]$$

$$= 5[-14 + 117]$$

$$= 515$$

Q4. (1) $S_3 = 3a + 3d = 54$

$$\Rightarrow a + d = 18$$

$$S_{20} = 10(2a + 19d)$$

$$\Rightarrow 10(36 + 17d)$$

$$\Rightarrow 1600 < 10(36 + 17d) < 1800$$

$$\Rightarrow 160 < 36 + 17d < 180$$

$$\Rightarrow 124 < 17d < 144$$

$$\Rightarrow 7\frac{5}{17} < d < 8\frac{8}{17}$$

Common difference will be natural number

$$\Rightarrow d = 8 \Rightarrow a = 10$$

$$\Rightarrow a_{11} = 10 + 10 \times 8 = 90$$

Q5. No. of 3 digits = $999 - 99 = 900$

(125) No. of 3 digit numbers divisible by 2 & 3 i.e. by 6

$$\frac{900}{6} = 150$$

No. of 3 digit numbers divisible by 4 & 9 i.e. by 36

$$\frac{900}{36} = 25$$

$\therefore$ No of 3 digit numbers divisible by 2&3 but not by 4&9

$$150 - 25 = 125$$

Q6. (2) $T_m = \frac{1}{25}, T_{25} = \frac{1}{20}, 20 \sum_{r=1}^{25} T_r = 13$

$$T_m = a + (m-1)d = \frac{1}{25} \dots\dots\dots (1)$$

$$T_{25} = a + 24d = \frac{1}{20}$$

$$20 \cdot \frac{25}{2} \left[a + \frac{1}{20} \right] = 13 \Rightarrow a = \frac{1}{500}$$

$$\text{also, } 20 S_{25} = 20 \cdot \frac{25}{2} [2a + 24d] = 13 \Rightarrow d = \frac{1}{500}$$

$$\text{from (1)} \quad \frac{1}{500} + \frac{m-1}{500} = \frac{1}{25} \Rightarrow m = 20$$

Now,

$$5 \sum_{r=m}^{2m} T_r = 100 \sum_{r=20}^{40} T_r = 126$$

Q7. $\frac{n}{2}[2a + (n-1)6] = (n-2)180^\circ$

(20) and $an + 3n^2 - 3n = 3n(n-2)180^\circ \dots (i)$

$$\therefore \text{Given } a + (n-1)6^\circ = 219^\circ$$

$$\Rightarrow a = 225^\circ - 6n^\circ$$

Putting value of a in (i)

$$\text{We get } (225 - 6n^2) + 3n^2 - 3n = 180n - 360^\circ$$

$$\Rightarrow 2n^2 - 42n - 360 = 0$$

$$\Rightarrow n^2 - 14n - 120 = 0$$

$$\Rightarrow (n-20)(n+6) = 0$$

$$\Rightarrow n = 20, -6 \text{ (Rejected)}$$

$$\therefore n = 20$$

Q8. As $a_1 + a_5 + a_{10} + \dots + a_{2020} + a_{2024} = 2233 \dots (1)$

(11132) We know in arithmetic progression.

Sum of terms equidistant from ends is equal

$\therefore$ from (1)

$$a_1 + a_{2024} = a_5 + a_{2020} = a_{10} + a_{2015} = \dots$$

$$\Rightarrow \underbrace{203 \text{ pairs}}_{203 \text{ pairs}} (a_1 + a_{2024}) = 2233$$

$$\Rightarrow a_1 + a_{2024} = 11$$

$$\text{Now } \sum_{i=1}^{2024} a_i = S_{2024} = \frac{2024}{2} [a_1 + a_{2024}]$$

$$= 1012(11)$$

$$= 11132$$

Q9. (4) $a_1 \cdot a_5 = 28 \Rightarrow a \cdot ar^4 = 28 \Rightarrow a^2 r^4 = 28 \dots (1)$

$$a_2 + a_4 = 29 \Rightarrow ar + ar^3 = 29$$

$$\Rightarrow ar(1 + r^2) = 29$$

$$\Rightarrow a^2 r^2 (1 + r^2)^2 = (29)^2 \dots (2)$$

By Eq. (1) & (2)

$$\frac{r^2}{(1 + r^2)^2} = \frac{28}{29 \times 29}$$

$$\Rightarrow \frac{r}{1 + r^2} = \frac{\sqrt{28}}{29} \Rightarrow r = \sqrt{28}$$

$$\therefore a^2 r^4 = 28 \Rightarrow a^2 \times (28)^2 = 28$$

$$\Rightarrow a = \frac{1}{\sqrt{28}}$$

$$\therefore a_6 = ar^5 = \frac{1}{\sqrt{28}} \times (28)^2 \sqrt{28} = 784$$

Q10. $S = a + (a + d)r + (a + 2d)r^2 + \dots$

(2) Then $S = \frac{a}{1-r} + \frac{dr}{(1-r)^2}, |r| < 1$

Since, $r = \frac{1}{7}$ and $a = 5, d = \alpha$

$$7 = \frac{5}{1 - \frac{1}{7}} + \frac{\alpha \cdot \frac{1}{7}}{\left(1 - \frac{1}{7}\right)^2}$$

$$\Rightarrow \alpha = 6$$

Q11. $T_n = S_n - S_{n-1}$

(2) $\Rightarrow T_n = \frac{1}{8}(2n-1)(2n+1)(2n+3)$

$$\Rightarrow \frac{1}{T_n} = \frac{8}{(2n-1)(2n+1)(2n+3)}$$

$$\lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{T_r} = \lim_{n \rightarrow \infty} 8 \sum_{r=1}^n \frac{1}{(2r-1)(2r+1)(2r+3)}$$

$$= \lim_{n \rightarrow \infty} \frac{8}{4} \sum \left(\frac{1}{(2n-1)(2n+1)} - \frac{1}{(2n+1)(2n+3)} \right)$$

$$= \lim_{n \rightarrow \infty} 2 \left[\left(\frac{1}{1.3} - \frac{1}{3.5} \right) + \left(\frac{1}{3.5} - \frac{1}{5.7} \right) + \dots \right]$$

$$= \frac{2}{3}$$

Q12.

(2)

$$S_n = \sum_{k=1}^n \frac{4}{K^2 + 5k + 6}$$

$$= \sum_{k=1}^n \frac{4}{(K+2)(K+3)} = 4 \sum_{K=1}^n \left(\frac{1}{K+2} - \frac{1}{K+3} \right)$$

$$= 4 \left[\frac{1}{3} - \frac{1}{4} \right]$$

$$= 4 \left[\frac{1}{4} - \frac{1}{5} \right]$$

$$= 4 \left[\frac{1}{n+2} - \frac{1}{n+3} \right]$$

$$S_n = 4 \left[\frac{1}{3} - \frac{1}{n+3} \right]$$

$$S_{2025} = 4 \left[\frac{1}{3} - \frac{1}{2028} \right]$$

$$S_{2025} = 4 \left[\frac{675}{2028} \right]$$

$$507 S_{2025} = 675$$

Q13. (4)
$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{k^3 + 6k^2 + 11k + 5}{(k+3)!}$$

$$= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{k^3 + 6k^2 + 11k + 6 - 1}{(k+3)!}$$

$$= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{(k+1)(k+2)(k+3) - 1}{(k+3)!}$$

$$= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{(k+1)(k+2)(k+3)}{(k+3)!} - \frac{1}{(k+3)!}$$

$$= \lim_{k=1}^n \sum_{k=1}^n \left(\frac{1}{k!} - \frac{1}{(k+3)!} \right)$$

$$= \lim_{k=1}^n \left(\frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} \dots + \frac{1}{n!} - \frac{1}{4!} - \frac{1}{5!} - \frac{1}{6!} \dots - \frac{1}{(n+3)!} \right)$$

$$= \frac{1}{1} + \frac{1}{2} + \frac{1}{6} = \frac{10}{6} = \frac{5}{3}$$

Q14. Let the A.P. be

(2) $a, a+2, a+2d, \dots, a+(2k-1)d$

Now, $a + a + 2d + a + 4d + \dots + a + (2k-2)d = 40$

$ka + 2d + 4d + \dots + (2k-2)d = 40$

$\Rightarrow ka + \frac{k-1}{2} [2d + 2kd - 2d] = 40$

$\Rightarrow ka + k(k-1)d = 40 \dots (1)$

And $a + d + a + 3d + \dots + a + (2k-1)d = 55$

$\Rightarrow ka + \frac{k}{2} (d + 2kd - d) = 55$

$\Rightarrow ka + k^2d = 55 \dots (2)$

Also, $a + (2k-1)d - a = 27$

$\Rightarrow (2k-1)d = 27 \Rightarrow d = \frac{27}{2k-1} \dots (3)$

From equation (1) and (2)

$k^2d - kd - k^2d = -15$

$\Rightarrow d = \frac{15}{k}$

From equation (3) and (4)

$\frac{27}{2k-1} = \frac{15}{k}$

$27k = 30k - 15$

$\Rightarrow 3k = 15$

$\Rightarrow k = 5$

Sequences and Series

JEE Main 2025 January

Q15. $S_n = \frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} \dots N \text{ terms}$

(4)
$$S_{2025} = \sum_{n=1}^{2025} \frac{1}{n(n+1)} = \sum_{n=1}^{2025} \left(\frac{1}{n} - \frac{1}{n+1} \right)$$
$$= \left(\frac{1}{1} - \frac{1}{2} \right) + \left(\frac{1}{2} - \frac{1}{3} \right) \dots \dots \left(\frac{1}{2025} - \frac{1}{2026} \right)$$
$$= \frac{2025}{2026}$$

$$\sqrt{2026} \cdot S_{2025} = \sqrt{2025} = 45$$

Given : $\frac{6}{2}[-2p + (6-1)p] = 45$

$$9p = 45$$

$$p = 5$$

$$|A_{20} - A_{15}| = |-5 + 19 \times 5| - [-5 + 14 \times 5]$$
$$= |90 - 65|$$
$$= 25$$

Q16. $a_0 = 0, a_1 = \frac{1}{2}$

(2) $2a_{n+2} = 5a_{n+1} - 3a_n$
$$2x^2 - 5x + 3 = 0 \Rightarrow x = 1, 3/2$$

$$\therefore a_n = A1^n + B\left(\frac{3}{2}\right)^n$$

$$\begin{array}{l} n=0 \quad 0 = A + B \\ n=1 \quad \frac{1}{2} = A + \frac{3}{2}B \end{array} \quad \left. \begin{array}{l} A = -1 \\ B = 1 \end{array} \right\}$$

$$\Rightarrow a_n = -1 + \left(\frac{3}{2}\right)^n$$

$$\sum_{k=1}^{100} a_k = \sum_{k=1}^{100} (-1) + \left(\frac{3}{2}\right)^k$$

$$= -100 + \frac{\left(\frac{3}{2}\right) \left(\left(\frac{3}{2}\right)^{100} - 1 \right)}{\frac{3}{2} - 1}$$

$$= -100 + 3 \left(\left(\frac{3}{2}\right)^{100} - 1 \right)$$

$$= 3 \cdot (a_{100}) - 100$$

Permutation Combination

JEE Main 2025 January

Q1. (3) A, K, N, P, R, U

$$\boxed{A} \dots\dots\dots 5! = 120$$

$$\boxed{K} \dots\dots\dots 5! = 120$$

$$\boxed{N} \dots\dots\dots 5! = 120$$

$$\boxed{P} \boxed{A} \dots\dots\dots 4! = 24$$

$$\boxed{P} \boxed{K} \dots\dots\dots 4! = 24$$

$$\boxed{P} \boxed{N} \dots\dots\dots 4! = 24$$

$$\boxed{P} \boxed{R} \boxed{A} \dots\dots\dots 3! = 6$$

$$\boxed{P} \boxed{R} \boxed{K} \boxed{A} \boxed{N} \boxed{U} = 1$$

$$\boxed{P} \boxed{R} \boxed{K} \boxed{A} \boxed{U} \boxed{N} = 1$$

Total = 440

$\Rightarrow 440^{\text{th}}$ word is P R K A U N

Q2. (3)

Group A		Group B		
B	G	B	G	Ways
4	1	0	3	${}^7C_4 \cdot {}^3C_1 \cdot {}^6C_0 \cdot {}^5C_3$
3	2	1	2	${}^7C_3 \cdot {}^3C_2 \cdot {}^6C_1 \cdot {}^5C_2$
2	3	2	1	${}^7C_2 \cdot {}^3C_3 \cdot {}^6C_2 \cdot {}^5C_1$

$$\text{Total ways} = 30 \cdot {}^7C_4 + 180 \cdot {}^7C_3 + 75 \cdot {}^7C_2 = 8925$$

Permutation Combination

JEE Main 2025 January

Q3. (4) Case I 5 0

Case II 5

5 1

5 3

6 0

6 1

6 2

7 0

7 1

$9 \times (8 \times 8 \times 8) = 4608$ but 50000 is not included, so total numbers $4608 - 1 = 4607$

Q4. Let the number be

(64) $2ab, a + b = 13$

$\Rightarrow a, b \in \{0, 9\}$

$\Rightarrow 6$ numbers $\{(9, 4), (8, 5) \dots (4, 9)\}$

Similarly, for $3ab, a + b = 12 \Rightarrow 7$ numbers

For $4ab, a + b = 11 \Rightarrow$ Numbers

For $5ab, a + b = 10 \Rightarrow 9$ numbers

For $6ab, a + b = 9 \Rightarrow 10$ numbers

For $7ab, a + b = 8 \Rightarrow 9$ numbers

For $8ab, a + b = 7 \Rightarrow 8$ numbers

For $9ab, a + b = 6 \Rightarrow 7$ numbers

$\therefore$ Total ways = 64.

Q5. (4) (i) number of numbers created using

$$1111133 = \frac{7!}{5!2!} \Rightarrow 21$$

(ii) number of numbers created using

$$1111223 = \frac{7!}{4!2!} \Rightarrow 105$$

(iii) number of numbers created using

$$1112222 = \frac{7!}{4!3!} \Rightarrow 35$$

Total = 161

Q6. (i) Single letter is used, then no. of words = 5

(1405) (ii) Two distinct letters are used, then no. of words

$${}^5C_2 \times \left(\frac{6!}{2!4!} \times 2 + \frac{6!}{3!3!} \right) = 10(30 + 20) = 500$$

(iii) Three distinct letters are used, then no. of words

$${}^5C_3 \times \frac{6!}{2!2!2!} = 900$$

Total no. of words = 1405

Q7. (1) $\underbrace{AB}_{12} \quad \underbrace{MN \dots \dots Z}_{13}$

$$= {}^{12}C_2 \times {}^{13}C_2 = 5148$$

Selection of two letters before M

Selection of two letter after M

Permutation Combination

JEE Main 2025 January

Q8. (2) Total - when B_1 and B_2 are together
 $= 2!(3!4!) - 2!(3!(3!2!)) = 144$

Q9. (1) DAUGHTER

Total words $= 8!$

Total words in which vowels are together $= 6! \times 3!$ words in which all vowels are not together
 $= 8! - 6! \times 3!$
 $= 6![56 - 6]$
 $= 720 \times 50$
 $= 36000$

Q10. A : number of ways that all boys sit together $= 5! \times 5!$

(17280) B : number of ways if no 2 boys

sit together $= 4! \times 5!$

$A \cap B = \phi$

Required no. of ways $= 5! \times 5! + 4! \times 5! = 17280$



Binomial Theorem

JEE Main 2025 January

Q1. The general term of multinomial expansion is

$$(612) \frac{6!}{\alpha! \beta! \gamma!} (1)^\alpha \left(2\frac{1}{3}\right)^\beta \left(3\frac{1}{2}\right)^\gamma$$

For terms to be rational $3 \mid \beta$ and $2 \mid \gamma$

β	γ	α	Term
0	0	6	1
0	2	4	$15 \cdot 3 = 45$
0	4	2	$15 \cdot 3^2 = 135$
0	6	0	$1 \cdot 3^3 = 27$
3	0	3	$20 \cdot 2 = 40$
3	2	1	$60 \cdot 2 \cdot 3 = 360$
6	0	0	$1 \cdot 4 = 4$

$\Rightarrow$ Sum of rational terms

$$= 1 + 45 + 135 + 27 + 40 + 360 + 4 = 612$$

Q2. $(1+x)^{n+4}$

(3) ${}^{n+4}C_4, {}^{n+4}C_5, {}^{n+4}C_6, \rightarrow$ A.P.

$$\Rightarrow 2 \times {}^{n+4}C_5 = {}^{n+4}C_4 + {}^{n+4}C_6$$

$$\Rightarrow 4 \times {}^{n+4}C_5 = ({}^{n+4}C_4 + {}^{n+4}C_5) + ({}^{n+4}C_5 + {}^{n+4}C_6)$$

$$\Rightarrow 4 \times {}^{n+4}C_5 = {}^{n+5}C_5 + {}^{n+5}C_6$$

$$\Rightarrow 4 \times \frac{(n+4)!}{5! \cdot (n-1)!} = \frac{(n+6)!}{6! \cdot n!}$$

$$\Rightarrow 4 = \frac{(n+6)(n+5)}{6n}$$

$$\Rightarrow n^2 + 11n + 30 = 24n$$

$$\Rightarrow n^2 - 13n + 30 = 0$$

$$\Rightarrow n = 3, 10 \text{ (rejected)}$$

$$\therefore n \neq 10$$

$\therefore$ Largest binomial coefficient in expansion of

$$(1+x)^7$$

$$(\because n+4=7)$$

is coeff. of middle term

$$\Rightarrow {}^7C_4 = {}^7C_3 = 35$$

Binomial Theorem

JEE Main 2025 January

Q3. $A = {}^{2n-1}C_{29}$ $B = {}^{2n-1}C_{11}$

(3) $2^{2n-1}C_{29} = 5^{2n-1}C_{11}$

$$2 \frac{(2n-1)!}{29!(2n-30)!} = 5 \frac{(2n-1)!}{(2n-12)!11!}$$

$$\frac{1}{29 \dots 12 \cdot 5} = \frac{1}{(2n-12)(2n-13) \dots (2n-29)2}$$

$$\frac{1}{30 \cdot 29 \dots 12} = \frac{1}{(2n-12)(2n-13) \dots (2n-29)12}$$

$$2n-12=30$$

$$n=21$$

Q4. $(x + \sqrt{x^3-1})^5 + (x - \sqrt{x^3-1})^5$

(1)
$$- \left[{}^5C_0 x^5 + {}^5C_1 x^4 (\sqrt{x^3-1}) + \dots + {}^5C_5 (\sqrt{x^3-1})^5 \right] +$$

$$\left[{}^5C_0 x^5 - {}^5C_1 x^4 (\sqrt{x^3-1}) + \dots - {}^5C_5 (\sqrt{x^3-1})^5 \right]$$

$$= 2 \left[x^5 + {}^5C_2 x^3 (x^3-1) + {}^5C_4 (x^3-1)^2 \right]$$

$$= 10x^7 + 20x^6 + 2x^5 - 20x^4 - 20x^3 + 10x$$

Now

$$\alpha = 10, \beta = 2, \gamma = -20, \delta = 10$$

Also,

$$\left. \begin{array}{l} 10u + 2v = 18 \\ -20u + 10v = 20 \end{array} \right\} u = 1, v = 4$$

$$u + v = 5$$

Q5. $(1+x)^p(1-x)^q = ({}^pC_0 + {}^pC_1x + {}^pC_2x^2 + \dots)({}^qC_0 - {}^qC_1x + {}^qC_2x^2 + \dots)$

(2) $\text{coff of } x \equiv {}^pC_0 {}^qC_1 + {}^pC_1 {}^qC_0 = 1$

$$p - q = 1$$

$$\text{coff of } x^2 \equiv {}^pC_0 {}^qC_2 - {}^pC_1 {}^qC_1 + {}^pC_2 {}^qC_0 = -2$$

$$\frac{q(q-1)}{2} - pq + \frac{p(p-1)}{2} = -2$$

$$q^2 - q - 2pq + p^2 - p = -4$$

$$1 - (p+q) = -4$$

$$p+q=5$$

$$p=3$$

$$q=2$$

$$\text{so } p^2 + q^2 = 13$$

Q6. Coefficient of

(1) $T_r, T_{r+1}, T_{r+2} \rightarrow GP$

$$\Rightarrow ({}^{12}C_r)^2 = {}^{12}C_{r-1} \cdot {}^{12}C_{r+1}$$

$$\Rightarrow ({}^{12}C_r)^2 = {}^{12}C_{r-1} \cdot {}^{12}C_{r+1}$$

but no three consecutive binomial coefficient are in

Binomial Theorem

JEE Main 2025 January

GP

$$\Rightarrow P = 0$$

$$\text{Now for } \left(3^{1/4} + 4^{1/3}\right)^{12}, T_{r+1} = {}^{12}C_r (4)^{K/3} (3)^{\frac{12-K}{4}}$$

for rational terms $K = 0, 12$

sum of rational terms

$$= {}^{12}C_0 4^0 \cdot 3^3 + {}^{12}C_{12} \cdot 4^4 \cdot 3^0$$

$$= 27 + 256 = 283 = q$$

$$\therefore p + q = 283$$

Q7. General term $= {}^nC_r \left\{7^{1/3}\right\}^{n-r} \left(11^{1/12}\right)^r$

(1) $= {}^nC_r \left\{7\right\}^{\frac{n-r}{3}} (11)^{r/12}$

For integral terms, r must be multiple of 12

$$\therefore r = 12k, k \in W$$

Total values of $r = 183$

$$\text{Hence max } r = 12(182)$$

$$= 2184$$

Min value of $n = 2184$

Q8. $7^{103} = 7(7^{102}) = 7(343)^{34} = 7(345 - 2)^{34}$

(4) $7^{103} = 23K_1 + 7 \cdot 2^{34}$

$$\text{Now } 7 \cdot 2^{34} = 7 \cdot 2^2 \cdot 2^{32}$$

$$= 28 \cdot (256)^4$$

$$= 28(253 + 3)^4$$

$$\therefore 28 \times 81 \Rightarrow (23 + 5)(69 + 12)$$

$$23K_2 + 60$$

$$\therefore \text{Remainder} = 14$$

Q9. $(1+x)^{11} = {}^{11}C_0 + {}^{11}C_1x + {}^{11}C_2x^2 + \dots + {}^{11}C_{11}x^{11}$

(2035) $\int_0^1 (1+x)^{11} dx = \int_0^1 ({}^{11}C_0 + {}^{11}C_1x + {}^{11}C_2x^2 + \dots + {}^{11}C_{11}x^{11}) dx$

$$\left[\frac{(1+x)^{12}}{12} \right]_0^1 = {}^{11}C_0x + \frac{{}^{11}C_1x^2}{2} + \frac{{}^{11}C_2x^3}{3} + \dots + \frac{{}^{11}C_{11}x^{12}}{12} \Bigg|_0^1$$

$$\frac{2^{12} - 1}{12} = C_0 + \frac{C_1}{2} + \frac{C_2}{3} + \dots + \frac{C_{11}}{12} \dots (1)$$

Now,

Binomial Theorem

JEE Main 2025 January

$$\int_{-1}^0 (1+x)^{11} dx = \int_{-1}^0 ({}^{11}C_0 + {}^{11}C_1 x + {}^{11}C_2 x^2 + \dots + {}^{11}C_{11} x^{11}) dx$$

$$\left[\frac{(1+x)^{12}}{12} \right]_{-1}^0 = {}^{11}C_0 x + \frac{{}^{11}C_1 x^2}{2} + \frac{{}^{11}C_2 x^3}{3} + \dots + \frac{{}^{11}C_{11} x^{12}}{12} \Bigg]_{-1}^0$$

$$\frac{1}{12} = C_0 - \frac{C_1}{2} + \frac{C_2}{3} \dots$$

$$(1) - (2)$$

$$= \frac{2^{12} - 2}{12} = 2 \left[\frac{C_1}{2} + \frac{C_3}{4} + \dots \right]$$

$$\Rightarrow \sum_{r=0}^5 \frac{C_{2r+1}}{2r+2} = \frac{2^{11} - 1}{12} = \frac{2047}{12} = \frac{m}{n}$$

$$= 2047 - 12 = 2035$$

Q10.

(5)

$$\alpha = 1 + \sum_{r=1}^6 (-1)^{r-1} {}^{12}C_{2r-1} 3^{r-1}$$

$$\alpha = 1 + \sum_{r=1}^6 {}^{12}C_{2r-1} \frac{(\sqrt{3}i)^{2r-1}}{\sqrt{3}i} \quad i = \text{iota, let } \sqrt{3}i = x$$

$$\alpha = 1 + \frac{1}{\sqrt{3}i} ({}^{12}C_1 x + {}^{12}C_3 x^3 + \dots + {}^{12}C_{11} x^{11})$$

$$= 1 + \frac{1}{\sqrt{3}i} \left(\frac{(1 + \sqrt{3}i)^{12} - (1 - \sqrt{3}i)^{12}}{2} \right)$$

$$= 1 + \frac{1}{\sqrt{3}i} \left(\frac{(-2w^2)^{12} - (2w)^{12}}{2} \right) = 1$$

so distance of $(12, \sqrt{3})$ from $x - \sqrt{3}y + 1 = 0$ is

$$\frac{12 - 3 + 1}{2} = 5$$

Q11.

(465)

$$\sum_{r=1}^{30} \frac{r^2 ({}^{30}C_r)^2}{{}^{30}C_{r-1}}$$

$$= \sum_{r=1}^{30} r^2 \left(\frac{31-r}{r} \right) \cdot \frac{30!}{r!(30-r)!}$$

$$\left(\because \frac{{}^{30}C_r}{{}^{30}C_{r-1}} = \frac{30-r+1}{r} = \frac{31-r}{r} \right)$$

Binomial Theorem

JEE Main 2025 January

$$\begin{aligned}&= \sum_{r=1}^{30} \frac{(31-r)30!}{(r-1)!(30-r)!} \\&= 30 \sum_{r=1}^{30} \frac{(31-r)29!}{(r-1)!(30-r)!} \\&= 30 \sum_{r=1}^{30} (30-r+1)^{29} C_{30-r} \\&= 30 \left(\sum_{r=1}^{30} (31-r)^{29} C_{30-r} + \sum_{r=1}^{30} {}^{29}C_{30-r} \right) \\&= 30 (29 \times 2^{28} + 2^{29}) = 30(29+2)2^{28} \\&= 15 \times 31 \times 2^{29} \\&= 465 (2^{29}) \\&\alpha = 465\end{aligned}$$



Q1. $8 + (n - 1)13 = 320$

(8788) $13n = 325$

$n = 25$

no. of terms = 25

$$\text{mean} = \frac{\sum x_i}{n} = \frac{8 + 21 + \dots + 320}{25} = \frac{\frac{25}{2}(8 + 320)}{25}$$

$$\text{variance } \sigma^2 = \frac{\sum x_i^2}{n} - (\text{mean})^2$$

$$= \frac{8^2 + 21^2 + \dots + 320^2}{13} - (164)^2$$

$$= 8788$$

Q2. (3) median $= \ell + \left(\frac{\frac{N}{2} - F}{f} \right) \times h$

$$= 12 + \left(\frac{\frac{N}{2} - 18}{12} \right) \times 6 = 14$$

$$\Rightarrow \left(\frac{\frac{N}{2} - 18}{12} \right) \times 6 = 2$$

$$\frac{N}{2} - 18 = 4 \Rightarrow N = 44$$

Q3. Mean $\bar{x} = 5.5$

(3) $= \sum_{i=1}^{10} x_i = 5.5 \times 10 = 55$

$$= \sum_{i=1}^{10} x_i^2 = 371$$

$$\left(\sum x_i \right)_{\text{new}} = 55 - (4 + 5) + (6 + 8) = 60$$

$$\left(\sum x_i \right)_{\text{new}} = 371 - (4^2 + 5^2) + (6^2 + 8^2) = 430$$

$$\text{Variance } \sigma^2 = \frac{\sum x_i^2}{10} - \left(\frac{\sum x_i}{10} \right)^2$$

$$\sigma^2 = \frac{430}{10} - \left(\frac{60}{10} \right)^2$$

$$\sigma^2 = 43 - 36$$

$$\sigma^2 = 7$$

Q4. (1) $\sum_{l=1}^{10} (x_l - 2) = 30$

$$\sum_{i=1}^{10} x_l = 50$$

$$\Rightarrow \text{Mean} = 5$$

$$\text{Variance} = \frac{4}{5} = \frac{\sum x_l^2}{10} - (\bar{x})^2$$

$$\frac{4}{5} = \frac{\sum x_l^2}{10} - 25$$

$$\Rightarrow \sum x_l^2 = 258$$

$$\text{Now, } \sum_{l=1}^{10} (x_l - \beta)^2 = 98$$

$$\sum_{l=1}^{10} x_l^2 - 2\beta \sum_{l=1}^{10} x_l + 10\beta^2 = 98$$

$$\Rightarrow 258 - 2\beta(50) + 10\beta^2 = 98$$

$$\Rightarrow 10\beta^2 - 100\beta + 160 = 0$$

$$\Rightarrow \beta^2 - 10\beta + 16 = 0$$

$$\Rightarrow \beta = 8 \text{ as } \beta > 2$$

Now, as per the question

$$2(x_1 - 1) + 4\beta, 2(x_2 - 1) + 4\beta, \dots, 2(x_{10} - 1) + 4\beta$$

Can be simplified as

$$2x_1 + 30, 2x_2 + 30, \dots, 2x_{10} + 30$$

$$\mu = 2(5) + 30 = 40$$

$$\sigma^2 = 2^2 \left(\frac{4}{5} \right) = \frac{16}{5}$$

$$\frac{\beta\mu}{\sigma^2} = \frac{8 \times 40}{\frac{16}{5}} = 100$$

Q1.

(1) Let $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$

$$\therefore \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\therefore b = 0, e = 0, h = 1$$

and $\begin{bmatrix} a & 0 & c \\ d & 0 & f \\ g & 1 & i \end{bmatrix} \begin{bmatrix} 4 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

$$\left. \begin{array}{l} 4a + 3c = 0 \\ 4d + 3f = 1 \\ 4g + 1 + 3i = 0 \end{array} \right\} \dots (1)$$

and $\begin{bmatrix} a & 0 & c \\ d & 0 & f \\ g & 1 & i \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

$$\left. \begin{array}{l} 2a + 2c = 1 \\ 2d + 2f = 0 \\ 2g + 1 + 2i = 0 \end{array} \right\} \dots (2)$$

From equation (1) and (2) we get

$$d = 1, f = -1$$

$$\therefore a_{23} = -1$$

Q2.

(2) $A = \begin{bmatrix} (\sqrt{2})^2 & (\sqrt{2})^3 & (\sqrt{2})^4 \\ (\sqrt{2})^3 & (\sqrt{2})^4 & (\sqrt{2})^5 \\ (\sqrt{2})^4 & (\sqrt{2})^5 & (\sqrt{2})^6 \end{bmatrix}$

$$A = \begin{bmatrix} 2 & 2\sqrt{2} & 4 \\ 2\sqrt{2} & 4 & 4\sqrt{2} \\ 4 & 4\sqrt{2} & 8 \end{bmatrix}$$

$$A^2 = 2^2 \begin{bmatrix} 1 & \sqrt{2} & 2 \\ \sqrt{2} & 2 & 2\sqrt{2} \\ 2 & 2\sqrt{2} & 4 \end{bmatrix} \begin{bmatrix} 1 & \sqrt{2} & 2 \\ \sqrt{2} & 2 & 2\sqrt{2} \\ 2 & 2\sqrt{2} & 4 \end{bmatrix}$$

$$= 4 \begin{bmatrix} - & - & - \\ - & - & - \\ (2+4+8) & (2\sqrt{2}+4\sqrt{2}+8\sqrt{2}) & (4+8+16) \end{bmatrix}$$

$$\text{Sum of elements of 3}^{\text{rd}} \text{ row} = 4(14 + 14\sqrt{2} + 28)$$

$$= 4(42 + 14\sqrt{2})$$

$$= 168 + 56\sqrt{2}$$

$$\alpha + \beta\sqrt{2}$$

$$\therefore \alpha + \beta = 168 + 56 = 224$$

Q3. $\therefore \text{tr}(A) = 3$ and $|A| = \frac{1}{2}$

(4) Now, $B = \text{adj}(\text{adj}(2A)) = |2A|^{3-2} \cdot (2A)$

Matrices

JEE Main 2025 January

$$= 2^3 |A| \cdot 2A = 8A$$

$$\therefore \text{tr}(B) = 8 \text{tr}(A) = 24$$

$$\text{and } |B| = |8A| = 8^3 \cdot \frac{1}{2} = 256$$

$$\therefore \text{trace}(B) + |B| = 24 + 256 = 280$$

Q4.
$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

(1613)

$$\text{No. of elements in } S_1 : A = A^T \Rightarrow 5^3 \times 5^3$$

$$\text{No. of elements in } A = -A^T \Rightarrow 0$$

since no. zero in 5

$$\text{No. of elements in } S_3 \Rightarrow$$

$$\left. \begin{array}{l} a_{11} + a_{22} + a_{33} = 0 \Rightarrow (1, 2, -3) \Rightarrow 31 \\ \text{or} \\ (1, 1, -2) \Rightarrow 3 \\ \text{or} \\ (-1, -1, 2) \Rightarrow 3 \end{array} \right\} \Rightarrow 12 \times 5^6$$

$$n(S_1 \cap S_3) = 12 \times 5^3$$

$$n(S_1 \cup S_2 \cup S_3) = 5^6(1 + 12) - 12 \times 5^3$$

$$\Rightarrow 5^3 \times [13 \times 5^3 - 12] = 125\alpha$$

$$\alpha = 1613$$

Q5.
$$A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$$

(2)

Now finding characteristic equation

$$\begin{vmatrix} 2 - \lambda & -1 \\ 1 & -\lambda \end{vmatrix} = 0$$

$$\Rightarrow (2 - \lambda)(-\lambda) - (-1)(1) = -2\lambda + \lambda^2 + 1 = 0$$

$$\Rightarrow \lambda^2 - 2\lambda + 1 = 0$$

$$\Rightarrow (\lambda - 1)^2 = 0$$

$$\Rightarrow \lambda = 1$$

$$\text{Since } A \text{ satisfies } (A - I)^2 = 0$$

$$\therefore A = I + N \text{ where}$$

$$N = A - I$$

$$N = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$$

$$N^2 = 0$$

$$A^m = (I + N)^m = I + mN$$

$$A^m \cdot A^m = (I + mN)(I + mN) = I + 2mN + m^2N^2$$

$$\text{Since } N^2 = 0$$

$$\Rightarrow A^{m^2} = I + 2mN$$

Now putting in given condition

Matrices

JEE Main 2025 January

$$I + m^2 N + I + mN = 3I - A^{-6}$$

$$A^{-1} = \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix}$$

$$A^{-6} = (A^{-1})^6 = I + (-6)N$$

$\therefore$ Putting in (i)

$$(m^2 + m)N = I - (I - 6N)$$

$$(m^2 + m)N = 6N$$

Since $N \neq 0$

$$\Rightarrow m^2 + m = 6$$

$$\Rightarrow m^2 + m - 6 = 0$$

$$\Rightarrow (m - 2)(m + 3) = 0$$

$$\Rightarrow m = 2, -3$$

$\therefore$ Number of elements in S is 2

Q6. As $A \operatorname{adj} A = |A|I$, $\det(\lambda A) = \lambda^n \det A$

$$\begin{aligned} (34) \quad \det(3 \operatorname{adj}(-6 \operatorname{adj}(3A))) &= 3^3 \det(\operatorname{adj}(-6 \operatorname{adj}(3A))) \\ &= 3^3 (-6 \operatorname{adj}(3A))^2 \\ &= 3^3 (-6)^6 |3A|^4 \\ &= 3^9 2^6 \cdot 3^{12} \cdot (-2)^4 \\ &= 3^{21} \cdot 2^{10} \end{aligned}$$

Now comparing with given condition

$$2^{m+n} 3^{mn} = 2^{10} \cdot 3^{21}$$

$$m + n = 10, mn = 21$$

$$\Rightarrow m = 7, n = 3 (m > n)$$

$$\therefore 4m + 2n = 28 + 6 = 34$$

Q7. $X^T A X = 0$

(44) $(xyz) \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = 0$

$$(xyz) \begin{bmatrix} a_1 x + a_2 y + a_3 z \\ b_1 x + b_2 y + b_3 z \\ c_1 x + c_2 y + c_3 z \end{bmatrix} = 0$$

$$x(a_1 x + a_2 y + a_3 z) + y(b_1 x + b_2 y + b_3 z) + z(c_1 x + c_2 y + c_3 z)$$

$$a_1 = 0, b_2 = 0, c_3 = 0$$

$$a_2 + b_1 = 0, a_3 + c_1 = 0, b_3 + c_2 = 0$$

A = skew symmetric matrix

$$A = \begin{bmatrix} 0 & x & y \\ -x & 0 & z \\ -y & -z & 0 \end{bmatrix}; A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \\ -5 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & x & y \\ -x & 0 & z \\ -y & -z & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \\ -5 \end{bmatrix}$$

$$x + y = 1$$

$$-x + z = 4$$

$$y + z = 5$$

$$\begin{bmatrix} 0 & x & y \\ -x & 0 & z \\ -y & -z & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \\ -8 \end{bmatrix}$$

$$2x + y = 0 \quad x = -1$$

$$-x + z = 4 \quad y = 2$$

$$-y - 2z = -8 \quad z = 3$$

$$A = \begin{bmatrix} 0 & -1 & 2 \\ 1 & 0 & 3 \\ -2 & -3 & 0 \end{bmatrix}$$

$$2(A + I) = \begin{bmatrix} 2 & -2 & 4 \\ 2 & 2 & 6 \\ -2 & -6 & 2 \end{bmatrix}$$

$$2(A + I) = 120$$

$$\Rightarrow \det(\text{adj}(2A + I))$$

$$= 120^2 = 2^6 \cdot 3^2 \cdot 5^2$$

$$\therefore \alpha = 6, \beta = 2, \gamma = 2$$

$$\text{Hence } \alpha^2 + \beta^2 + \gamma^2 = 6^2 + 2^2 + 2^2 = 44$$

Matrices

JEE Main 2025 January

Q8. $|A| = \frac{11}{2}$

(3) $C_{11} = \sum_{k=1}^2 a_{1k} \cdot A_{1k} = a_{11} A_{11} + a_{12} A_{12} = |A| = \frac{11}{2}$

$C_{12} = \sum_{k=1}^2 a_{1k} \cdot A_{2k} = 0$

$C_{21} = \sum_{k=1}^2 a_{2k} \cdot A_{1k} = 0$

$C_{22} = \sum_{k=1}^2 a_{2k} \cdot A_{2k} = |A| = \frac{11}{2}$

$C = \begin{bmatrix} 11/2 & 0 \\ 0 & 11/2 \end{bmatrix}$

$|C| = \frac{121}{4}$

$8|C| = 242$

Q9. $\left[A \left(\text{adj}(A^{-1}) + \text{adj}(B^{-1}) \right)^{-1} \cdot B \right]^{-1}$

(4) $B^{-1} \cdot \left(\text{adj}(A^{-1}) + \text{adj}(B^{-1}) \right) \cdot A^{-1}$

$B^{-1} \text{adj}(A^{-1}) A^{-1} + B^{-1} \left(\text{adj}(B^{-1}) \right) \cdot A^{-1}$

$B^{-1} |A^{-1}| I + |B^{-1}| I A^{-1}$

$\frac{B^{-1}}{|A|} + \frac{A^{-1}}{|B|}$

$\Rightarrow \frac{\text{adj} B}{|B||A|} + \frac{\text{adj} A}{|A||B|}$

$= \frac{1}{|A||B|} (\text{adj} B + \text{adj} A)$

Q10. $P = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$

(3) $\therefore P^T P = I$

$B = P A P^T$

Pre multiply by P^T (Given)

$P^T B = P^T P A P^T = A P^T$

Now post multiply by P

$P^T B P = A P^T P = A$

So $A^2 = \underbrace{P^T B P}_{I} P^T B P$

$A^2 = P^T B^2 P$

Similarly $A^{10} = P^T B^{10} P = C$

Matrices

JEE Main 2025 January

$$A = \begin{bmatrix} \frac{1}{\sqrt{2}} & -2 \\ 0 & 1 \end{bmatrix} \text{ (Given)}$$

$$\Rightarrow A^2 = \begin{bmatrix} \frac{1}{2} & -\sqrt{2} - 2 \\ 0 & 1 \end{bmatrix}$$

Similarly check A^3 and so on since $C = A^{10} \Rightarrow$ Sum of diagonal elements of C is $\left(\frac{1}{\sqrt{2}}\right)^{10} + 1$

$$= \frac{1}{32} + 1 = \frac{33}{32} = \frac{m}{n}$$

gcd(m, n) = 1 (Given)

$$\Rightarrow m + n = 65$$



Determinants

JEE Main 2025 January

Q1.
$$(1) \begin{vmatrix} 1 + \sin^2 x & \cos^2 x & 4 \sin 4x \\ \sin^2 x & 1 + \cos^2 x & 4 \sin 4x \\ \sin^2 x & \cos^2 x & 1 + 4 \sin 4x \end{vmatrix}, x \in R$$

$$R_2 \rightarrow R_2 - R_1 \& R_3 \rightarrow R_3 - R_1$$

$$f(x) = \begin{vmatrix} 1 + \sin^2 x & \cos^2 x & 4 \sin 4x \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{vmatrix}$$

Expand about R_1 , use get

$$f(x) = 2 + 4 \sin 4x$$

$$\therefore M = \text{max value of } f(x) = 6$$

$$M = \text{min value of } f(x) = -2$$

$$\therefore M^4 - M^4 = 1280$$

Q2.
$$(3) D = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 5 \\ 1 & 5 & \lambda \end{vmatrix} = 0$$

$$\lambda = 17$$

$$D_z = \begin{vmatrix} 1 & 1 & 6 \\ 1 & 2 & 9 \\ 1 & 5 & \mu \end{vmatrix} \neq 0$$

$$\mu \neq 18$$

Q3.
$$(1) \lim_{x \rightarrow 0} \begin{vmatrix} a + \frac{\sin x}{x} & 1 & b \\ a & 1 + \frac{\sin x}{x} & b \\ a & 1 & b + \frac{\sin x}{x} \end{vmatrix} = \lambda + \mu a + vb$$

At $\lim x \rightarrow 0$,

$$f(x) = \begin{vmatrix} a + 1 & 1 & b \\ a & 1 + 1 & b \\ a & 1 & b + 1 \end{vmatrix} = \lambda + \mu a + vb$$

$$R_1 \rightarrow R_1 - R_2$$

$$R_2 \rightarrow R_2 - R_3$$

$$\begin{vmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ a & 1 & b + 1 \end{vmatrix} = \lambda + \mu a + vb$$

$$C_2 \rightarrow C_1 - C_2$$

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ a & a + 1 & b + 1 \end{vmatrix} = \lambda + \mu a + vb$$

$$a + b + 2 = \lambda + \mu a + vb$$

$$\lambda = 2, \mu = 1, v = 1$$

$$(\lambda + \mu + v) = (2 + 1 + 1)^2 = 16$$

Determinants

JEE Main 2025 January

Q4. The given equations are

(1) $x + y + 2z = 6$

$$2x + 3y + az = a + 1$$

$$-x - 3y + bz = 2b, \text{ where } a, b, \in \mathbf{R}$$

For infinite many solutions:

$$D = D_1 = D_2 = D_3 = 0$$

$$\therefore D = \begin{vmatrix} 1 & 1 & 2 \\ 2 & 3 & a \\ -1 & -3 & b \end{vmatrix} = 2a + b - 6$$

$$D_1 = \begin{vmatrix} 6 & 1 & 2 \\ a+1 & 3 & a \\ 2b & -3 & b \end{vmatrix} = 12a + 5b + ab - 6$$

$$D_2 = \begin{vmatrix} 1 & 6 & 2 \\ 2 & a+1 & a \\ -1 & 2b & b \end{vmatrix} = -4a - 3b - ab + 2$$

$$\text{and } D_3 = \begin{vmatrix} 1 & 1 & 6 \\ 2 & 3 & a+1 \\ -1 & -3 & 2b \end{vmatrix} = 2a + 2b - 16$$

from above relations

$$a = -2, b = 10$$

$$\therefore 7a + 3b = 16$$

Q5. $(\lambda - 1)x + (\lambda - 4)y + \lambda z = 5$

(4) $\lambda x + (\lambda - 1)y + (\lambda - 4)z = 7$

$$(\lambda + 1)x + (\lambda + 2)y - (\lambda + 2)z = 9$$

For infinitely many solutions

$$D = \begin{vmatrix} \lambda - 1 & \lambda - 4 & \lambda \\ \lambda & \lambda - 1 & \lambda - 4 \\ \lambda + 1 & \lambda + 2 & -(\lambda + 2) \end{vmatrix} = 0$$

$$(\lambda - 3)(2\lambda + 1) = 0$$

$$D_x = \begin{vmatrix} 5 & \lambda - 4 & \lambda \\ 7 & \lambda - 1 & \lambda - 4 \\ 9 & \lambda + 2 & -(\lambda + 2) \end{vmatrix} = 0$$

$$2(3 - \lambda)(23 - 2\lambda) = 0$$

$$\lambda = 3$$

$$\therefore \lambda^2 + \lambda = 9 + 3 = 12$$

Determinants

JEE Main 2025 January

Q6. $2x - y + z = 4$

(1) $5x + \lambda y + 3z = 12$

$100x - 47y + \mu z = 212$

$\Delta x = \Delta y = \Delta z = 0$

$$\Delta z = \begin{vmatrix} 2 & -1 & 4 \\ 5 & \lambda & 12 \\ 100 & -47 & 212 \end{vmatrix} = 0$$

$$\Rightarrow 2(212\lambda + 564) + 1(1060 - 1200)$$

$$\Rightarrow 424\lambda + 1128 - 140 - 940 - 400\lambda = 0$$

$$\Rightarrow \lambda = -2$$

$$\Delta y = \begin{vmatrix} 2 & -1 & 1 \\ 5 & -2 & 3 \\ 100 & -47 & \mu \end{vmatrix} = 0$$

$$\Rightarrow 2(-2\mu + 141) + 1(5\mu - 300) + 1(-235 + 200) = 0$$

$$\Rightarrow \mu = 53$$

$$\mu - 2\lambda = 53 - 2(-2) = 57$$

Q7. $\Delta = \begin{vmatrix} 1 & 2 & -3 \\ 2 & \lambda & 5 \\ 14 & 3 & \mu \end{vmatrix} = 0 \Rightarrow \lambda\mu + 42\lambda - 4\mu + 107 = 0$

$$\Delta_1 = \begin{vmatrix} 2 & 2 & -3 \\ 5 & \lambda & 5 \\ 33 & 3 & \mu \end{vmatrix} = 0 \Rightarrow 2\lambda\mu + 99\lambda - 10\mu + 255 = 0$$

$$\Delta_2 = \begin{vmatrix} 1 & 2 & -3 \\ 2 & 5 & 5 \\ 14 & 33 & \mu \end{vmatrix} = 0 \Rightarrow \mu = 13$$

Also, $\lambda = -1$

Hence, $\lambda + \mu = 13 - 1 = 12$

Q8. $\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 4 & 10 \end{vmatrix} = 1(20 - 16) - 1(10 - 4) + 1(4 - 2) = 4 - 6 + 2 = 0$

For infinite solutions

$\Delta_x = \Delta_y = \Delta_z = 0$

$m^2 - 3x + 2 = 0$

$m = 1, 2$

$\alpha = 1, \beta = 2$

$$\begin{aligned} \therefore \sum_{n=1}^{10} (n^\alpha + n^\beta) &= \sum_{n=1}^{10} n^1 + \sum_{n=1}^{10} n^2 \\ &= \frac{10(11)}{2} + \frac{10(11)(21)}{6} \\ &= 55 + 385 \\ &= 440 \end{aligned}$$

Probability

JEE Main 2025 January

Q1. $A = [a_{ij}]_{2 \times 2}$ and entries are 0 or 1.

(3) $\therefore \begin{vmatrix} a & b \\ c & d \end{vmatrix} = 0$

$$\Rightarrow ad - bc = 0$$

Case I: $ad = bc = 1$

$$\therefore a = b = c = d = 1$$

Case II: $ad = bc = 0$

$$a = 0, d = 0 \quad b = 0, c = 0$$

$$a = 0, d = 1 \quad b = 0, c = 1$$

$$a = 1, d = 0 \quad b = 1, c = 0$$

$\therefore$ Total 10 cases when matrix is non invertible Total possible matrix $= 2^4 = 16$

Required probability of invertible

$$= \frac{16-10}{16} = \frac{6}{16} = \frac{3}{8}$$

Q2. Let $k_1 = 4\lambda_1 + r_1, r_1 \in \{0, 1, 2, 3\}$

(2) $k_2 = 4\lambda_2 + r_2$

$$(i)^{k_1} + (i)^{k_2} = (i)^{r_1} + (i)^{r_2}$$

$$(i)^{r_1} \in \{1, i, -1, -i\}$$

$$\text{Zero} \Rightarrow 1, (-1) \text{ pair} \Rightarrow \left\{ \begin{array}{l} 1, -1 \\ i, -i \\ -i, +i \\ -1, 1 \end{array} \right\}$$

$i, (-i)$ pair

$$\text{Zero probability} = \frac{4}{{}^4C_1 \cdot {}^4C_1} = \frac{1}{4}$$

$$\text{Probability (non-zero)} = 1 - \frac{1}{4} = \frac{3}{4}$$

Q3. A, E, G, R, D, N

(1) Probability (P) $= \frac{\text{favourable case}}{\text{Total case}}$

(when A & E are in order)

Total case $= 6!$

Favourable case $= {}^6C_2 \cdot 4!$

$$P = \frac{(15)4!}{(30)4!}$$

$$\text{Probability when not in order} = 1 - \frac{1}{2} = \frac{1}{2}$$

Q4. Bag contains 4 white and 6 black balls

(2) A : first ball selected is black

B : Second ball is also black

$$P\left(\frac{A}{B}\right) = \frac{\frac{6}{10} \times \frac{5}{9}}{\frac{4}{10} \times \frac{6}{9} + \frac{6}{10} \times \frac{5}{9}} = \frac{30}{24 + 30}$$

$$= \frac{30}{54} = \frac{5}{9}$$

$$m + n = 5 + 9 = 14$$

Q5.

(4)

	x_i	P_i
HHH	0	$\frac{1}{8}$
TTT	0	$\frac{1}{8}$
HHT	1	$\frac{1}{8}$
HTH	1	$\frac{1}{8}$
THH	0	$\frac{1}{8}$
TTH	0	$\frac{1}{8}$
THT	1	$\frac{1}{8}$
HTT	1	$\frac{1}{8}$

$$\mu = \sum x_i P_i = \frac{1}{2}$$

$$\sigma^2 = \sum x_i^2 P_i - \mu^2$$

$$= \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$$

$$64(\mu + \sigma^2) = 64\left[\frac{1}{2} + \frac{1}{4}\right]$$

$$= 64 \times \frac{3}{4} = 48$$

Q6. a = number of dice 1

(2) b = number on dice 2

$$(a, b) = (1, 3), (3, 1), (2, 2), (2, 3), (3, 2), (1, 4), (4, 1)$$

Required probability

$$= \frac{2}{6} \times \frac{2}{6} + \frac{1}{6} \times \frac{1}{6} + \frac{2}{6} \times \frac{2}{6} + \frac{2}{6} \times \frac{2}{6} + \frac{1}{6} \times \frac{2}{6} + \frac{2}{6} \times \frac{1}{6} + \frac{1}{6} \times \frac{2}{6}$$


$$= \frac{18}{36} = \frac{1}{2}$$

Probability

JEE Main 2025 January

Q7.

(2)



$$\text{Total} = {}^{16}C_2$$

$$\text{Required ways} = \text{Total} - (\text{adjacent square})$$

$$= {}^{16}C_2 -$$

[3 pair in vertical \& horizontal for each row and column]

$$= {}^{16}C_2 - [3 \times 4 + 3 \times 4]$$

$$= 96$$

$$\text{Probability} = \frac{96}{120} = \frac{4}{5}$$

Q8. For sum '5' $\rightarrow (1, 4), (2, 3), (3, 2)$

(2) $(4, 1) \Rightarrow P(A) = \frac{4}{36}$

For sum '8' $\rightarrow (2, 6), (3, 5), (4, 4)$

For sum '5' $\rightarrow (1, 4), (2, 3), (3, 2)$

$(4, 1) \Rightarrow P(A) = \frac{4}{36}$

For sum '8' $\rightarrow (2, 6), (3, 5), (4, 4)$

$(5, 3), (6, 2) \Rightarrow P(B) = \frac{5}{36}$

$P(\bar{A}) = \frac{32}{36}, P(\bar{B}) = \frac{31}{36}$

$P(A \text{ wins}) = P(A) + P(\bar{A})P(\bar{B})P(A) +$

$+ P(\bar{A})P(\bar{B})P(\bar{A})P(\bar{B})P(A) + \dots$

$= \frac{P(A)}{1 - P(\bar{A})P(\bar{B})} = \frac{9}{19}$

Q9. $P(A \cap B) = 0.1, P(A | B)$ and $P(B | A)$ are the roots of the equation $12x^2 - 7x + 1 = 0$

(4) $\Rightarrow P(A | B)P(B | A) = \frac{1}{12}$

$\Rightarrow \frac{P(A \cap B)}{P(B)} \times \frac{P(A \cap B)}{P(A)} = \frac{1}{12}$

$\Rightarrow P(A)P(B) = 12(0.1)^2$

$= 0.12$

$$\begin{aligned}
 \text{Also, } P(A | B) + P(B | A) &= \frac{7}{12} \\
 \Rightarrow P(A \cap B) \left(\frac{1}{P(B)} + \frac{1}{P(A)} \right) &= \frac{7}{12} \\
 \Rightarrow P(A) + P(B) &= \frac{7}{12} \times \frac{0.12}{0.1} \\
 \Rightarrow P(A) + P(B) &= 0.7 \\
 \frac{P(\bar{A} \cup \bar{B})}{P(\bar{A} \cap \bar{B})} &= \frac{P(\overline{A \cap B})}{P(\overline{A \cup B})} \\
 &= \frac{1 - P(A \cap B)}{1 - P(A \cup B)} \\
 &= \frac{1 - 0.1}{1 - (0.7 - 0.1)} = \frac{0.9}{0.4} = \frac{9}{4}
 \end{aligned}$$

Q10. Bag 1 $\rightarrow 4w, 5B$

(1) Bag 2 $\rightarrow nw, 3B$

(I) $\rightarrow$ Transferred ball is white $P(w) = \frac{n+1}{n+4} \cdot \frac{4}{9}$

(II) $\rightarrow$ Transferred ball is black $P(w) = \frac{5}{9} \cdot \frac{n}{n+4}$

$$\begin{aligned}
 \frac{4n+4}{9n+36} + \frac{5n}{9n+36} &= \frac{29}{45} \\
 \frac{9n+4}{9n+36} &= \frac{29}{45} \Rightarrow n = 6
 \end{aligned}$$

Q11. E_1 : Bag B_1 is selected

(1) B_1 B_2 B_3
 6W4B 4W6B 5W5B

E_2 : bag B_2 is selected

E_3 : Bag B_3 is selected

A : Drawn ball is white

We have to find $P\left(\frac{E_2}{A}\right)$

$$\begin{aligned}
 P\left(\frac{E_2}{A}\right) &= \frac{P(E_2)P\left(\frac{A}{E_2}\right)}{P(E_1)P\left(\frac{A}{E_1}\right) + P(E_2)P\left(\frac{A}{E_2}\right) + P(E_3)P\left(\frac{A}{E_3}\right)} \\
 &= \frac{\frac{1}{3} \times \frac{4}{10}}{\frac{1}{3} \times \frac{6}{10} + \frac{1}{3} \times \frac{4}{10} + \frac{1}{3} \times \frac{5}{10}} \\
 &= \frac{4}{15}
 \end{aligned}$$

Probability

JEE Main 2025 January

Q12.

x	0	1	-1
$P(x)$	$\frac{10}{16}$	$\frac{3}{16}$	$\frac{3}{16}$

(3)

$$\begin{aligned}\text{Var}(x) &= E(x^2) - [E(x)]^2 \\&= \sum_{i=1}^3 x_i^2 P(x_i) - (\mu)^2 \\&= 1 \times \frac{3}{16} + 1 \times \frac{3}{16} \quad [\mu = 0] \\&= \frac{6}{16} = \frac{3}{8}\end{aligned}$$

Q13. There are 3 bad oranges and 7 good oranges. $\therefore X$ = number of bad oranges drawn.

(1)

X	0	1	2
$P(X)$	$\frac{{}^7C_2}{{}^{10}C_2}$	$\frac{{}^3C_1 \times {}^7C_1}{{}^{10}C_2}$	$\frac{{}^3C_2}{{}^{10}C_2}$

$\therefore$ Variance

$$\begin{aligned}&= 0^2 \cdot \frac{{}^7C_2}{{}^{10}C_2} + 1^2 \cdot \left(\frac{3 \times 7}{{}^{10}C_2} \right) + 2^2 \cdot \left(\frac{3}{{}^{10}C_2} \right) \\&\quad - \left(0 + 1 \cdot \frac{3 \times 7}{{}^{10}C_2} + 2 \cdot \frac{3}{{}^{10}C_2} \right)^2 \\&= \frac{28}{75}\end{aligned}$$

Q1. Let R be the required relation

(2) $A = \{(1, 1)(2, 2), (3, 3)\}$

(i) $|R| = 3$, when $R = A$

(ii) $|R| = 5$, e.g. $R = A \cup \{(1, 2), (2, 1)\}$

Number of R can be [3]

(iii) $R = \{1, 2, 3\} \times \{1, 2, 3\}$

Ans. (5)

Q2. $A = \{1, 2, 3, 4\}$

(2) For relation to be reflexive

$R = \{(1, 2), (2, 3), (3, 3)\}$

Minimum elements added will be

$(1, 1), (2, 2), (4, 4)(2, 1)(3, 2)(3, 2)(3, 1)(1, 3)$

$\therefore$ Minimum number of elements = 7

Q3. $\sec^2 x - \tan^2 x = 1$ (on replacing y with x)

(2) $\Rightarrow$ Reflexive

$\sec^2 x - \tan^2 y = 1$

$\Rightarrow 1 + \tan^2 x + 1 - \sec^2 y = 1$

$\Rightarrow \sec^2 y - \tan^2 x = 1$

$\Rightarrow$ symmetric

$\sec^2 x - \tan^2 y = 1$

$\sec^2 y - \tan^2 z = 1$

Adding both

$\Rightarrow \sec^2 x - \tan^2 y + \sec^2 y - \tan^2 z = 1 + 1$

$\sec^2 x + 1 - \tan^2 z = 2$

$\sec^2 x - \tan^2 z = 1$

$\Rightarrow$ Transitive

hence equivalence relation

Q4. $A = \{1, 2, \dots, 10\}$

(2) $B = \left\{ \frac{m}{n} = m, n \in A, m < n, \gcd(m, n) = 1 \right\}$

$n(B)$

$n = 2 \quad \left\{ \frac{1}{2} \right\}$

$n = 3 \quad \left\{ \frac{1}{3}, \frac{2}{3} \right\}$

$n = 4 \quad \left\{ \frac{1}{4}, \frac{3}{4} \right\}$

$n = 5 \quad \left\{ \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5} \right\}$

Sets and Relations

JEE Main 2025 January

$$n = 6 \quad \left\{ \frac{1}{6}, \frac{5}{6} \right\}$$

$$n = 7 \quad \left\{ \frac{1}{7}, \frac{2}{7}, \frac{3}{7}, \frac{4}{7}, \frac{5}{7}, \frac{6}{7} \right\}$$


$$n = 8 \quad \left\{ \frac{1}{8}, \frac{3}{8}, \frac{5}{8}, \frac{7}{8} \right\}$$

$$n = 9 \quad \left\{ \frac{1}{9}, \frac{2}{9}, \frac{4}{9}, \frac{5}{9}, \frac{7}{9}, \frac{8}{9} \right\}$$

$$n = 10 \quad \left\{ \frac{1}{10}, \frac{3}{10}, \frac{7}{10}, \frac{9}{10} \right\}$$

$$n(B) = 31$$

- Q5.** $A = \{(x, y) \in \mathbf{R} \times \mathbf{R} : |x + y| \geq 3\}$
(4) and $B = \{(x, y) \in \mathbf{R} \times \mathbf{R} : |x| + |y| \leq 3\}$
 $C = \{(x, y) \in A \cap B : x = 0 \text{ or } y = 0\}$



$A \cap B$ will have only common points lying on the line PQ and RS

Now, $C = \{(-3, 0), (3, 0), (0, 3), (0, -3)\}$

$$\sum_{(x,y) \in C} |x + y| = 3 + 3 + 3 + 3 = 12$$

Q6. Let $\frac{y}{x} = \lambda$

(5120) $y = \lambda x$

$$= 10 \times ({}^9C_0 + {}^9C_1 + {}^9C_2 + {}^9C_3 + \dots + {}^9C_9)$$


$$= 10 \times 2$$

$$= 10 \times 512 = 5120$$

Q7. $A : \log_{2\pi} |\sin x| + \log_{2\pi} |\cos x| = 2$

(2) $\Rightarrow \log_{2\pi} (|\sin x \cdot \cos x|) = 2$

$$\Rightarrow |\sin 2x| = \frac{8}{\pi^2}$$



Number of solution 4

B : let $\sqrt{x} = t < 2$

$$\text{Then } \sqrt{x}(\sqrt{x} - 4) + 3(\sqrt{x} - 2) + 6 = 0$$

$$\Rightarrow t^2 - 4t + 3t - 6 + 6 = 0$$

$$\Rightarrow t^2 - t = 0, t = 0, t = 1$$

$$x = 0, x = 1$$

$$\text{again let } \sqrt{x} = t > 2$$

$$\text{then } t^2 - 4t - 3t + 6 + 6 = 0$$

$$\Rightarrow t^2 - 7t + 12 = 0$$

$$\Rightarrow t = 3, 4$$

$$x = 9, 16$$


Total number of solutions

$$n(A \cup B) = 4 + 4 = 8$$

Q8. $S = \{0, 1, 2, 3, \dots\}$

(4) $\log_y y = \log_c \left(\frac{2}{5}\right)$

$$\Rightarrow y = \left(\frac{2}{5}\right)^x$$



Required

$$\text{Sum} = 1 + \left(\frac{2}{5}\right)^1 + \left(\frac{2}{5}\right)^2 + \left(\frac{2}{5}\right)^3 + \dots = \frac{1}{1 - \frac{2}{5}} = \frac{5}{3}$$

Q9. R is reflexive $\Rightarrow R$ have $(1, 1), (2, 2), (3, 3)$

(3) R is transitive

$$\because (1, 2), (2, 3) \in R \quad \therefore (1, 3) \in R$$

$$\therefore R_1 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$$

Clearly R_1 is reflexive and transitive but not symmetric.

Similarly,

$$R_2 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3), (3, 2)\}$$

$$R_3 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3), (2, 1)\}$$

Therefore, 3 relations are possible

Q10. Reflexive : $(a_1, b) R (a_1, b_1) \Rightarrow b_1 = b_1$ True

(2) Symmetric : $(a_1, b_1) R (a_2, b_2) \Rightarrow b_1 = b_2$
 $(a_2, b_2) R (a_1, b_1) \Rightarrow b_2 = b_1$ } True

Sets and Relations

JEE Main 2025 January

$$\begin{aligned}\text{Transitive : } (a_1, b_1) R (a_2, b_2) &\Rightarrow b_1 = b_2 \\ &\& (a_2, b_2) R (a_3, b_3) \Rightarrow b_2 = b_3 \\ &\Rightarrow (a_1, b_1) R (a_3, b_3) \Rightarrow \text{True } \} b_1 = b_3\end{aligned}$$

Hence Relation R is an equivalence relation Statement-I is true.

For statement –II $\Rightarrow y = b$ so False

Q11. For reflexive $(x, x) \in \mathbb{R}, x \in \mathbb{Z}$

(2) $\Rightarrow x + x = 2x \rightarrow \text{even}$

For symmetric of $(x, y) \in \mathbb{R}$ then $(y, x) \in \mathbb{R}$ when $x, y \in \mathbb{Z}$

$$x + y \rightarrow \text{even}$$

$$\Rightarrow y + x \rightarrow \text{even}$$

for transitive if $(x, y) \in \mathbb{R} \Rightarrow x + y \rightarrow \text{even}$

$$(y, z) \in \mathbb{R} \Rightarrow y + z \rightarrow \text{even}$$

$$x + 2y + z \rightarrow \text{even}$$

$$\Rightarrow x + z \text{ is even}$$

$$\Rightarrow (x, z) \in \mathbb{R}$$

$\Rightarrow \mathbb{R}$ is an equivalence relation.



Q1. as $f(x)$ is a polynomial of degree two let it be

(2) $f(x) = ax^2 + bx + c \quad (a \neq 0)$

on satisfying given conditions we get

$$C = 1 \& a = \pm 1$$

hence $f(x) = 1 \pm x^2$

also range $\in (-\infty, 1]$ hence

$$f(x) = 1 - x^2$$

now $f(k) = -2k$

$$1 - k^2 = -2k \rightarrow k^2 - 2k - 1 = 0$$

let roots of this equation be $\alpha \& \beta$ then $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$
 $= 4 - 2(-1) = 6$

Q2.

(392)

ways to connect
one of
 $\{1, 2, \dots, 98\}$ to 1

99 can have
Image 1 or 0

100 can have
Image 0 or 1

392 Ans.

Q3. $f(x) = \sec^{-1}(2[x] + 1)$

(3) $\Rightarrow 2[x] + 1 \geq 1$

$$\Rightarrow 2[x] \geq 0$$

$$\Rightarrow [x] \geq 0$$

$$\Rightarrow x \geq 0$$

Domain of $f(x)$ is $(-\infty, \infty)$

$$\text{or } 2[x] + 1 \leq -1$$

$$\text{or } 2[x] \leq -2$$

$$\text{or } [x] \leq -1$$

$$\text{or } x \leq 0$$

Functions

JEE Main 2025 January

Q4. $f_1(x) = \log_5(18x - x^2 - 77)$

(3) $\therefore 18x - x^2 - 77 > 0$

$$x^2 - 18x + 77 < 0$$

$$x \in (7, 11)$$

$$\alpha = 7, \beta = 11$$

$$f_2(x) = \log_{(x-1)} \left(\frac{2x^2 + 3x - 2}{x^2 - 3x - 4} \right)$$

$$x > 1, x - 1 \neq 1, \frac{2x^2 + 3x - 2}{x^2 - 3x - 4} > 0$$

$$x > 1, x \neq 2, \frac{(2x-1)(x+2)}{(x-4)(x+1)} > 0$$

$$x > 1, x \neq 2,$$



$$\therefore x \in (4, \infty)$$

$$\therefore \gamma = 4$$

$$\therefore \alpha^2 + \beta^2 + \gamma^2 = 49 + 121 + 16 = 186$$

Q5. $f(x) = 6 + 16 \cos x \cdot \cos \left(\frac{\pi}{3} - x \right)$

(1) $\cos \left(\frac{\pi}{3} + x \right) \cdot \sin 3x \cdot \cos 6x$

$$f(x) = 6 + 4 \cos 3x \cdot \sin 3x \cdot \cos 6x$$

$$\therefore f(x) = 6 + \sin 12x$$

$$\therefore \text{Range of } f(x) = [5, 7]$$

$$\therefore [\alpha, \beta] = [5, 7]$$

$$\therefore \text{Distance of point from } 3x + 4y + 12 = 0$$

$$= \left| \frac{3.5 + 4.7 + 12}{\sqrt{3^2 + 4^2}} \right|$$

$$= 11 \text{ units}$$

Q6. $f(g(x)) = \ln \left(\frac{x^4 - 2x^3 + 3x^2 - 2x + 2}{2x^2 - 2x + 1} \right)$

(4) Since $2x^2 - 2x + 1 > 0 \quad \forall x \in \mathbb{R} \therefore (-2)^2 - 4(2) < 0$

Consider

$$x^4 - 2x^3 + 3x^2 - 2x + 2$$

$$= (x^4 - 2x^3 + x^2) + (x^2 - 2x + 1) + (1 + x^2)$$

$$= x^2(x-1)^2 + (x-1)^2 + (x^2 + 1) > 0 \quad \forall x \in \mathbb{R}$$

$$\Rightarrow g(x) > 0 \quad \forall x \in \mathbb{R}$$

$$\Rightarrow \ln f(g(x)), f(x) > 0 \quad \forall x \in \mathbb{R}$$

$$\Rightarrow x \in \mathbb{R} \text{ is domain}$$

Functions

JEE Main 2025 January

Q7. $A = \{1, 2, 3, 4\}$

(1) $B = \{1, 4, 9, 16\}$

Total number of functions $= 4^4$

Total number of one-one functions $= 4!$

Total number of many one functions $= 4^4 - 4! = 232$

Total number of many-one functions in which $1 \notin f(A) = 3 \times 3 \times 3 \times 3 = 81$

$\therefore$ Total number of many one functions $1 \notin f(A)$

$= 232 - 81$

$= 151$

Q8. as $f(x)$ is onto hence A is range of $f(x)$

(2) $f'(x) = 6x^2 - 30x + 36$

$= 6(x-2)(x-3)$

now $f(2) = 16 - 60 + 72 + 7 = 35$

$f(3) = 54 - 135 + 108 + 7 = 34$

$f(0) = 7$

hence range $\in [7, 35] = A$

also for range of $g(x)$

$g(x) = 1 - \frac{1}{x^{2025} + 1} \in [0, 1) = B$

$s = \{0, 7, 8, \dots, 35\}$ hence $n(s) = 30$

Q9. $f(x) = \frac{42^x + 16}{2 \cdot 2^{2x} + 16 \cdot 2^x + 32}$

(2) $f(x) = \frac{2(2^x + 4)}{2^{2x} + 8 \cdot 2^x + 16}$

$f(x) = \frac{2}{2^x + 4}$

$f(4-x) = \frac{2^x}{2(2^x + 4)}$

$f(x) + f(4-x) = \frac{1}{2}$

So, $f\left(\frac{1}{15}\right) + f\left(\frac{59}{15}\right) = \frac{1}{2}$

Similarly $= f\left(\frac{29}{15}\right) + f\left(\frac{31}{15}\right) = \frac{1}{2}$


$f\left(\frac{30}{15}\right) = f(2) = \frac{2}{2^2 + 4} = \frac{2}{8} = \frac{1}{4}$

$\Rightarrow 8\left(29 \times \frac{1}{2} + \frac{1}{4}\right)$

Ans. 118

Q10. $x^2 + 3x + 2 = \min\{|x - 3|, |x + 2|\}$

(3)



$$y = x^2 + 3x + 2$$

$$y = x^2 + 2\left(\frac{3}{2}\right)x + \frac{9}{4} - \frac{9}{4} + 2$$

$$y = \left(x + \frac{3}{2}\right)^2 - \frac{1}{4}$$

$$y + \frac{1}{4} = \left(x + \frac{3}{2}\right)^2$$

$$\Rightarrow \text{Parabola vertex } \left(\frac{-3}{2}, \frac{-1}{4}\right)$$

$\Rightarrow$ By graph 2 solution possible

Q11. $f(x) - 6f\left(\frac{1}{x}\right) = \frac{35}{3x} - \frac{5}{2} \dots (1)$

(3)

$$6\left(f\left(\frac{1}{x}\right) - 6f(x)\right) = \frac{35x}{3} - \frac{5}{2}$$

$$6f\left(\frac{1}{x}\right) - 36f(x) = \frac{210x}{3} - \frac{30}{2} \dots (2)$$

$$(1) + (2)$$

$$-35f(x) = \frac{35}{3}\left[\frac{1}{x} + 6x\right] - \frac{5}{2}(1 + 6)$$

$$-f(x) = \frac{1}{3}\left(\frac{1}{x} + 6x\right) - \frac{1}{2}$$

$$f(x) = -\frac{1}{3x} - 2x + \frac{1}{2}$$

$$\lim_{x \rightarrow 0} \left[\frac{1}{\alpha x} - \frac{1}{3x} - 2x + \frac{1}{2} \right] = \beta$$

$$\Rightarrow \alpha = 3$$

$$\beta + 2\beta = 3 + 2 \times \frac{1}{2} = 4$$

Q12. $f(x) = \frac{2^x}{2^x + \sqrt{2}}$
(4)

$$f(x) + f(1-x) = \frac{2^x}{2^x + \sqrt{2}} + \frac{2^{1-x}}{2^{1-x} + \sqrt{2}}$$

$$= \frac{2^x}{2^x + \sqrt{2}} + \frac{2}{2 + \sqrt{2}2^x} = \frac{2^x + \sqrt{2}}{2^x + \sqrt{2}} = 1$$

Now, $\sum_{k=1}^{81} f\left(\frac{k}{82}\right) = f\left(\frac{1}{82}\right) + f\left(\frac{2}{82}\right) + \dots + f\left(\frac{81}{82}\right)$

$$= f\left(\frac{1}{82}\right) + f\left(\frac{1}{82}\right) + \dots + f\left(1 - \frac{2}{82}\right) + f\left(1 - \frac{1}{82}\right)$$

$$\left[f\left(\frac{1}{82}\right) + f\left(1 - \frac{1}{82}\right) \right] + \left[f\left(\frac{2}{82}\right) + f\left(1 - \frac{2}{82}\right) \right] + \dots + 40 \text{ cases} + f\left(\frac{41}{82}\right)$$

$$(1 + 1 + \dots + 1)40 \text{ times} + \frac{2^{1/2}}{2^{1/2} + 2^{1/2}}$$

$$40 + \frac{1}{2} = \frac{81}{2}$$

Q13. $f(x) = \frac{2^{2x} - 1}{2^{2x} + 1}$
(4)

$$= 1 - \frac{2}{2^{2x} + 1}$$

$$f'(x) = \frac{2}{(2^{2x} + 1)^2} \cdot 2 \cdot 2^{2x} \cdot \ln 2 \text{ i.e always +ve}$$

so $f(x)$ is $\uparrow$ function

$$\therefore f(-\infty) = -1$$

$$f(\infty) = 1$$

$$\therefore f(x) \in (-1, 1) \neq \text{co-domain}$$

so function is one-one but not onto

Q14. Put $y = 0$

(4) $f(x) = f(0) + f(x) + 1 - 0$

$$f(0) = -1$$

$$f(0) = 0 + 0 + b$$

$$\Rightarrow b = -1$$

$$f(-1+1) = f(-1) + f(1) + 1 + \frac{2}{7}$$

$$f(0) = f(-1) + f(1) + \frac{9}{7}$$

$$-1 = (2 + 3a) + \left(\frac{a+2}{a-1}\right)(-1) + b + (2 + 3a)$$

$$+ \frac{a+2}{a-1} + b + \frac{9}{7}$$

Functions

JEE Main 2025 January

$$-1 = 4 + 6a - 2 + \frac{9}{7}$$

$$-1 = 2 + \frac{9}{7} + 6a$$

$$6a = -1 - 2 - \frac{9}{7}$$

$$a = \frac{-5}{7}$$

$$f(x) = \frac{-x^2}{7} + \frac{\frac{9}{7}}{\frac{-12}{7}}x - 1$$


$$f(x) = \frac{-x^2}{7} - \frac{3}{4}x - 1$$

$$\sum_{i=1}^5 f(i) = -\frac{1}{7} \left(\frac{5 \times 6 \times 11}{6} \right) - \frac{3}{4} \left(\frac{5 \times 6}{2} \right) - 5$$

$$= \frac{-55}{7} - \frac{45}{4} - 5$$

$$= \frac{675}{28}$$

$$\Rightarrow 28 \left| \sum_{i=1}^5 f(i) \right| = 675$$



Limits

JEE Main 2025 January

Q1. $\lim_{x \rightarrow 0} \operatorname{cosec} \left(\sqrt{2 \cos^2 x + 3 \cos x} - \sqrt{\cos^2 x + \sin x + 4} \right)$

(4)
$$\lim_{x \rightarrow 0} \frac{\operatorname{cosec} x (\cos^2 x + 3 \cos x - \sin x - 4)}{\left(\sqrt{2 \cos^2 x + 3 \cos x} + \sqrt{\cos^2 x + \sin x + 4} \right)}$$
$$\lim_{x \rightarrow 0} \frac{1}{\sin x} \frac{(\cos^2 x + 3 \cos x - 4) - \sin x}{\left(\sqrt{2 \cos^2 x + 3 \cos x} + \sqrt{\cos^2 x + \sin x + 4} \right)}$$
$$\lim_{x \rightarrow 0} \frac{(\cos^2 x + 4)(\cos x - 1) - \sin x}{\sin x \left(\sqrt{2 \cos^2 x + 3 \cos x} + \sqrt{\cos^2 x + \sin x + 4} \right)}$$
$$\lim_{x \rightarrow 0} \frac{-2 \sin^2 \frac{x}{2} (\cos x + 4) - 2 \sin \frac{x}{2} \cos \frac{x}{2}}{2 \sin \frac{x}{2} \cos \frac{x}{2} \left(\sqrt{2 \cos^2 x + 3 \cos x} + \sqrt{\cos^2 x + \sin x + 4} \right)}$$
$$\lim_{x \rightarrow 0} \frac{-\left(\sin^2 x (\cos x + 4) + \cos \frac{x}{2} \right)}{\cos \frac{x}{2} \left(\sqrt{2 \cos^2 x + 3 \cos x} + \sqrt{\cos^2 x + \sin x + 4} \right)}$$
$$-\frac{1}{2\sqrt{5}}$$

Q2. $\alpha = \lim_{x \rightarrow \infty} \left(\left(\frac{e}{1-e} \right) \left(\frac{1}{e} - \frac{x}{1+x} \right) \right)^x$ (1^∞ form)

(4) $\therefore \alpha = e^L$

Where $L = \lim_{x \rightarrow \infty} x \left(\left(\frac{e}{1-e} \right) \left(\frac{1}{e} - \frac{x}{1+x} \right) - 1 \right)$

$\Rightarrow L = \lim_{x \rightarrow \infty} \left(\frac{e}{1-e} \right) x \left(\frac{1}{e} - \frac{x}{1+x} - \left(\frac{1-e}{e} \right) \right)$

$\Rightarrow L = \frac{e}{1-e} \lim_{x \rightarrow \infty} x \left(1 - \frac{x}{1+x} \right)$

$\Rightarrow L = \frac{e}{1-e} \lim_{x \rightarrow \infty} \frac{x}{x+1}$

$\Rightarrow L = \frac{e}{1-e} \cdot 1$

$\Rightarrow L = \frac{e}{1-e}$

$\therefore \alpha = e^{\frac{e}{1-e}} \Rightarrow \log \alpha = \frac{e}{1-e}$

$\therefore \text{Required value} = \frac{\frac{e}{1-e}}{1 + \frac{e}{1-e}} = e$

Limits

JEE Main 2025 January

Q3. $\frac{\alpha}{5e} = \exp\left(\lim_{t \rightarrow 0} \frac{1}{t} \left(\int_0^1 (3x+5)^t dx - 1 \right)\right)$

(64)
$$= \exp\left(\lim_{t \rightarrow 0} \frac{1}{t} \left(\frac{(3x+5)^{t+1}}{3(t+1)} \Big|_0^1 - 1 \right)\right)$$

$$= \exp\left(\lim_{t \rightarrow 0} \frac{1}{t} \left(\frac{8^{t+1} - 5^{t+1}}{3(t+1)} - 1 \right)\right)$$

$$= \exp\left(\lim_{t \rightarrow 0} \frac{1}{t} \left(\frac{8^{t+1} - 5^{t+1} - 3t - 3}{3(t+1)} \right)\right)$$

$$= \exp\left(\lim_{t \rightarrow 0} \left(\frac{8^{t+1} \cdot \ln 8 - 5^{t+1} \ln 5 - 3}{3(t+1)} \right)\right)$$

$$= \exp\left(\frac{\ln 8^8 - \ln 5^5 - 3}{5}\right)$$

$$= \left(\frac{8}{5}\right)^{2/3} \frac{\alpha}{5e} = \exp\left(\frac{\ln\left(\frac{8^8}{5^5}\right)}{5} - 1\right)$$

$$\Rightarrow \left(\frac{8}{5}\right)^{2/3} \frac{\alpha}{5} = \left(\frac{8^8}{5^5}\right)^{1/3} = \left(\frac{8^6 \cdot 8^2}{5^3 \cdot 5^2}\right)^{1/3} = \frac{64}{5} \left(\frac{8}{5}\right)^{2/3}$$

$$\Rightarrow \alpha = 64$$

Q4. $\lim_{x \rightarrow 0^+} \left(x \left(\left[\frac{1}{x} \right] + \left[\frac{2}{x} \right] + \dots + \left[\frac{p}{x} \right] \right) - x^2 \left(\left[\frac{1}{x^2} \right] + \left[\frac{2^2}{x^2} \right] + \left[\frac{9^2}{x^2} \right] \right) \right) \geq 1$

(24) $(1 + 2 + \dots + p) - (1^2 + 2^2 + \dots + 9^2) \geq 1$

$$\frac{p(p+1)}{2} - \frac{9 \cdot 10 \cdot 19}{6} \geq 1$$

$$p(p+1) \geq 572$$

Least natural value of p is 24

Q5. $\lim_{x \rightarrow \infty} \frac{(2x^2 - 3x + 5)(3x - 1)^{x/2}}{(3x^2 + 5x + 4)\sqrt{(3x+2)^x}}$

(3)
$$= \lim_{x \rightarrow \infty} \frac{x^2 \left(2 - \frac{3}{x} + \frac{5}{x^2}\right) (3x)^{\frac{x}{2}} \left(1 - \frac{1}{3x}\right)^{\frac{x}{2}}}{x^2 \left(3 + \frac{5}{x} + \frac{4}{x^2}\right) (3x)^{\frac{x}{2}} \left(1 + \frac{2}{3x}\right)^{\frac{x}{2}}}$$

$$\lim_{x \rightarrow \infty} \left(1 - \frac{1}{3x}\right)^{\frac{x}{2}} = e^{\lim_{x \rightarrow \infty} \left(1 - \frac{1}{3x} - 1\right) \times \frac{x}{2}} = e^{-\frac{1}{6}}$$

$$\lim_{x \rightarrow \infty} \left(1 + \frac{2}{3x}\right)^{\frac{x}{2}} = e^{\lim_{x \rightarrow \infty} \left(1 + \frac{2}{3x} - 1\right) \times \frac{x}{2}} = e^{\frac{1}{3}}$$

$$= e^{\frac{1}{3}}$$

So, $\frac{2}{3} \times \frac{e^{-\frac{1}{6}}}{e^{\frac{1}{3}}} = \frac{2}{3} \times \frac{1}{e^{\frac{1}{3} + \frac{1}{6}}} = \frac{2}{3\sqrt{e}}$

Limits

JEE Main 2025 January

Q6. (1) $f(x) = \lim_{x \rightarrow \infty} \left(\frac{\tan(x/2^{r+1}) + \tan^3(x/2^{r+1})}{1 - \tan^2(x/2^{r+1})} \right)$

$$= \lim_{x \rightarrow \infty} \frac{\tan\left(\frac{x}{2^{r+1}}\right)}{\cos\left(\frac{x}{2^r}\right)}$$

$$= \lim_{x \rightarrow \infty} \frac{\sin(x/2^{r+1})}{\cos\left(\frac{x}{2^{r+1}}\right) \cos\left(\frac{x}{2^r}\right)}$$

$$= \lim_{x \rightarrow \infty} \frac{\sin\left(\frac{x}{2^r} - \frac{x}{2^{r+1}}\right)}{\cos\left(\frac{x}{2^{r+1}}\right) \cos\left(\frac{x}{2^r}\right)}$$

$$= \lim_{x \rightarrow \infty} \tan\left(\frac{x}{2^r}\right) - \tan\left(\frac{x}{2^{r+1}}\right)$$

From condition given question

$$\therefore \lim_{n \rightarrow \infty} \sum_{r=0}^n \left[\tan\left(\frac{x}{2^r}\right) - \tan\left(\frac{x}{2^{r+1}}\right) \right] = \tan x$$


$$\therefore \lim_{x \rightarrow 0} \left(\frac{e^x - e^{\tan x}}{x - \tan x} \right)$$

$$\Rightarrow \lim_{x \rightarrow 0} e^{\tan x} \left(\frac{e^{x - \tan x} - 1}{x - \tan x} \right)$$

$$\Rightarrow \lim_{x \rightarrow 0} e^{\tan x} \lim_{x \rightarrow 0} \left(\frac{e^{x - \tan x} - 1}{x - \tan x} \right)$$

$$\Rightarrow 1.1 \left(\because \lim_{x \rightarrow 0} \frac{e^{x-1}}{x} = 1 \right)$$

$$= 1$$



Q1. $\lim_{x \rightarrow 0^-} \frac{2}{x} \{\sin(k_1 + 1)x + \sin(k_2 - 1)x\} = 4$

(4) $\Rightarrow 2(k_1 + 1) + 2(k_2 - 1) = 4$
 $\Rightarrow k_1 + k_2 = 2$
 $\Rightarrow \lim_{x \rightarrow 0^+} \frac{2}{x} \ln\left(\frac{2 + k_1 x}{2 + k_2 x}\right) = 4$
 $\Rightarrow \lim_{x \rightarrow 0^+} \frac{1}{x} \ln\left(1 + \frac{(k_1 - k_2)x}{2 + k_2 x}\right) = 2$
 $\Rightarrow \frac{k_1 - k_2}{2} = 2$
 $\Rightarrow k_1 - k_2 = 4$
 $\therefore k_1 = 3, k_2 = -1$
 $k_1^2 + k_2^2 = 9 + 1 = 10$

Q2. $\therefore f(x + y) = f(x) \cdot f(y) + f(x) \cdot f(y), \forall x, y \in R \dots (i)$

(1) And $f(0) = 1 \dots (ii)$

Now replace x by zero and y by zero we get

$$f(0) = f(0)f(0) + f(0)f(0)$$

$$1 = f(0) + f(0)$$

$$\therefore f'(0) = \frac{1}{2} \dots (iii)$$

Now replace y by zero in equation (i), we get

$$f(x) = \frac{1}{2}f(x) + f'(x)$$

$$\text{or, } \frac{1}{2}f(x) = f'(x)$$

$$\text{then } \frac{f'(x)}{f(x)} = \frac{1}{2}$$

$$\text{hence } \ln|f(x)| = \frac{x}{2} + c$$

$$\text{Put } x = 0, \text{ we get } c = 0$$

$$\therefore \ln|f(x)| = \frac{x}{2}$$

$$\text{Then } \sum_{n=1}^{100} \ln(f(\eta)) = \left(\frac{1}{2} + \frac{2}{2} + \frac{3}{2} + \dots + \frac{100}{2}\right)$$


$$= \frac{5050}{2} = 2525$$

Q3.

(2)

$$f(x) = [x] + |x - 2|, -2 < x < 3$$

$$\therefore f(x) = \begin{cases} -x, & -2 < x < -1 \\ 1 - x, & -1 \leq x < 0 \\ 2 - x, & 0 \leq x < 1 \\ 3 - x, & 1 \leq x < 2 \\ x, & 2 \leq x < 3 \end{cases}$$




Continuity and Differentiability

JEE Main 2025 January

It is clearly discontinuous at 4 points and nondifferentiable at 4 points.

$$\therefore m + n = 8$$

Q4. (5) $f(x) = \begin{cases} 3x & ; x < 0 \\ \min\{1+x, x\} & ; 0 \leq x < 1 \\ \min\{2+x, x+2\} & ; 1 \leq x < 2 \\ 5 & ; x > 2 \end{cases}$



$$f(x) = \begin{cases} 3x & ; x < 0 \\ x & ; 0 \leq x < 1 \\ x+2 & ; 1 \leq x < 2 \\ 5 & ; x > 2 \end{cases}$$

Not continuous at $x \in \{1, 2\} \Rightarrow \alpha = 2$

Not diff. at $x \in \{0, 1, 2\} \Rightarrow \beta = 3 \quad \alpha + \beta = 5$

Q5. $f(x) = (x^2 + 1)|x^2 - ax + 2| + \cos|x|$

(3) Notice that $\cos(-x) = \cos x = \cos|x|$ which means $\cos|x|$ is differentiable everywhere in $x \in \mathbb{R}$

$$\Rightarrow f(x) \text{ can be non differentiable where } |x^2 - ax + 2| = 0$$

$$\begin{aligned} &\swarrow x = \alpha = 2 \\ &\searrow x = \beta \end{aligned}$$

$$\Rightarrow x^2 - ax + 2 = 0$$

$$\Rightarrow 4 - 2a + 2 = 0 \Rightarrow a = 3$$

$$\Rightarrow (x^2 - 3x + 2) = 0 \Rightarrow x = 1, 2$$

$$\beta = 1$$

distance of (α, β) from line

$$12x + 5y + 10 = 0$$

$$\Rightarrow \frac{|2(12) + 5(1) + 10|}{13} = \frac{39}{13} = 3$$

Continuity and Differentiability

JEE Main 2025 January

Q6. $f(x)$ is continuous and differentiable


(34)

at $x = 1$, $\text{LHL} = \text{RHL}$, $\text{LHD} = \text{RHD}$

$$-3a - 2 = a^2 + b, -6a = b$$

$$a = 2; b = -12$$

$$f(x) = \begin{cases} -6x^2 - 2, & x < 1 \\ 4 - 12x, & x \geq 1 \end{cases}$$



$$\text{Area} = \int_{-\sqrt{3}}^1 (-6x^2 - 2 + 20) dx + \int_1^2 (4 - 12x + 20) dx$$


$$= 16 + 12\sqrt{3} + 6 = 22 + 12\sqrt{3}$$

$$\therefore \alpha + \beta = 34$$



Q1.

(2)



$$v = \frac{4}{3}\pi r^3$$

$$\frac{dv}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$81 = 4\pi r^2 \times \frac{1}{4\pi}$$

$$r^2 = 81$$

$$r = 9$$

$$\text{surface area of chocolate} = 4\pi(r-1)^2 = 256\pi$$

Q2. $f'(x) = \frac{2}{x-2} - 2x + a \geq 0$

(2)

$$f''(x) = \frac{-2}{(x-2)^2} - 2 < 0$$

$$f'(x) \downarrow$$

$$f'(3) \geq 0$$

$$2 - 6 + a \geq 0$$

$$a \geq 4$$

$$a_{\min} = 4$$

$$g(x) = (x-1)^3(x+2-a)^2$$

$$g(x) = (x-1)^3(x-2)^2$$

$$g'(x) = (x-1)^3 2(x-2) + (x-2)^2 3(x-1)^2$$

$$= (x-1)^2(x-2)(2x-2+3x-6)$$

$$= (x-1)^2(x-2)(5x-8) < 0$$

$$x \in \left(\frac{8}{5}, 2\right)$$

$$100(a+b-c) = 100\left(4 + \frac{8}{5} - 2\right) = 360$$


Application of Derivatives

JEE Main 2025 January

Q3. $5x^3 - 15x - a = 0$

(30) $f(x) = 5x^3 - 15x$

$$f(x) = 15x^2 - 15 = 15(x-1)(x+1)$$




$$a \in (-10, 10)$$

$$\alpha = -10, \beta = 10$$

$$\beta - 2\alpha = 10 + 20 = 30$$

Q4.

(1)



∴ Local minimum values at A & B

$$\frac{7}{3} - \frac{11}{72}$$

$$\Rightarrow \frac{168 - 11}{72} \Rightarrow \frac{157}{72}$$

Q1. $I(x) = \int \frac{dx}{(x-11)^{\frac{11}{13}}(x+15)^{\frac{15}{13}}}$

(2) Put $\frac{x-11}{x+15} = t \Rightarrow \frac{26}{(x+15)^2} dx = dt$

$$I(x) = \frac{1}{26} \int \frac{dt}{t^{11/13}} = \frac{1}{26} \cdot \frac{t^{2/13}}{2/13}$$

$$I(x) = \frac{1}{4} \left(\frac{x-11}{x+15} \right)^{2/13} + C$$

$$I(37) - I(24) = \frac{1}{4} \left(\frac{26}{52} \right)^{2/13} - \frac{1}{4} \left(\frac{13}{39} \right)^{2/13}$$

$$= \frac{1}{4} \left(\frac{1}{2^{2/13}} - \frac{1}{3^{2/13}} \right)$$

$$= \frac{1}{4} \left(\frac{1}{4^{1/13}} - \frac{1}{9^{1/13}} \right)$$

$$\therefore b = 4, c = 9$$

$$3(b+c) = 39$$

Q2. $2x^2 + 5x + 9 = A(x^2 + x + 1) + B(2x + 1) + C$

(16) $A = 2 \quad B = \frac{3}{2} \quad C = \frac{11}{2}$

$$2 \int \sqrt{x^2 + x + 1} dx + \frac{3}{2} \int \frac{2x+1}{\sqrt{x^2 + x + 1}} dx + \frac{11}{2} \int \frac{dx}{\sqrt{x^2 + x + 1}}$$

$$2 \int \sqrt{\left(x + \frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} dx + 3\sqrt{x^2 + x + 1} + \frac{11}{2} \int \frac{dx}{\sqrt{\left(x + \frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2}}$$

$$2 \left(\frac{x + \frac{1}{2}}{2} \sqrt{x^2 + x + 1} + \frac{3}{8} \ln \left(x + \frac{1}{2} + \sqrt{x^2 + x + 1} \right) \right) + 3\sqrt{x^2 + x + 1}$$

$$+ \frac{11}{2} \ln \left(x + \frac{1}{2} + \sqrt{x^2 + x + 1} \right) + C$$

$$\alpha = \frac{7}{2} \quad \beta = \frac{25}{4}$$

$$\alpha + 2\beta = 16$$

Indefinite Integration

JEE Main 2025 January

Q3. (2) $\frac{d}{dx} \left(\frac{x \cdot \sin^{-1} x}{\sqrt{1-x^2}} \right) - \sin^{-1} x \cdot \left(\frac{1 \cdot \sqrt{1-x^2} + \frac{x \cdot 2x}{2\sqrt{1-x^2}}}{1-x^2} \right)$

$$= \frac{x}{\sqrt{1-x^2}} \cdot \frac{1}{\sqrt{1-x^2}}$$

$$= \frac{\sin^{-1} x}{(1-x^2)^{3/2}} + \frac{x}{1-x^2}$$

$$\text{Hence, } I = \int e^x (f(x) + f'(x)) dx$$

$$= e^x \cdot f(x) + C$$

$$I = e^x \cdot \frac{x \cdot \sin^{-1} x}{\sqrt{1-x^2}} + C = g(x) + C$$

$$\Rightarrow g(x) = \frac{x e^x \sin^{-1} x}{\sqrt{1-x^2}} \text{ and } g(1/2) = \frac{\pi}{6} \sqrt{\frac{e}{3}}$$

Q4. (2) $\int x^3 \sin x dx = -x^3 \cos x + \int 3x^2 \cos x dx$

$$= -x^3 \cos x + 3x^2 \sin x - \int 6x \sin x dx$$

$$= -x^3 \cos x + 3x^2 \sin x + 6x \cos x - 6 \sin x + c$$

$$\text{So } g(x) = -x^3 \cos x + 3x^2 \sin x + 6x \cos x - 6 \sin x$$

$$g\left(\frac{\pi}{2}\right) = \frac{3\pi^2}{4} - 6$$

$$g'(x) = -3x^2 \cos x + x^3 \sin x + 6 \cos x - 6 \sin x$$

$$g'\left(\frac{\pi}{2}\right) = \frac{\pi^3}{8}$$

$$8 \left(g\left(\frac{\pi}{2}\right) + g'\left(\frac{\pi}{2}\right) \right) = \pi^3 + 6\pi^2 - 48$$

$$\text{So } \alpha + \beta - \gamma = 55$$

Q5. Put $x^{1/4} = t \Rightarrow dx = 4t^3 dt$

(1) $\int \frac{4t^3 dt}{t(t+1)} = 4 \int \left(\frac{t^2-1}{t+1} + \frac{1}{t+1} \right) dt$

$$f(x) = 4 \left[\frac{x^{1/2}}{2} - x^{1/4} + \ln|x^{1/4} + 1| \right] + C$$

$$f(0) = -6$$

$$\Rightarrow C = -6$$

$$\Rightarrow f(1) = 4(\log_e 2 - 2)$$

Q1. $I = \int_0^{\frac{\pi}{4}} \left(\frac{\sin \theta + \cos \theta}{9 - 16 \sin 2\theta} \right) d\theta$

(2)

Take $\sin \theta - \cos \theta = t$

$(\cos \theta + \sin \theta) d\theta = dt$

$(\sin \theta - \cos \theta)^2 = t^2$

$\Rightarrow \sin 2\theta = 1 - t^2$

$\theta = 0 \rightarrow t = -1$

$\theta = \frac{\pi}{4} \rightarrow t = 0$

$I = \int_{-1}^0 \frac{dt}{9 + 16(1 - t^2)}$

$= \frac{1}{16} \int_{-1}^0 \frac{dt}{\frac{25}{16} - t^2}$

$= \frac{1}{4} \left[\frac{1}{10} \log \left| \frac{5-4t}{5+4t} \right| \right]_{-1}^0$

$= \frac{1}{40} [0 + \log_e 9]$

$I = \frac{\log_e 9}{40}$

$80I = 2 \log_e 9$

$80I = 4 \log_e 3$

Q2. $\int_0^2 xF'(x) dx = 6$

(1)

$= xF(x) \Big|_0^2 - \int_0^2 f(x) dx = 6$

$= 2F(2) - \int_0^2 xF(x) dx = 6 \left[\because f(2) = 2F(2) = 2 \right]$

$\int_0^2 xF(x) dx = -2 \dots (1)$

$\Rightarrow \int_0^2 F(x) dx = -2 \dots (2)$

Also

$\int_0^2 x^2 F''(x) dx = x^2 F'(x) \Big|_0^2 - 2 \int_0^2 x' F'(x) dx = 40$

$= 4F'(2) - 2 \times 6 = 40$

$F'(2) = 13$

$\therefore F'(2) + \int_0^2 F(x) dx = 13 - 2 = 11$

Q3. Let $I = 24 \int_0^{\frac{\pi}{2}} \left(\sin \left| 4x - \frac{\pi}{2} \right| + [2 \sin x] \right) dx \dots (i)$

(12)

Now $\left| 4x - \frac{\pi}{2} \right| = \begin{cases} -4x + \frac{\pi}{12} & ; x < \frac{\pi}{48} \\ 4x - \frac{\pi}{12} & ; x \geq \frac{\pi}{48} \end{cases}$

∴ from (i)

$$I = 24 \int_0^{\frac{\pi}{48}} -\sin\left(4x - \frac{\pi}{12}\right) dx + \int_{\frac{\pi}{48}}^{\frac{\pi}{4}} \sin\left(4x - \frac{\pi}{12}\right)$$

$$+ \int_0^{\frac{\pi}{6}} [2 \sin x] dx + \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} [2 \sin x] dx$$

$$I = 24 \left[\frac{\left(1 - \cos \frac{\pi}{12}\right)}{4} - \frac{\left(-\cos \frac{\pi}{12} - 1\right)}{4} \right] + \frac{\pi}{4} - \frac{\pi}{6}$$

$$I = 24 \left(\frac{1}{2} \right) + \frac{\pi}{4} - \frac{\pi}{6}$$

$$I = 2\pi + 12 = 2\pi + \alpha \text{ (from above)}$$

$$\therefore \alpha = 12$$


Q4.
(1) $f(x) = \int_0^{x^2} \frac{t^2 - 8t + 15}{e^t} dt, x \in R$

$$f'(x) = \frac{x^4 - 8x^2 + 15}{e^{x^2}} (2x) = 0$$

$$\Rightarrow \frac{2 \times (x^2 - 5)(x^2 - 3)}{e^{x^2}} = 0$$

$$\Rightarrow x(x + \sqrt{5})(x - \sqrt{5})(x + \sqrt{3})(x - \sqrt{3}) = 0$$

By using wavy curve method



Number of local maximum = 2

Number of local minimum = 3

Q5.
(1) $\int_0^x t f(t) dt = x^2 f(x)$

$$\Rightarrow x f(x) = 2x f(x) + x^2 f'(x)$$

$$\Rightarrow x^2 \cdot f'(x) = -x f(x)$$

$$\Rightarrow \frac{f'(x)}{f(x)} = -\frac{1}{x}$$

$$\int \frac{f'(x)}{f(x)} dx = - \int \frac{1}{x} dx$$

$$\Rightarrow \ln f(x) = -\ln x + C$$

$$f(2) = 3 \Rightarrow \ln 3 = -\ln 2 + C$$

$$C = \ln 6$$

$$\Rightarrow f(x) = \frac{6}{x} \Rightarrow f(6) = 1$$

Definite Integration

JEE Main 2025 January

Q6. $g(x) = x^2 + x^{7/3}$

(4) $g'(x) = 2x + \frac{7}{3}x^{4/3}$

$$f(x) = \frac{g'(x)}{x}$$

$$f(x) = 2 + \frac{7}{3}x^{1/3}$$


$$f(r^3) = 2 + \frac{7}{3}r$$

$$\sum_{r=1}^{15} \left(2 + \frac{7}{3}r \right) = 2(15) + \frac{7}{3} \left(\frac{15(16)}{2} \right)$$

$$= 310$$

Q7. $f'(x) = x(x^2 - 9x + 20), x \in (1, 5)$

(4) $= (x-4)x(x-5)$



$$\Rightarrow f'(x) > 0 \forall x \in (1, 4)$$

$$\Rightarrow f'(x) < 0 \forall x \in (4, 5)$$

$$\Rightarrow f(x) \text{ increasing in } (1, 4)$$

$$f(x) \text{ decreasing in } (4, 5)$$

$$\Rightarrow \text{critical points to check:}$$

$$x = 1, 4, 5$$

$$f(x) = \int_0^x (t^3 - 9t^2 + 20t) dt$$

$$= \frac{t^4}{4} - 3t^3 + 10t^2 \Big|_0^x = \frac{x^4}{4} - 3x^3 + 10x^2$$

$$f(1) = \frac{1}{4} - 3 + 10 = \frac{29}{4}$$

$$f(4) = 4^3 - 3 \cdot 4^3 + 10 \cdot 4^2 = -2 \cdot 4^3 + 10 \cdot 4^2 = 32$$

$$f(5) = \frac{5^4}{4} - 3 \cdot 5^3 + 10 \cdot 25 = \frac{5^4}{4} - 125 = \frac{125}{4}$$

$$\text{Range} \Rightarrow \left[\frac{29}{4}, 32 \right] \Rightarrow 4(\alpha + \beta) = 128 + 29$$

$$= 157$$

Definite Integration

JEE Main 2025 January

Q8. $I(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx$

(4) Let $x = \sin^2 \theta$ $dx = 2 \sin \theta \cos \theta d\theta$

$$I(m, n) = 2 \int_0^{\pi/2} (\sin \theta)^{2m-1} (\cos \theta)^{2n-1} d\theta$$

$$I(9, 14) + I(10, 13) = 2 \int_0^{\pi/2} (\sin \theta)^{17} (\cos \theta)^{27} d\theta$$

$$+ 2 \int_0^{\pi/2} (\sin \theta)^{19} (\cos \theta)^{25} d\theta$$

$$= 2 \int_0^{\pi/2} (\sin \theta)^{17} (\cos \theta)^{25} d\theta$$

$$= I(9, 13)$$

Q9. $f(x) = 7 \tan^8 x + 7 \tan^6 x - 3 \tan^4 x - 3 \tan^2 x$

(2) $= 7 \tan^6 x (1 + \tan^2 x) - 3 \tan^2 x (1 + \tan^2 x)$

$$= (7 \tan^6 x - 3 \tan^2 x) (1 + \tan^2 x)$$

$$= (7 \tan^6 x - 3 \tan^2 x) \sec^2 x$$

$$I_1 = \int_0^{\pi/4} f(x) dx = \int_0^{\pi/4} (7 \tan^6 x - 3 \tan^2 x) \sec^2 x dx$$

$$= \left(\frac{7 \tan^7 x}{7} - \frac{3 \tan^3 x}{3} \right) \Big|_0^{\pi/4} = 1 - 1 = 0$$

$$I_2 = \int_0^{\pi/4} x f(x) dx = \int_0^{\pi/4} x (7 \tan^6 x - 3 \tan^2 x) \sec^2 x dx$$

$$= x (\tan^7 x - \tan^3 x) \Big|_0^{\pi/4} - \int_0^{\pi/4} 1 \cdot (\tan^7 x - \tan^3 x) dx$$

$$= 0 - \int_0^{\pi/4} \tan^3 x (\tan^2 x - 1) (\tan^2 x + 1) dx$$

$$= \int_0^{\pi/4} (\tan^3 x - \tan^5 x) \sec^2 x dx = \frac{\tan^4 x}{4} - \frac{\tan^6 x}{6} \Big|_0^{\pi/4}$$

$$= \frac{1}{12}$$

$$\text{Hence } 7I_1 + 12I_2 = 1$$

Definite Integration

JEE Main 2025 January

Q10.

$$(3) \quad \text{Put } \ln x = t \Rightarrow \frac{1}{x} dx = dt \quad \begin{array}{c|c} x & t \\ \hline e^2 & 2 \\ e^4 & 4 \end{array}$$

$$I = \int_2^4 \frac{e^{(t^2+1)^{-1}}}{e^{(t^2+1)^{-1}} + e^{((6-t)^2+1)^{-1}}} dt \quad \dots (i)$$

$$I = \int_2^4 \frac{e^{((6-t)^2+1)^{-1}}}{e^{((6-t)^2+1)^{-1}} + e^{(t^2+1)^{-1}}} dt \quad \dots (ii)$$

$$\left\{ \text{Using } \int_a^0 f(x) dx = \int_a^0 f(a+b-x) dx \right\}$$

Adding (i) and (ii) gives

$$2I = \int dt \Rightarrow I = 1$$

Q11.

$$(3) \quad I = \int_0^{\frac{\pi}{2}} \frac{(\sin x)^{\frac{3}{2}} dx}{(\sin x)^{\frac{3}{2}} x + (\cos x)^{\frac{3}{2}}} = \int_0^{\frac{\pi}{2}} \frac{\sin^{\frac{3}{2}}\left(\frac{\pi}{2} - x\right) dx}{\sin^{\frac{3}{2}}\left(\frac{\pi}{2} - x\right) + \cos^{\frac{3}{2}}\left(\frac{\pi}{2} - x\right)}$$

$$\Rightarrow \text{Adding } 2I = \int_0^{\frac{\pi}{2}} \frac{(\sin x)^{\frac{3}{2}} + (\cos x)^{\frac{3}{2}}}{(\sin x)^{\frac{3}{2}} + (\cos x)^{\frac{3}{2}}} dx = \frac{\pi}{2}$$

$$I_0 = \int_0^{\frac{\pi}{2}} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} dx = \int_0^{\frac{\pi}{2}} \frac{\left(\frac{\pi}{2} - x\right) \sin x \cos x}{(\sin x)^4 + (\cos x)^4} dx$$

$$\text{Adding, } 2I_0 = \int_0^{\frac{\pi}{2}} \frac{\frac{\pi}{2} (\sin x) \cos x}{(\sin^4 x + \cos^4 x)} dx$$

$$\Rightarrow I_0 = \frac{\pi}{4} \int_0^{\frac{\pi}{2}} \frac{\tan x (\sec^2 x) dx}{1 + \tan^4 x}$$

$$\text{put } \tan^2 x = t \Rightarrow 2 \tan x \sec^2 x dx = dt$$

$$\Rightarrow I_0 = \frac{\pi}{4} \int_0^{\infty} \frac{\frac{dt}{2}}{(1+t^2)} = \frac{\pi}{8} (\tan^{-1} t) \Big|_0^{\infty} = \frac{\pi}{8} \left(\frac{\pi}{2} - 0 \right)$$

$$\Rightarrow I_0 = \frac{\pi^2}{16}$$

Q12.

$$(4) \quad I = \int_0^{\frac{\pi}{2}} \frac{96x^2 \cos^2 x}{1 + e^x} dx$$

$$2I = 2 \int_0^{\frac{\pi}{2}} 96x^2 \cos^2 x dx$$

$$I = 96 \int_0^{\frac{\pi}{2}} x^2 \cos^2 x dx$$

$$= 48 \int_0^{\frac{\pi}{2}} x^2 (1 + \cos 2x) dx$$

$$= 2\pi^2 + 48(0 - 0) - 48 \int_0^{\frac{\pi}{2}} x \sin 2x dx$$

$$= 2\pi^2 - 12\pi + [0 - 0] = \pi (2\pi^2 - 12)$$

$$= \pi (\alpha\pi^2 + \beta)$$

$$\Rightarrow (\alpha + \beta)^2 = 100$$

Definite Integration

JEE Main 2025 January

Q13. Given, $\int_0^1 f(\lambda x) d\lambda = af(x)$

(112) Let $\lambda x = u$

$$d\lambda = \frac{1}{x} du$$

$$\therefore \text{From (1)} \quad \frac{1}{x} \int_0^x f(u) du = af(x)$$

$$\Rightarrow \int_0^x f(u) du = axf(x)$$

Differentiate both sides

$$f(x) = a(xf'(x) + f(x))$$

$$\Rightarrow f(x) = axf'(x) + af(x)$$

$$\Rightarrow (1-a)f(x) = axf'(x)$$

$$\Rightarrow \frac{f'(x)}{f(x)} = \frac{(1-a)}{a} \cdot \frac{1}{x}$$

Integrate both side w.r.t. (x)

$$\Rightarrow \int \frac{f'(x)}{f(x)} dx = \frac{(1-a)}{a} \int \frac{1}{x} dx$$

$$\Rightarrow \ln f(x) = \left(\frac{1-a}{a} \right) \ln x + c$$

Now at $x = 1$, $f(1) = 1$

$$\Rightarrow c = 0$$

Also given $f(16) = \frac{1}{8}$

$$\therefore \frac{1}{8} = (16)^{\frac{1-a}{a}}$$

$$\Rightarrow 2^{-3} = 2^{\frac{4-4a}{a}}$$

$$\Rightarrow -3 = \frac{4-4a}{a}$$

$$\Rightarrow -3a = 4 - 4a$$

$$\Rightarrow a = 4$$

$$\therefore f(x) = x^{-3/4}$$

$$f(x) = \frac{-3}{4} x^{-7/4}$$


$$\text{Put } x = \frac{1}{16}$$

$$f' \left(\frac{1}{16} \right) = \frac{-3}{4} \left(\frac{1}{16} \right)^{-7/4} = \frac{-3}{4} \cdot 2^{-4x \left(\frac{-7}{4} \right)} = -96$$

$$\therefore 16 - f' \left(\frac{1}{16} \right) \Rightarrow 16 - (-96) = 112$$

Q1.

(2)



$$\text{Required area} = 2 \int_0^{2\sqrt{3}} \left(\sqrt{4\sqrt{3}x - x^2} - \sqrt{2\sqrt{3}x} \right) dx$$

$$= 2 \int_0^{2\sqrt{3}} \left(\sqrt{12 - (x - 2\sqrt{3})^2} - \sqrt{2\sqrt{3}x} \right) dx$$

$$= 2 \left[\frac{x - 2\sqrt{3}}{2} \sqrt{12 - (x - 2\sqrt{3})^2} + \frac{12}{2} \sin^{-1} \left(\frac{x - 2\sqrt{3}}{2\sqrt{3}} \right) \right.$$


$$\left. - \frac{\sqrt{2\sqrt{3}x^{\frac{3}{2}}}}{\frac{3}{2}} \right]_0^{2\sqrt{3}}$$

$$= 2\{3\pi - 8\}$$

$$= 6\pi - 16 \text{ sq. units.}$$

Q2.

(1)



$$\text{Area} = \int_0^2 (\sqrt{16 - 8x} - (x^2 - 4x + 4)) dx$$

$$= \left[\frac{-(16 - 8x)^{3/2}}{12} - \frac{x^3}{3} + 2x^2 + 4x \right]_0^2$$


$$= \frac{8}{3}$$

Area Under Curves

JEE Main 2025 January

Q3.

(2)



$$\begin{aligned}\text{Area} &= 2 \left[\int_0^3 (x^2 + 1) dx + \frac{1}{2} [5 + 7] \times 1 \right] \\ &= \frac{64}{3}\end{aligned}$$

Q4. $x(1 + y^2) = 1$

(2) $y^2 = 2x$


From equation (1) & (2)

$$x(1 + 2x) = 1 \Rightarrow 2x^2 + x - 1 = 0$$

$$\Rightarrow x = \frac{1}{2}, x = -1 \text{ (Reject)}$$


$$\Rightarrow y^2 = 2 \left(\frac{1}{2} \right)$$

$$\Rightarrow y = \pm 1$$



$$\begin{aligned} \text{Area bounded} &= \int_{-1}^1 \left(\frac{1}{1+y^2} - \frac{y^2}{2} \right) dy \\ &= \left(\tan^{-1} y - \frac{y^3}{6} \right) \Big|_{-1}^1 \\ &= \frac{\pi}{2} - \frac{1}{3} \end{aligned}$$

Q5.
(1)



$$e^x = 1 - e^x \Rightarrow 2e^x = 1$$

$$\Rightarrow e^x = \frac{1}{2}$$

$$\Rightarrow x = \ln \frac{1}{2}$$

$$\int_{\ln(1/2)}^0 [e^x - (1 - e^x)] dx$$


$$= \int_{\ln 2}^0 (2e^x - 1) dx = 2e^x - x \Big|_{-\ln 2}^0$$

$$= 2 - (1 + \ln 2)$$

$$= 1 - \log_e 2$$

Q6.

(77)



$$x^2 + y^2 = 5$$

$$x^2 + (x - 1)^2 = 25 \Rightarrow x = 4$$

$$x^2 + (-x + 1)^2 = 5 \Rightarrow x = -3$$

$$A = 25\pi - \int_{-3}^4 \sqrt{25 - x^2} dx + \frac{1}{2} \times 4 \times 4 + \frac{1}{2} \times 3 \times 3$$

$$A = 25\pi + \frac{25}{2} - \left[\frac{x}{2} \sqrt{25 - x^2} + \frac{25}{2} \sin^{-1} \frac{x}{5} \right]_{-3}^4$$

$$A = 25\pi + \frac{25}{2} - \left[6 + \frac{25}{2} \sin^{-1} \frac{4}{5} + 6 + \frac{25}{2} \sin^{-1} \frac{3}{5} \right]$$

$$A = 25\pi + \frac{1}{2} - \frac{25}{2} \cdot \frac{\pi}{2}$$

$$A = \frac{75\pi}{4} + \frac{1}{2}$$



$$A = \frac{1}{4}(75\pi + 2)$$

$$b = 75, c = 2$$

$$b + c = 75 + 2 = 77$$

Q7. $x^2 + 4x + 2 \leq y \leq |x + 2|$


(4)



$$\begin{aligned}
 A_1 &= \int_{-4}^0 [2 - (x^2 + 4x + 2)] dx - \frac{1}{2} \times 4 \times 2 \\
 &= \left(\frac{-x^3}{3} - 2x^2 \right)_{-4}^0 - 4 \\
 &= 0 - \left(\frac{64}{3} - 32 \right) - 4 \\
 &= 32 - \frac{64}{3} - 4 = \frac{20}{3}
 \end{aligned}$$

Q8.

(2)



$$\text{Required area} = \pi - 4 \int_0^1 (1 - x^2) dx$$

$$= \pi - 4 \left[x - \frac{x^3}{3} \right]_0^1$$

$$= \pi - 4 \times \frac{2}{3} = \pi - \frac{8}{3}$$

$$\therefore \alpha = \pi - \frac{8}{3}$$

$$9\alpha = 9\pi - 24 \rightarrow \beta = 9, \gamma = -24$$

$$|\beta - \gamma| = |9 + 24| = 33$$

Q9.


(3)

$$y \in [0, a + e^{|x|} - e^{-x}]$$

$$(i) \text{ If } x \geq 0 \Rightarrow y \in \left(0, a + e^x - \frac{1}{e^x} \right)$$


$$\text{if } x < 0 \Rightarrow y \in (0, a + e^{-x} - e^{-x})$$

$$\Rightarrow y \in (0, a)$$



$$\begin{aligned} \text{Area} &= (a) + \int_0^1 (a + e^x - e^{-x}) dx = \frac{e^2 + 8e + 1}{e} \\ &= a + (ax + e^x + e^{-x}) \Big|_0^1 = e + 8 + \frac{1}{e} \\ &= a + \left(a + e + \frac{1}{e} - 2 \right) = e + \frac{1}{e} + 8 \\ &\Rightarrow 2a - 2 = 8 \Rightarrow a = 5 \end{aligned}$$

Q10.



$A \Rightarrow$ Rectangle ABDE - Area of region EDC

$$A \Rightarrow 4 - 2 \int_0^1 (3 - x) - \left(\frac{x^2 + 3}{2} \right) dx$$


$$A \Rightarrow 4 - 2 \left\{ 3x - \frac{x^2}{2} - \frac{x^3}{6} - \frac{3}{2}x \right\}_0^1$$

$$A \Rightarrow 4 - 2 \left\{ 3 - \frac{1}{2} - \frac{1}{6} - \frac{3}{2} \right\} = \frac{7}{3}$$

So $6A = 14$

Q11. $t, \left(9 - \frac{11t^2}{3} - t\right)$

(4)



$$A = 9t - t^2 - \frac{11}{3}t^3$$

$$\frac{dA}{dt} = 9 - 2t - 11t^2$$

$$\Rightarrow 11t^2 + 2t - 9 = 0$$

$$11t^2 + 11t - 9t - 9 = 0$$

$$t = -1 \text{ \& } t = \frac{9}{11}$$

$$\therefore \text{largest area} = \frac{9}{11} \left(9 - \frac{11}{3}, \frac{81}{121} - \frac{9}{11}\right)$$

$$= \frac{9}{11} \cdot \frac{63}{11} = \frac{567}{121}$$

Q12. $f(x+y) = f(x) \cdot f(y)$

(1) $\Rightarrow f(x) = e^{\lambda x} f'(0) = 4a$

$$\Rightarrow f'(x) = \lambda e^{\lambda x} \Rightarrow \lambda = 4a$$

So, $f(x) = e^{4ax}$


Area Under Curves

JEE Main 2025 January

$$f''(x) - 3af'(x) - f(x) = 0$$

$$\Rightarrow \lambda^2 - 3a\lambda - 1 = 0$$

$$\Rightarrow 16a^2 - 12a^2 - 1 = 0 \Rightarrow 4a^2 = 1 \Rightarrow a = \frac{1}{2}$$



$$F(x) = e^{2x}$$

$$\text{Area} = \int_0^2 e^x dx = e^2 - 1$$



Q1. $\frac{dy}{dx} + 2y \tan x = \sin x$
(1) I.F. = $e^{2 \int \tan x dx} = \sec^2 x$

$$y \sec^2 x = \int \frac{\sin x}{\cos^2 x} dx$$

$$= \int \tan x \sec x dx$$

$$= \sec x + C$$

$$C = -2$$

$$y = \cos x - 2 \cos^2 x$$

$$y\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} - 1$$

$$y' = -\sin x + 4 \cos x \sin x$$

$$y'\left(\frac{\pi}{4}\right) = -\frac{1}{\sqrt{2}} + 2$$

$$y'\left(\frac{\pi}{4}\right) + y\left(\frac{\pi}{4}\right) = 1$$

Q2. $y dy = (x dy - y dx) \sin\left(\frac{x}{y}\right)$
(4)

$$\frac{dy}{y} = \left(\frac{x dy - y dx}{y^2}\right) \sin\left(\frac{x}{y}\right)$$

$$\frac{dy}{y} = \sin\left(\frac{x}{y}\right) d\left(-\frac{x}{y}\right)$$

$$\rightarrow \ln y = \cos \frac{x}{y} + C$$

$$x(1) = \frac{\pi}{2} \Rightarrow 0 = \cos \frac{\pi}{2} + C \Rightarrow C = 0$$

$$\ln y = \cos \frac{x}{y}$$

$$\text{but } y = 2 \Rightarrow \cos \frac{x}{2} = \ln 2$$

$$\cos x = 2 \cos^2 \frac{x}{2} - 1$$

$$= 2(\ln 2)^2 - 1$$

Q3. $y^2 dx + \left(x - \frac{1}{y}\right) dy = 0$
(3)

$$y^2 dx = \left(\frac{1}{y} - x\right) dy$$

$$\Rightarrow y^2 \frac{dx}{dy} = \frac{1}{y} - x$$

$$\Rightarrow \frac{dx}{dy} + \frac{x}{y^2} = \frac{1}{y^3}$$

$$\text{I.F.} = e^{\int \frac{1}{y^2} dy} = e^{-\frac{1}{y}}$$

$\therefore$ Solution is

$$x e^{-\frac{1}{y}} = \int e^{-\frac{1}{y}} \times \frac{1}{y^3} dy + C$$

Differential Equations

JEE Main 2025 January

$$\text{Let } \frac{-1}{y} = t$$

$$\Rightarrow \frac{1}{y^2} dy = dt$$

$$\Rightarrow x e^{-\frac{1}{y}} = - \int e^t dt + C$$

$$\Rightarrow x e^{-\frac{1}{y}} = -e^t(t-1) + C$$

$$\Rightarrow x e^{-\frac{1}{y}} = -e^{-\frac{1}{y}} \left(\frac{-1}{y} - 1 \right) + C$$

$$x(1) = 1$$

$$\Rightarrow e^{-1} = -e^{-1}(-2) + C$$

$$\Rightarrow C = -e^{-1}$$

$$\Rightarrow x = \frac{1}{y} + 1 - e^{-1+\frac{1}{y}}$$

$$x\left(\frac{1}{2}\right) = 3 - e$$

Q4. $\frac{dx}{dy} + \frac{x}{1+y^2} = \frac{2e^{\tan^{-1}y}}{1+y^2}$

(4)

$$\text{I.F.} = e^{\tan^{-1}y}$$

$$x e^{\tan^{-1}y} = \int \frac{2(e^{\tan^{-1}y})^2 dy}{1+y^2}$$

$$\text{Put } \tan^{-1}y = t, \frac{dy}{1+y^2} = dt$$

$$x e^{\tan^{-1}y} = \int 2e^{2t} dt$$

$$x e^{\tan^{-1}y} = e^{2\tan^{-1}y} + c$$

$$x = e^{\tan^{-1}y} + c e^{-\tan^{-1}y}$$

$$\because y = 0, x = 1$$

$$1 = 1 + c \Rightarrow c = 0$$

$$y = \frac{1}{\sqrt{3}}, x = e^{\pi/6}$$

Differential Equations

JEE Main 2025 January

Q5. $\frac{dy}{dx} = \frac{2(3+y) \cdot e^{2x}}{7+e^{2x}}$

(3) $\frac{dy}{dx} = \frac{2ye^{2x}}{7+e^{2x}} = \frac{6 \cdot e^{2x}}{7+e^{2x}}$

$$\frac{dy}{dx} = \frac{2ye^{2x}}{7+e^{2x}} = \frac{6 \cdot e^{2x}}{7+e^{2x}}$$

$$\text{I.F.} = e^{-\int \frac{2e^{2x}}{7+e^{2x}} dx}$$

$$\Rightarrow e^{-\ln(7+e^{2x})}$$

$$= \frac{1}{7+e^{2x}}$$

$$y \cdot \frac{1}{7+e^{2x}} = \int \frac{6e^{2x}}{(7+e^{2x})^2} dx$$

$$\frac{y}{7+e^{2x}} = \frac{-3}{7+e^{2x}+C}$$

$$\therefore y(0) = 5$$

$$\Rightarrow \frac{5}{8} = \frac{-3}{8} + C$$

$$\Rightarrow C = 1$$

$$\therefore y = -3 + 7 + e^{2x}$$

$$y = e^{2x} + 4$$

$$\therefore k = 8$$

Q6. $(1+x^2) \frac{dy}{dx} + xy = 5x^1 \sqrt{1+x^2}$

(2) $\frac{dy}{dx} + \frac{xy}{1+x^2} = \frac{5x^2}{\sqrt{1+x^2}}$

$$\therefore \text{I.F.} = e^{\int \frac{x}{1+x^2} dx} = e^{\frac{\ln(1+x^2)}{2}} = \sqrt{1+x^2}$$

$$\therefore y\sqrt{1+x^2} = \int \frac{5x^2}{\sqrt{1+x^2}} \cdot \sqrt{1+x^2} dx$$

$$\therefore y\sqrt{1+x^2} = \int \frac{5x^2}{\sqrt{1+x^2}} \cdot \sqrt{1+x^2} dx$$

$$y\sqrt{1+x^2} = \frac{5x^3}{3} + C$$

$$\therefore y(0) = 0 \Rightarrow 0 = 0 + C \Rightarrow C = 0$$

$$\therefore y = \frac{5x^3}{3\sqrt{1+x^2}}$$

$$y(\sqrt{3}) = \frac{15\sqrt{3}}{32} = \frac{5\sqrt{3}}{2}$$

Q7. $2(x+2)^2 f(x) - 3(x+2)^2 = 10 \int_0^x (t+2)f(t)dt$

(19) Differentiating both side

Differential Equations

JEE Main 2025 January

$$4(x+2)f(x) + 2f(x)(x+2)^2 - 6(x+2) = 10(x+2)f(x)$$

$$= (x+2) \frac{dy}{dx} - 3y = 3$$

$$\frac{1}{3} \int \frac{dy}{y+1} = \int \frac{dx}{x+2}$$

$$\ln|y+1| = 3 \ln|x+2| + \ln c$$

$$y+1 = (x+2)^3 c$$

$$\therefore y(0) = \frac{3}{2}$$

$$\Rightarrow \frac{5}{16} = c$$

$$\therefore y = \frac{5}{16}(x+2)^3 - 1$$

$$y(2) = \frac{5}{16} \times 64 - 1 = 19$$

Q8. $x^2 f'(x) - 2xf(x) = 3$

(1) $\left(\frac{x^2 f'(x) - 2xf(x)}{(x^2)^2} \right) = \frac{3}{(x^2)^2}$

$$\Rightarrow \frac{d}{dx} \left(\frac{f(x)}{x^2} \right) = \frac{3}{x^4}$$

Integrating both sides

$$\frac{f(x)}{x^2} = -\frac{1}{x^3} + C$$

$$f(x) = -\frac{1}{x} + Cx^2$$

put $x = 1$

$$4 = -1 + C \Rightarrow C = 5$$

$$f(x) = -\frac{1}{x} + 5x^2$$

$$\text{Now } 2 \times f(2) = 2 \times \left[-\frac{1}{2} + 5 \times 2^2 \right]$$

$$= 39$$

Q9. (4) $\frac{dy}{dx} + \frac{\left(\sin^{-1} \frac{x}{2}\right)}{\sqrt{4-x^2}} y = \frac{\left(\sin^{-3} \frac{x}{2}\right)^3}{\sqrt{4-x^2}}$

$$ye^{\frac{\left(\sin^{-1} \frac{x}{2}\right)^2}{2}} = \int \frac{\left(\sin^{-3} \frac{x}{2}\right)^3}{4-x^2} e^{\frac{\left(\sin^{-1} \frac{x}{2}\right)^2}{2}} dx$$


$$y = \left(\sin^{-1} \frac{x}{2}\right)^2 - 2 + c \cdot e^{-\frac{\left(\sin^{-1} \frac{x}{2}\right)^2}{2}}$$

$$y(2) = \frac{\pi^2}{4} - 2 \Rightarrow c = 0$$

$$y(0) = -2$$

Q10. I.F. $e^{-\frac{1}{2} \int \frac{2x}{1-x^2} dx} = e^{-\frac{1}{2} \ln(1-x^2)} = \sqrt{1-x^2}$

(27) $y \times \sqrt{1-x^2} = \int (x^6 + 4x) dx = \frac{x^7}{7} + 2x^2 + c$



Differential Equations

JEE Main 2025 January

$$\text{Given } y(0) = 0 \Rightarrow c = 0$$

$$y = \frac{\frac{x^7}{7} + 2x^2}{\sqrt{1-x^2}}$$

$$\text{Now, } 6 \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\frac{x^7}{7} + 2x^2}{\sqrt{1-x^2}} dx = 6 \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{2x^2}{\sqrt{1-x^2}} dx$$

$$= 24 \int_0^{\frac{1}{2}} \frac{x^2}{\sqrt{1-x^2}} dx$$

$$\text{Put } x = \sin \theta$$

$$dx = \cos \theta d\theta$$

$$= 24 \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta}{\cos \theta} \cos \theta d\theta$$

$$= 24 \int_0^{\frac{\pi}{6}} \left(\frac{1 - \cos 2\theta}{2} \right) d\theta = 12 \left[\theta - \frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{6}}$$

$$= 12 \left(\frac{\pi}{6} - \frac{\sqrt{3}}{4} \right)$$

$$= 2\pi - 3\sqrt{3}$$

$$\alpha^2 = (3\sqrt{3})^2 = 27$$

Q11. (1) $\frac{dy}{dx} - \frac{3 \sin x}{\cos x \ln \cos x} y = -\frac{\sin x}{\cos x (\ln \cos x)^2}$

$$\text{I.F.} = e^{-\int \frac{3 \tan x}{\ln \cos x} dx}$$

$$\text{Let } \ln \cos x = t$$

$$-\tan x dx = dt$$

$$e^{3 \int \frac{dt}{t}} = e^{3 \ln t} = t^3 = (\ln \cos x)^3$$

$\therefore$ Solution will be

$$y(\ln \cos x)^3 = - \int (\tan x)(\ln \cos x) dx$$

$$y(\ln \cos x)^3 = \frac{(\ln \cos x)^2}{2} + c$$

$$\therefore y\left(\frac{\pi}{4}\right) = -\frac{1}{\ln 2}$$

$$\Rightarrow c = 0$$

$$\therefore y = \frac{1}{2(\ln \cos x)}$$

$$y\left(\frac{\pi}{6}\right) = \frac{1}{2} \times \frac{1}{\ln\left(\cos \frac{\pi}{6}\right)}$$

$$= \frac{1}{2} \left[\frac{1}{\ln\left(\frac{\sqrt{3}}{2}\right)} \right]$$

$$= \frac{1}{2} \times \frac{1}{\ln \sqrt{3} - \ln 2}$$

$$= \frac{1}{\ln 3 - \ln 4}$$

Differential Equations

JEE Main 2025 January

Q12. If $e^{\int \tan x dx} = e^{\ln(\sec x)} = \sec x$

$$(4) \quad \therefore y \cdot \sec x = \int \left\{ \frac{2 + \sec x}{(1 + 2 \sec x)^2} \right\} \sec x dx$$

$$= \int \frac{2 \cos x + 1}{(\cos x + 2)^2} dx \quad \text{Let } \cos x = \frac{1 - t^2}{1 + t^2}$$

$$= \int \frac{2 \left(\frac{1 - t^2}{1 + t^2} \right) + 1}{\left(\frac{1 - t^2}{1 + t^2} + 2 \right)^2} 2dt$$

$$= \int \frac{2 - 2t^2 + 1 + t^2}{(1 - t^2 + 2 + 2t^2)^2} \times 2dt$$

$$= 2 \int \frac{3 - t^2}{(t^2 + 3)^2} dt$$

$$\text{Let } t + \frac{3}{t} = u$$

$$\left(1 - \frac{3}{t^2} \right) dt = du$$

$$= -2 \int \frac{du}{u^2}$$

$$y \cdot (\sec x) = \frac{2}{u} + c$$

$$y \cdot \sec x = \frac{2}{t + \frac{3}{t}} + c \dots (I)$$

$$\text{At } x = \frac{\pi}{3}, t = \tan \frac{x}{2} = \frac{1}{\sqrt{3}}$$

$$2 \cdot \frac{\sqrt{3}}{10} = \frac{2}{\frac{1}{\sqrt{3}} + 3\sqrt{3}} + c$$

$$2 \cdot \frac{\sqrt{3}}{10} = \frac{2\sqrt{3}}{10} + c \Rightarrow C = 0$$

$$\text{At } x = \frac{\pi}{4}, t = \tan \frac{x}{2} = \sqrt{2} - 1$$


$$\therefore y \cdot \sqrt{2} = \frac{2}{\sqrt{2} - 1 + \frac{3}{\sqrt{2} - 1}}$$

$$y \cdot \sqrt{2} = \frac{2(\sqrt{2} - 1)}{6 - 2\sqrt{2}}$$

$$y = \frac{\sqrt{2}(\sqrt{2} - 1)}{2(3 - \sqrt{2})} = \frac{1}{\sqrt{2}} \times \frac{2\sqrt{2} - 1}{7}$$

$$= \frac{4 - \sqrt{2}}{14}$$

Q1.
(4)



Equation of lines $QR = 5x + 2y + 2 = 0$

Equation of lines $PR = 10x - 3y - 38 = 0$


$\therefore$ Point $R(2, -6)$

Centroid $= \left(\frac{5 - 2 + 2}{3}, \frac{4 + 4 - 6}{3} \right)$

$= \left(\frac{5}{3}, \frac{2}{3} \right)$

$c + 2d = \frac{5}{3} + \frac{4}{3} = 3$

Q2. The centroid $G''(\alpha, \beta)$ of $\triangle PQR$ be image of centroid of given triangle $P'Q'R'$.
(4)



Centroid of $\triangle P'Q'R' = \left(\frac{1+3+2}{3}, \frac{3+1+4}{3} \right) = G' \left(2, \frac{8}{3} \right)$.

Image of $G \left(2, \frac{8}{3} \right)$, w.r.t. line $x + 2y = 2$ is (α, β)

Straight Lines

JEE Main 2025 January

$$\text{Then } \frac{\alpha-2}{1} = \frac{\beta-\frac{8}{3}}{2} = \frac{-2\left(2+\frac{16}{3}-2\right)}{1+4}$$


$$\therefore \frac{\alpha-2}{1} = \frac{\beta-\frac{8}{3}}{2} = -\frac{32}{15}$$

$$\therefore \alpha = -\frac{2}{15} \text{ and } \beta = -\frac{8}{5}$$

$$\text{Then } 15(\alpha - \beta) = 15\left(-\frac{2}{15} + \frac{24}{15}\right) = 22$$

Q3.

(145)



$$\frac{a+0}{3} = h \Rightarrow a = 3h$$

$$\frac{9+0}{3} = k \Rightarrow k = 3$$

$$\therefore (h, k) = \left(\frac{6 + 10 \cos \alpha - 10 \sin \alpha}{3}, \frac{8 - 10 \sin \alpha - 10 \cos \alpha}{3} \right)$$

$$6 + 10 \cos \alpha - 10 \sin \alpha = 3h$$

$$10 \cos \alpha - 10 \sin \alpha = 3h - 6$$

$$10(\cos \alpha - \sin \alpha) = 1$$

$$\frac{8 - 10 \sin \alpha + 10 \cos \alpha}{3} = k$$

$$\Rightarrow 100 \sin 2\alpha = 99$$

$$h = \frac{7}{3}$$


$$\Rightarrow a = 7$$

$$\text{Now, } 5a - 3h + 6k + 100 \sin 2\alpha = 35 - 7 + 18 + 99$$

$$= 145$$

Q4.

(2)



$\therefore$ centroid of $\triangle ABC = \left(\frac{9+3+5}{3}, \frac{11+4+13}{3} \right)$
 $= \left(\frac{17}{3}, \frac{28}{3} \right)$

Let image of centroid with respect to line mirror is

(h, k)

$$\therefore \left(\frac{k - \frac{28}{3}}{h - \frac{17}{3}} \right) \left(-\frac{1}{2} \right) = -1$$


$$\&3 \left(\frac{h + \frac{17}{3}}{2} \right) + 6 \cdot \left(\frac{\frac{k+28}{3}}{2} \right) = 53$$

Solving (1) \& (2) we get $h = 3, k = 4$

$$\therefore h^2 + k^2 + hk = 37$$

Q5.

(3)



Clearly, $x = \frac{11}{2}$ intersect $x + y - 11 = 0$ at $\left(\frac{11}{2}, \frac{11}{2} \right)$ and $2x + 3y - 29 = 0$ at $\left(\frac{11}{2}, 6 \right) \Rightarrow \alpha = \left[\frac{11}{2}, 6 \right]$


$$\alpha_{\min} \cdot \alpha_{\max} = \frac{11}{2} \cdot 6 = 33$$

Straight Lines

JEE Main 2025 January

Q6.

(3)



Slope of the third side = slope of the perpendicular bisector of given lines

$$h : \frac{-x+2y-4}{\sqrt{5}} = \pm \frac{x+y-4}{\sqrt{2}}$$

$$h_1 : \sqrt{2}(-x+2y-4) = \sqrt{5}(x+y-4)$$

$$h_2 : \sqrt{2}(-x+2y-4) = -\sqrt{5}(x+y-4)$$

$$M_{L_1} : -\left[\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - 2\sqrt{2}} \right]$$

$$M_{L_2} : -\left[\frac{\sqrt{5} - \sqrt{2}}{\sqrt{5} + 2\sqrt{2}} \right]$$


$$M_{L_1} + M_{L_2} = -\left[\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - 2\sqrt{2}} + \frac{\sqrt{5} - \sqrt{2}}{\sqrt{5} + 2\sqrt{2}} \right]$$

$$= -\left[\frac{(\sqrt{5} + \sqrt{2})(\sqrt{5} + 2\sqrt{2}) + (\sqrt{5} - \sqrt{2})(\sqrt{5} - 2\sqrt{2})}{-3} \right]$$

$$= 6$$

Q7.

(3)



$$\therefore PM = QM$$

$$\text{So, } M \left(\frac{\frac{57}{13} + 1}{2}, \frac{\frac{40}{13} + 2}{2} \right)$$

$$= \left(\frac{35}{13}, \frac{-7}{13} \right)$$

$\therefore$ M lies on the line

Straight Lines

JEE Main 2025 January

$$2x - 3y + \lambda = 0$$

$$2\left(\frac{35}{13}\right) - 3\left(\frac{-7}{13}\right) + \lambda = 0$$

$$\lambda = -\frac{70}{13} + \frac{21}{13}$$

$$= \frac{-91}{13} = -7$$

$$\begin{vmatrix} 3 & -4 & -\alpha \\ 8 & -11 & -33 \\ 2 & 3 & \lambda \end{vmatrix} = 0$$

$$\Rightarrow 3(-11\lambda - 99) + 4(8\lambda + 66) - \alpha(-24 + 22) = 0$$

$$\Rightarrow 33\lambda - 297 + 32\lambda + 264 + 24\alpha - 22\alpha = 0$$

$$\Rightarrow -\lambda + 2\alpha - 33 = 0$$

$$\therefore \lambda = -7$$

$$-(-7) + 2\alpha - 33 = 0$$

$$2\alpha = 26$$

$$\alpha = 13$$

$$\therefore |\alpha\lambda| = |13 \times (-7)|$$

$$= 91$$

Q8. ${}^nC_{r-1} = 28$

(4) ${}^nC_r = 56$


$${}^nC_{r+1} = 70$$

$$\frac{{}^nC_{r-1}}{{}^nC_r} = \frac{28}{56} \Rightarrow \frac{r}{n-r+1} = \frac{1}{2}$$

$$\frac{{}^nC_r}{{}^nC_{r+1}} = \frac{56}{70} \Rightarrow \frac{r+1}{n-r} = \frac{70}{56}$$

$$\left. \begin{array}{l} n = 8 = 3 \end{array} \right\}$$

$$A(4\cos t, 4\sin t)$$



$$h = \frac{4\cos t + 2\sin t + 1}{3} \quad k = \frac{4\sin t - 2\cos t}{3}$$

$$3h - 1 = 4\cos t + 2\sin t$$

$$3k - 1 = 4\sin t - 2\cos t$$

$$(1)^2 + (2)^2$$

$$(3h - 1)^2 + (3k)^2 = 20$$

$$\text{Locus of centroid: } (3x - 1)^2 + (3y)^2 = 20$$


$$\Rightarrow \alpha = 20$$

Straight Lines

JEE Main 2025 January

Q9.

(1)



$$\text{Area of } \triangle AOB = \frac{1}{2}$$

$$\text{Area of } \triangle AMN = \frac{4}{9} \times \frac{1}{2} = \frac{2}{9}$$

Equation of AB is $x + y = 1$

$$OA = 1, AM = \sec(45^\circ - \theta)$$

$$AN = \sec(45^\circ - \theta) \cos \theta$$

$$MN = \sec(45^\circ - \theta) \sin \theta$$

$$\text{Ar}(\triangle AMN) = \frac{1}{2} \times \sec^2(45^\circ - \theta) \sin \theta \cdot \cos \theta = \frac{2}{9}$$


$$\Rightarrow \tan \theta = 2, \frac{1}{2}$$

$\tan \theta = 2$ is rejected

$$\frac{AN}{NB} = \frac{\lambda}{1} = \cot \theta = 2$$

Q10. $AB = 8$

(3) $AB^2 = 64$



$$\Rightarrow (a - b)^2 + (b + 4)^2 = 64 \dots (1)$$

Now P divides AB in the ratio 2 : 1 internally

$$\Rightarrow h = \frac{2a+b}{3} \text{ and } k = \frac{-4+b+2}{3}$$

$$\Rightarrow 2a + b = 3h \dots (2) \quad k = \frac{b-2}{3}$$

$$\text{From equation (2) and (3)} \Rightarrow b = 3k + 2$$

Straight Lines

JEE Main 2025 January

$$\Rightarrow 2a = 3h - 3k - 2$$

$$\Rightarrow a = \frac{3h - 3k - 2}{2}$$

Now by putting value of a and b in equation

$$\Rightarrow \left(\frac{3h - 3k - 2}{2} - (3k + 2) \right)^2 + (3k + 2 + 4)^2 = 64$$

$$\Rightarrow \left(\frac{3h - 3k - 2 - 6k - 4}{2} \right)^2 + (3k + 6)^2 = 64$$

$$\Rightarrow (3h - 9k - 6)^2 + 4(3k + 6)^2 = 4 \times 64$$

$$\Rightarrow 9(h - 3k - 2)^2 + 36(k + 2)^2 = 256$$

$$\Rightarrow 9(h^2 + 9k^2 + 4 - 6hk - 4h + 12k)$$

$$+ 36(k^2 + 4 + 4k) = 256$$

$$\Rightarrow 9(h^2 + 13k^2 + 20 - 6hk - 4h + 28k) = 256$$


Replacing h by x and k by y

$$\Rightarrow 9(x^2 + 13y^2 - 6xy - 4x + 28y) + 180 - 256 = 0$$

$$\Rightarrow 9(x^2 + 13y^2 - 6xy - 4x + 28y) - 76 = 0$$

By comparing $\alpha = 13, \beta = -6, \gamma = -4$

$$\alpha - \beta - \gamma = 13 + 6 + 4 = 23$$




Circle

JEE Main 2025 January

Q1.

(4)



By pythagorus $r^2 = a^2 + \frac{b^2}{4} = P^2$

$$r = \sqrt{\frac{4a^2 + b^2}{4}}$$

Equation of circle is $(x - \alpha)^2 + (y - \beta)^2 = r^2$

$$x^2 + y^2 - 2ax - 2py + \alpha^2 + \beta^2 - r^2 = 0$$

$$\text{comparision } x^2 + y^2 - \alpha x + \beta y + r = 0$$

$$-\alpha = -2a, \beta = -2p, r = a^2$$

$$\Rightarrow 2a = \alpha, 4a^2 + b^2 = 4p^2$$


$$\alpha^2 + b^2 = 4p^2$$

$$\alpha^2 + b^2 = \beta^2$$

$$\text{So, } (2a, b^2) = (\alpha, \beta^2 - 4r)$$

Q2.

(3)



By solving $x = y$ with circle We get

$$C(\sqrt{2}, \sqrt{2})$$

$$D(-\sqrt{2}, -\sqrt{2})$$

By solving $x + y = 1$ with circle $x^2 + y^2 = 4$

we set

Circle

JEE Main 2025 January

$$A \left(\frac{1 + \sqrt{7}}{2}, \frac{1 - \sqrt{7}}{2} \right)$$

$$\& B \left(\frac{1 - \sqrt{7}}{2}, \frac{1 + \sqrt{7}}{2} \right)$$

$\therefore$ Area of Quadrilateral ACBD

$$= 2 \times \text{Area of } \triangle BCD$$

$$= 2 \times \frac{1}{2} \begin{vmatrix} \sqrt{2} & \sqrt{2} & 1 \\ \frac{1-\sqrt{7}}{2} & \frac{1+\sqrt{7}}{2} & 1 \\ -\sqrt{2} & -\sqrt{2} & 1 \end{vmatrix}$$

$$= 2\sqrt{14}$$

Q3. Let the centre be

$$(3) \left(-2a, \frac{6a-2}{2} \right) \equiv (-2a, 3a-1)$$


Centre is equal distance from $(4, 2)$ and $(0, 2)$

$$\Rightarrow \sqrt{(4+2a)^2 + (3a-3)^2} = \sqrt{(-2a-0)^2 + (3a-3)^2}$$

$$\Rightarrow (2a+4)^2 + 9(a-1)^2 = 4a^2 + 9(a-1)^2$$

$$\Rightarrow 4a^2 + 16 + 16a = 4a^2 \Rightarrow a = -1$$

$$\Rightarrow \text{centre} \equiv (2, -4) \Rightarrow \text{Radius} = \sqrt{40}$$



$$\Rightarrow AM^2 = (\sqrt{40})^2 - (\sqrt{37})^2$$


$$\Rightarrow 2AM = AB = 2\sqrt{3}$$

Circle

JEE Main 2025 January

Q4.

(2)



$$C_1 C_2 = \sqrt{(2+2)^2 + (5-2)^2}$$

$$= \sqrt{16+9}$$

$$= 5$$

$$r + 2 > 5$$

$$r > 3$$

$$r < 5 + 2$$


$$r < 7$$

$$\therefore \alpha = 3, \beta = 7$$

$$3\beta - 2\alpha = 3(7) - 2(3) = 21 - 6 = 15$$

Q5.

(2)



$$\text{Centre } (1, -2), r = 3$$


$$\text{Reflection of } (1, -2) \text{ about } 2x - 3y + 5 = 0$$

$$\frac{x-1}{2} = \frac{y+2}{-3} = \frac{-2(2+6+5)}{13} = -2$$

$$x = -3, y = 4$$

Equation of circle 'C'

$$C : (x+3)^2 + (y-4)^2 = 9$$



$$AB = 3\sqrt{3}$$

A.T.Q.

$$\ell(\text{arc AB}) = \frac{1}{6} \times 2\pi r$$

$$r\theta = \frac{1}{6} \times 2\pi r$$

$$\theta = \frac{\pi}{3}$$

$$(\alpha + 6)^2 + (\beta - 4)^2 = 27$$

$$(\alpha + 3)^2 + (\beta - 4)^2 = 9$$

$$\frac{(\alpha + 6)^2 + (\beta - 4)^2 - (\alpha + 3)^2 - (\beta - 4)^2}{2} = 18$$

$$\Rightarrow 6\alpha = -9$$

$$\Rightarrow \alpha = -\frac{3}{2}, \beta = \left(4 - \frac{3\sqrt{3}}{2}\right)$$

$$\therefore \beta - \sqrt{3}\alpha$$

$$\left(4 - \frac{3\sqrt{3}}{2}\right) + \frac{3\sqrt{3}}{2}$$

$$= 4$$

Parabola

JEE Main 2025 January

Q1. $y^2 = 4(x + 4)$

(15) Equation of circle

$$(x + 3)^2 + y^2 = 25$$

Passes through the point of intersection of two lines $3x - y = 0$ and $x + \lambda y = 4$

$\left(\frac{4}{3\lambda+1}, \frac{12}{3\lambda+1}\right)$, we get


$$\lambda = -\frac{7}{6}, 1$$

$$12\lambda_1 + 29\lambda_2$$

$$-14 + 29 = 15$$

Q2.

(2)



$\therefore$ Equation of parabola is $PS = (1)PM$

$$\Rightarrow PS^2 = PM^2$$

$$(x - 3)^2 + (y - 6)^2 = \left(\frac{x + 2y}{\sqrt{5}}\right)^2$$

$$5x^2 - 30x + 5y^2 - 60y + 225 = x^2 + 4y^2 + 4xy$$

$$4x^2 + y^2 - 4xy - 30x - 60y + 225 = 0$$

We get : $\alpha = 4, \beta = 1, \gamma = 4$

$$\therefore \alpha + \beta + \gamma = 9$$

Q3. The parabolas are

(3) $(x - 4)^2 + (y - 3)^2 = x^2 \dots(i)$

and $(x - 4)^2 + (y - 3)^2 = y^2 \dots$

If point of intersection are $A(x_1, y_1)$ and $B(x_2, y_2)$ By solving (i) and (ii), we get

$$x_1 + x_2 = 14 \text{ and } x_1 x_2 = 25$$

$$(AB)^2 = 2 \left((x_1 + x_2)^2 - 4x_1 x_2 \right) = 192$$

Q4. $y = x^2 + px - 3$

(3) Let $P(\alpha, 0), Q(\beta, 0), R(0, -3)$

Circle with centre $(-1, -1)$ is $(x + 1)^2 + (y + 1)^2 = r^2$

Passes through $(0, -3)$

$$1^2 + (-2)^2 = r^2$$

$$r^2 = 5$$

Parabola

JEE Main 2025 January

$$(x+1)^2 + (y+1)^2 = 5$$

Put $y = 0$

$$(x+1)^2 = 5 - 1$$

$$(x+1)^2 = 4$$

$$x+1 = \pm 2$$


$$x = 1 \text{ or } x = -3$$

$$\therefore P(1, 0) \text{ and } Q(-3, 0)$$

$$\text{Area of } \triangle PQR = \frac{1}{2} \begin{vmatrix} 1 & 0 & 1 \\ -3 & 0 & 1 \\ 0 & -3 & 1 \end{vmatrix} = 6$$

Q5.

(2)



Normal at P

$$y + tx = 2t + t^3$$

↑

$$(a, 0)$$

$$at = 2t + t^3$$

$$a = 2 + t^2$$

$$R(2 + t^2, 0)$$

$$PR = 4 \Rightarrow 4 + 4t^2 = 16$$

$$4t^2 = 12 \Rightarrow t^2 = 3$$

$$a = 5 \Rightarrow R(5, 0)$$

Focus (1, 0)

(1, 0) & (5, 0) will be the end pts. of diameter

$\Rightarrow$ Eqⁿ of circle is

$$(x-1)(x-5) + y^2 = 0$$


$$x^2 + y^2 - 6x + 5 = 0$$

Parabola

JEE Main 2025 January

Q6.

(14)



To find image of $P(t^2, 2t)$

$$\frac{x - t^2}{1} = \frac{y - 2t}{1} = \frac{-2(t^2 + 2t + 4)}{1^2 + 1^2} = -(t + 1)^2 - 3$$

$$x = t^2 - (t + 1)^2 - 3 = -2t - 4$$

$$y = 2t - (t + 1)^2 - 3 = -t^2 - 4$$

$$t = \frac{-x - 4}{2}$$

$$\Rightarrow y + 4 = -\left(\frac{-x - 4}{2}\right)^2$$

$$\Rightarrow (y + 4) = -\frac{(x + 4)^2}{4}$$

$$\Rightarrow x^2 = -4y$$

$$\Rightarrow \text{Focus } (-4, -5)$$

Also, $y = -5$ intersect

$$\therefore (-4)(-1) = (x + 4)^2$$

$$4 = (x + 4)^2$$

$$x + 4 = \pm 2$$

$$x = -2, -6$$

$$\Rightarrow d = 4$$

$$a = \frac{1}{2} \begin{vmatrix} 1 & 0 & 1 \\ -2 & -5 & 1 \\ -6 & -5 & 1 \end{vmatrix}$$

$$= \frac{1}{2} |1(-5 + 5) + 1(10 - 30)|$$

$$= \frac{1}{2} (20)$$

$$a = 10$$


$$\therefore a + d = 14$$

Parabola

JEE Main 2025 January

Q7.

(2)



$$3x - 2y + 12 = 0$$

$$4y = 3x^2$$

$$\therefore 2(3x + 12) = 3x^2$$

$$\Rightarrow x^2 - 2x - 8 = 0$$

$$\Rightarrow x = -2, 4$$


$$m_{OA} = -3/2, m_{OB} = 3$$

$$\tan \theta = \left(\frac{\frac{-3}{2} - 3}{1 - \frac{9}{2}} \right) = \frac{9}{7}$$

$$\theta = \tan^{-1} \left(\frac{9}{7} \right) \text{ (angle will be acute)}$$

Q8.

(1)



Parabola

JEE Main 2025 January

$$A(at_1^2, 2at) \text{ \& } C\left(\frac{a}{t_1^2}, -\frac{2a}{t_1}\right)$$


$$\text{Length AC} = a\left(t_1 + \frac{1}{t_1}\right)^2 = \frac{25}{4}, t_1 + \frac{1}{t_1} = \pm \frac{5}{2}$$

$$\Rightarrow t_1 = 2 \text{ or } \frac{1}{2}, A\left(\frac{1}{2}, 1\right), D\left(\frac{1}{4}, -1\right), B(4, 4), C(4, -4)$$

$$\text{So, area of trapezium} = \frac{1}{2}(8+2)\left(4 - \frac{1}{4}\right) = \frac{75}{4}$$

Q9.

(4)



$$(4, 4\sqrt{3}) \text{ lies on } y^2 = 4ax$$

$$\Rightarrow 48 = 4a \cdot 4$$

$$4a = 12$$

$$\Rightarrow y^2 = 12x \text{ is equation of parabola}$$

$$\text{Now, parameter of } P \text{ is } t_1 = \frac{2}{\sqrt{3}} \Rightarrow \text{Parameters of } Q \text{ is } t_2 = -\frac{\sqrt{3}}{2} \Rightarrow Q\left(\frac{9}{4}, -3\sqrt{3}\right)$$

$$\text{Area of trapezium PQNM}$$

$$= \frac{1}{2}MN \cdot (PM + QN)$$

$$= \frac{1}{2}MN \cdot (PS + QS)$$

$$= \frac{1}{2}MN \cdot PQ$$

$$= \frac{1}{2}7\sqrt{3} \cdot \frac{49}{4} = (343)\frac{\sqrt{3}}{8} = 3$$

Q10. $y^2 = 12x$ $a = 3$ $SP \times SQ = \frac{147}{4}$

(1328) Let $P(3t^2, 6t)$ and $t_1 t_2 = -1$

(ends of focal chord)

$$\text{So, } Q\left(\frac{3}{t^2}, \frac{-6}{t}\right)$$

Parabola

JEE Main 2025 January

$$S(3, 0)$$

$$SP \times SQ = PM_1 \times QM_2$$

(dist. from directrix)

$$= (3 + 3t^2) \left(3 + \frac{3}{t^2} \right) = \frac{147}{4}$$

$$\Rightarrow \frac{(1 + t^2)^2}{t^2} = \frac{49}{12}$$

$$t^2 = \frac{3}{4}, \frac{4}{3}$$

$$t = \pm \frac{\sqrt{3}}{2}, \pm \frac{2}{\sqrt{3}}$$

considering $t = \frac{-\sqrt{3}}{2}$

$$P\left(\frac{9}{4}, -3\sqrt{3}\right) \text{ and } Q(4, 4\sqrt{3})$$


Hence, diametric circle:

$$(x - 4) \left(x - \frac{9}{4} \right) + (y + 3\sqrt{3})(y - 4\sqrt{3}) = 0$$

$$\Rightarrow x^2 + y^2 - \frac{25}{4}x - \sqrt{3}y - 27 = 0$$

$$\Rightarrow \alpha = 400, \beta = 1728$$

$$\beta - \alpha = 1328$$



Q1. Product of focal distances = $(a + ex_1)(a - ex_1)$

$$(2) \quad = a^2 - e^2 x_1^2 = a^2 - e^2(3) \\ = a^2 - 3e^2 = \frac{7}{4} \Rightarrow a^2 = \frac{7}{4} + 3e^2 \\ \Rightarrow 4a^2 = 7 + 12e^2 \\ \& \left(\sqrt{3}, \frac{1}{2} \right) \text{ lines on } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\therefore \frac{3}{a^2} + \frac{1}{4b^2} = 1$$

$$\frac{3}{a^2} + \frac{1}{4(a^2)(1-e^2)} = 1$$

$$12(1-e^2) + 1 = 4a^2(1-e^2)$$

$$13 - 12e^2 = (7 + 12e^2)(1 - e^2)$$

$$\Rightarrow 13 - 12e^2 = 7 - 7e^2 + 12e^2 - 12e^4$$

$$\Rightarrow 12e^4 - 17e^2 + 6 = 0$$

$$\therefore e^2 = \frac{17 \pm \sqrt{289 - 288}}{24} = \frac{17 \pm 1}{24} = \frac{3}{4} \& \frac{2}{3}$$

$$\therefore e = \frac{\sqrt{3}}{2} \& \sqrt{\frac{2}{3}}$$

$$\therefore \text{difference} = \frac{\sqrt{3}}{2} - \sqrt{\frac{2}{3}} = \frac{3 - 2\sqrt{2}}{2\sqrt{3}}$$

Q2. $2ae = 4$

$$(4) \quad \Rightarrow a = 2\sqrt{3}$$

$$\Rightarrow 1 - \frac{b^2}{12} = \frac{1}{3} \Rightarrow b^2 = 8$$

$$\frac{2b^2}{a} \times \frac{2A^2}{B} = \frac{32}{\sqrt{3}}$$

$$\Rightarrow$$

$$\frac{2 \times 8}{2\sqrt{3}} \times \frac{2A^2}{B} = \frac{32}{\sqrt{3}}$$

$$\Rightarrow$$

$$\frac{A^2}{B} = 2 \Rightarrow A^2 = 2B$$

$$1 - \frac{A^2}{B} = \frac{1}{3}$$

$$\Rightarrow$$

$$B = 3 \Rightarrow A^2 = 6$$

$$E_1 : \frac{x^2}{12} + \frac{y^2}{8} = 1 \dots (i)$$

$$E_1 : \frac{x^2}{6} + \frac{y^2}{9} = 1 \dots (ii)$$

On solving (i) \& (ii)

Ellipse

JEE Main 2025 January

$$(x, y) = \left(\frac{\sqrt{6}}{\sqrt{5}}, \frac{6}{\sqrt{5}} \right), \left(\frac{-\sqrt{6}}{\sqrt{5}}, \frac{6}{\sqrt{5}} \right), \left(\frac{\sqrt{6}}{\sqrt{5}}, \frac{-6}{\sqrt{5}} \right), \left(\frac{-\sqrt{6}}{\sqrt{5}}, \frac{-6}{\sqrt{5}} \right)$$

Four points are vertices of rectangle area = $\frac{24\sqrt{6}}{5}$

Q3. $T = S_1$

$$(3) \quad \frac{x \cdot 1}{4} + \frac{y}{4} = \frac{1}{4} + \frac{1}{8}$$

$$\Rightarrow 2x + 2y = 3$$

$$\frac{x^2}{4} + \frac{\left(\frac{3-2x}{2}\right)^2}{2} = 1$$

$$\Rightarrow x = \frac{12 \pm \sqrt{120}}{12} \Rightarrow y = \frac{1}{2} \mp \frac{\sqrt{120}}{12}$$

So length of chord

$$= \frac{2\sqrt{15}}{3}$$

Q4. Equation of chord with given middle point

(1) $T = S_1$


$$\Rightarrow \frac{3x}{25} + \frac{y}{16} - 1 = \frac{9}{25} + \frac{1}{16} - 1$$

$$48x + 25y = 144 + 25$$

$$48x + 25y = 169 \text{ Ans.}$$

Q5.

(2)



$$E: \frac{x^2}{9} + \frac{y^2}{4} = 1$$

$T = S_1$

$$\Rightarrow \frac{\sqrt{2}x}{9} + \frac{1}{4} \left(\frac{4}{3}y \right) - 1 = \frac{2}{9} + \frac{16}{9(4)} - 1$$

$$\frac{\sqrt{2}x}{9} + \frac{y}{3} = \frac{2}{9} + \frac{4}{9}$$

$$\frac{\sqrt{2}x}{9} + \frac{y}{3} = \frac{2}{3} \Rightarrow \sqrt{2}x + 3y = 6$$

Now point of intersection of chord and ellipse is

Ellipse

JEE Main 2025 January

$$\frac{(6-3y)^2}{18} + \frac{y^2}{4} = 1$$

$$\frac{(2-y)^2}{2} + \frac{y^2}{4} = 1$$

$$2(4+y^2-4y) + y^2 = 4$$

$$\Rightarrow 3y^2 - 8y + 4 = 0$$

$$\Rightarrow y = 2, \frac{2}{3}$$

So, points are $(0, 2)$ are $(2\sqrt{2}, \frac{2}{3})$

$$\text{Length of chord} = \sqrt{(2\sqrt{2})^2 + \left(\frac{2}{3} - 2\right)^2}$$


$$= \sqrt{8 + \frac{16}{9}}$$

$$= \frac{\sqrt{88}}{3} = \frac{2\sqrt{22}}{3}$$

On comparing $\alpha = 22$

Q6.

(1)



Equation of chord $T = S_1$

$$\frac{5}{2} \left(\frac{x}{9} \right) + \frac{1}{2} \left(\frac{y}{4} \right) = \frac{25}{36} + \frac{1}{16}$$

$$\Rightarrow \frac{5x}{18} + \frac{y}{8} = \frac{100+9}{144} = \frac{109}{144}$$


$$\Rightarrow 40x + 18y = 109$$

$$\Rightarrow \alpha = 40, \beta = 18$$

$$\Rightarrow \alpha + \beta = 58$$

Q7.

(54)



$$E_1: \frac{x^2}{9} + \frac{y^2}{4} = 1 \Rightarrow e = \sqrt{1 - \frac{4}{9}} = \frac{\sqrt{5}}{3}$$

$$E_2: \frac{x^2}{a^2} + \frac{y^2}{4} = 1$$

$$e = \frac{\sqrt{5}}{3} = \sqrt{1 - \frac{a^2}{4}} \Rightarrow \frac{5}{9} = 1 - \frac{a^2}{4}$$

$$a^2 = \frac{16}{9}$$

$$E_2: \frac{x^2}{\frac{16}{9}} + \frac{y^2}{4} = 1$$

$$E_3: \frac{x^2}{\frac{16}{9}} + \frac{y^2}{b^2} = 1$$

$$e = \frac{\sqrt{5}}{3} = \sqrt{1 - \frac{b^2}{\frac{16}{9}}} \Rightarrow b^2 = \frac{64}{81}$$

$$E_3: \frac{x^2}{\frac{16}{9}} + \frac{y^2}{\frac{64}{81}} = 1$$

$$A_1 = \pi \times 3 \times 2 \Rightarrow 6\pi$$

$$A_2 = \pi \times \frac{4}{3} \times 2 = \frac{8\pi}{3}$$


$$A_3 = \pi \times \frac{4}{3} \times \frac{8}{9} = \frac{32\pi}{81}$$

$$\sum_{i=1}^{\infty} A_i = 6\pi + \frac{8\pi}{3} + \frac{32\pi}{81} + \dots \Rightarrow \frac{6\pi}{1 - \frac{4}{9}} \Rightarrow \frac{54\pi}{5}$$

$$\therefore \frac{5}{\pi} \sum_{i=1}^{\infty} A_i \Rightarrow \frac{5}{\pi} \times \frac{54\pi}{5} = 54$$

Q1.

(4)



$$be = 13, b = 5$$

$$a^2 = b^2 (e^2 - 1)$$

$$= b^2 e^2 - b^2$$

$$= 169 - 25 = 144$$

$$\ell(\text{LR}) = \frac{2a^2}{b} = \frac{2 \times 144}{5} = \frac{288}{5}$$

Q2. $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

(55) $\frac{2b^2}{a} = 15\sqrt{2} \dots (i)$

$$\sqrt{1 + \frac{b^2}{a^2}} = \sqrt{\frac{5}{2}} \dots (ii)$$

From (i) and (ii)

$$a = 5\sqrt{2} \text{ and } b^2 = 75$$

$$\frac{x^2}{A^2} - \frac{y^2}{B^2} = -1$$

$$\frac{2A^2}{B} = 12\sqrt{5} \dots (iii)$$

Since, product of transverse axis is $= 100\sqrt{10}$

$$(2A) \cdot (2B) = 100\sqrt{10}$$

From (iii) and (iv)

$$A^2 = 150 \text{ and } B = 5\sqrt{5}$$

$$e_2 = \sqrt{1 + \frac{A^2}{B^2}} = \sqrt{\frac{11}{5}}$$

$$\therefore 25e_2^2 = 25 \left(\frac{11}{5} \right) = 55$$

Hyperbola

JEE Main 2025 January

Q3. C: $x^2 + y^2 - 8x = 0$ H: $\frac{x^2}{9} - \frac{y^2}{4} = 1$

(3) By solving $\frac{x^2}{9} - \left(\frac{8x-x^2}{4}\right) = 1$

$$4x^2 - 72x + 9x^2 = 36$$

$$\Rightarrow 13x^2 - 72x - 36 = 0$$

$$\Rightarrow 13x^2 - 78x + 6x - 36 = 0$$

$$\Rightarrow 13x(x-6) + 6(x-6) = 0$$

$$\Rightarrow x = 6 \text{ or } -\frac{13}{6} \times \text{neglected}$$

$$\Rightarrow y^2 = 8(6) - (6)^2$$

$$\Rightarrow y = \pm\sqrt{12}$$

So, points A and B are $(6, \sqrt{12}), (6, -\sqrt{12})$

$$P\left(h, \frac{2h+4}{3}\right)$$

Centroid of $\triangle PAB$ is $\left(\frac{12+h}{3}, \frac{2h+4}{9}\right)$

By options this centroid lies on the line $6x - 9y = 20$

Q4. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ foci are $(ae, 0)$ and $(-ae, 0)$

(3) $\frac{x^2}{A^2} + \frac{y^2}{B^2} = 1$ foci are $(Ae', 0)$ and $(-Ae', 0)$

$$\Rightarrow 2ae = 2\sqrt{3} \Rightarrow ae = \sqrt{3}$$

$$\text{and } 2Ae' = 2\sqrt{3} \Rightarrow Ae' = \sqrt{3}$$

$$\Rightarrow ae = Ae' \Rightarrow \frac{e}{e'} = \frac{A}{a}$$

$$\Rightarrow \frac{1}{3} = \frac{A}{a} \Rightarrow a = 3A$$

$$\text{Now } a - A = 2 \Rightarrow a - \frac{a}{3} = 2 \Rightarrow a = 3 \text{ and } A = 1$$

$$Ae = \sqrt{3} \Rightarrow e = \frac{1}{\sqrt{3}} \text{ and } e' = \sqrt{3}$$

$$b^2 = a^2(1 - e^2)$$


$$b^2 = 6$$

$$\text{and } B^2 = A^2((e')^2 - 1) = (2) \Rightarrow B^2 = 2$$

$$\text{sum of LR} = \frac{2b^2}{a} + \frac{2B^2}{A} = 8$$

Q5.

(19)



$$r = \left| \frac{a+1}{\sqrt{2}} \right| \Rightarrow (a+1)^2 = 2r^2$$

Hyperbola

JEE Main 2025 January

$$\text{Also } \left(\frac{3a-1}{\sqrt{13}}\right)^2 + \left(\frac{2}{\sqrt{13}}\right)^2 = r^2$$
$$\Rightarrow \left(\frac{3a-1}{\sqrt{13}}\right)^2 + \frac{4}{13} = \frac{(a+1)^2}{2}$$

$$5a^2 - 14a - 3 = 0$$

$$\therefore a = -\frac{1}{5}, 3$$

$$\therefore a \neq -\frac{1}{5} \Rightarrow$$

$$\Rightarrow r = 2\sqrt{2}$$

$$\therefore \text{One focus of } \frac{x^2}{\alpha^2} - \frac{y^2}{\beta^2} = 1 \text{ is } (3, 0)$$

$$\Rightarrow \alpha e = 3 \text{ and } 2\alpha = 4\sqrt{2}$$

$$\Rightarrow \alpha = 2\sqrt{2} \Rightarrow \alpha^2 = 8$$

$$\alpha^2 \left[1 + \frac{\beta^2}{\alpha^2}\right] = 9$$

$$\alpha^2 + \beta^2 = 9$$

$$\Rightarrow \beta^2 = 1$$

$$\therefore 2\alpha^2 + 3\beta^2 = 19$$



Trigonometric Ratios

JEE Main 2025 January

Q1. $\sin x + \sin^2 x = 1$

(4) $\Rightarrow \sin x = \cos^2 x \Rightarrow \tan x = \cos x$

$\therefore$ Given expression

$$= 2 \cos^{12} x + 6 [\cos^{10} x + \cos^8 x] + 2 \cos^6 x$$

$$= 2 [\sin^6 x + 3 \sin^5 x + 3 \sin^4 x + \sin^3 x]$$

$$= 2 \sin^3 x [(\sin x + 1)^3]$$

$$= 2 [\sin^2 x + \sin x]^3$$

$$= 2$$

Q2. $(\sin 70^\circ)(\cot 10^\circ \cot 70^\circ - 1)$

(2) $= \sin 70^\circ \cot 10^\circ \cot 70^\circ - \sin 70^\circ$

$$= \cot 10^\circ \cos 70^\circ - \sin 70^\circ$$

$$= \frac{\cos 10^\circ \cos 70^\circ - \sin 70^\circ \sin 10^\circ}{\sin 10^\circ}$$

$$= \frac{\cos(10^\circ + 70^\circ)}{\sin 10^\circ}$$

$$= \frac{\cos 80^\circ}{\sin 10^\circ} = 1$$

Q3. $\frac{12}{13} \cos x + \frac{5}{13} \sin x$

(3) Let $\tan \alpha = \frac{5}{12}, \alpha \in \left(0, \frac{\pi}{2}\right)$

$$\Rightarrow \sin \alpha = \frac{5}{13}, \cos \alpha = \frac{12}{13}$$

$$\Rightarrow \frac{12}{13} \cos x + \frac{5}{13} \sin x = \cos \alpha \cos x + \sin \alpha \sin x$$

$$= \cos(x - \alpha)$$

$$\Rightarrow \cos^{-1}[\cos(x - \alpha)] = x - \alpha$$

$$= x - \tan^{-1}\left(\frac{5}{12}\right)$$

Q4. $\frac{1}{\sin \frac{\pi}{6}} \sum_{r=1}^{13} \frac{\sin\left[\left(\frac{\pi}{4} + \frac{r\pi}{6}\right) - \left(\frac{\pi}{4}\right) - (r-1)\frac{\pi}{6}\right]}{\sin\left(\frac{\pi}{4} + (r-1)\frac{\pi}{6}\right) \sin\left(\frac{\pi}{4} + \frac{r\pi}{6}\right)}$

(4) $\frac{1}{\sin \frac{\pi}{6}} \sum_{r=1}^{13} \left(\cot\left(\frac{\pi}{4} + (r-1)\frac{\pi}{6}\right) - \cot\left(\frac{\pi}{4} + \frac{r\pi}{6}\right) \right)$

$$= 2\sqrt{3} - 2 = a\sqrt{3} + b$$

$$\text{So } a^2 + b^2 = 8$$

Trigonometric Equations

JEE Main 2025 January

Q1. $2 \sin^2 \theta = \cos 2\theta$

(3) $2 \sin^2 \theta = 1 - 2 \sin^2 \theta$

$$4 \sin^2 \theta = 1$$

$$\sin^2 \theta = \frac{1}{4}$$

$$\sin \theta = \pm \frac{1}{2}$$

$$2 \cos^2 \theta = 3 \sin \theta$$

$$2 - 2 \sin^2 \theta + 3 \sin \theta - 2 = 0$$

$$(2 \sin \theta - 1)(2 \sin \theta - 2) = 0$$

$$\sin \theta = \frac{1}{2}$$

so common equation which satisfy both equations is $\sin \theta = \frac{1}{2}$

$$\theta = \frac{\pi}{6}, \frac{5\pi}{6} \quad (\theta \in [0, 2\pi])$$

$$\text{Sum} = \pi$$



Inverse Trigonometric Functions

JEE Main 2025 January

Q1. Let $\tan^{-1} \alpha = A \Rightarrow \tan A = \alpha$

(14) $\cot^{-1} \beta = B \Rightarrow \cot B = \beta$

$$\sec^2 A + \operatorname{cosec}^2 B = 36$$

$$\Rightarrow 1 + \tan^2 A + 1 + \cot^2 B = 36$$

$$\Rightarrow \alpha^2 + \beta^2 = 34$$

Also $\alpha + \beta = 8$ (Given)

$$\therefore (\alpha + \beta)^2 = 34 + 2\alpha\beta = 64$$

$$\Rightarrow \alpha\beta = 15$$

$\Rightarrow \alpha, \beta$ are roots of equation

$$x^2 - 8x + 15 = 0$$

$$\Rightarrow (x - 3)(x - 5) = 0$$

$$\Rightarrow x = 3, 5$$

$$\therefore \alpha = 3, \beta = 5 \quad (\alpha < \beta)$$

$$\therefore \alpha^2 + \beta = 9 + 5 = 14$$

Q2. $\cos\left(\sin^{-1} \frac{3}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{33}{65}\right)$

(2) $\cos\left(\tan^{-1} \frac{3}{4} + \tan^{-1} \frac{5}{12} + \tan^{-1} \frac{33}{56}\right)$

$$\cos\left(\tan^{-1}\left(\frac{\frac{3}{4} + \frac{5}{12}}{1 + \frac{3}{4} \cdot \frac{5}{12}}\right) + \tan^{-1} \frac{33}{56}\right)$$

$$\cos\left(\tan^{-1} \frac{56}{33} + \cot^{-1} \frac{56}{33}\right)$$

$$\cos\left(\frac{\pi}{2}\right) = 0$$

Q3. $16(\sec^{-1} x)^2 + (\operatorname{cosec}^{-1} x)^2$

(2) $\sec^{-1} x = a \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$

$$\operatorname{cosec}^{-1} x = \frac{\pi}{2} - a$$

$$= 16\left[a^2 + \left(\frac{\pi}{2} - a\right)^2\right] = 16\left[2a^2 - \pi a + \frac{\pi^2}{4}\right]$$

$$\max]_{a=\pi} = 16\left[2\pi^2 - \pi^2 + \pi \frac{2}{4}\right] = 20\pi^2$$

$$\min]_{a=\frac{\pi}{4}} = 16\left[\frac{2 \times \pi^2}{16} - \frac{\pi^2}{4} + \frac{\pi^2}{4}\right] = 2\pi^2$$

$$\text{Sum} = 22\pi^2$$

Q4. $\Rightarrow \cot^{-1}\left(\frac{\alpha\beta + 1}{\alpha - \beta}\right) + \cot^{-1}\left(\frac{\beta\gamma + 1}{\beta - \gamma}\right) + \cot^{-1}\left(\frac{\alpha\gamma + 1}{\gamma - \alpha}\right)$

(1) $\Rightarrow \tan^{-1}\left(\frac{\alpha - \beta}{1 + \alpha\beta}\right) + \tan^{-1}\left(\frac{\beta - \gamma}{1 + \beta\gamma}\right) + \pi + \tan^{-1}\left(\frac{\gamma - \alpha}{1 + \gamma\alpha}\right)$

$$\Rightarrow (\tan^{-1} \alpha - \tan^{-1} \beta) + (\tan^{-1} \beta - \tan^{-1} \gamma) + (\pi + \tan^{-1} \gamma - \tan^{-1} \alpha)$$

$$\Rightarrow \pi$$

Inverse Trigonometric Functions

JEE Main 2025 January

Q5. $\cos^{-1} x = \pi + \sin^{-1} x + \sin^{-1}(2x + 1)$

(5) $2 \cos^{-1} x - \sin^{-1}(2x + 1) = \frac{3\pi}{2}$

$$2\alpha - \beta = \frac{3\pi}{2} \text{ where } \cos^{-1} x = \alpha, \sin^{-1}(2x + 1) = \beta$$

$$2\alpha = \frac{3\pi}{2} + \beta$$

$$\cos 2\alpha = \sin \beta$$

$$2 \cos^2 \alpha - 1 = \sin \beta$$


$$2x^2 - 1 = 2x + 1$$

$$x^2 - x - 1 = 0$$

$$\Rightarrow n = \frac{1 \pm \sqrt{5}}{2} = \begin{cases} n = \frac{1+\sqrt{5}}{2} \text{ rejected} \\ n = \frac{1-\sqrt{5}}{2} \end{cases}$$

$$\therefore 4x^2 - 4x = 4$$

$$(2x - 1)^2 = 5$$



Q1. $\vec{b} = \vec{a} \times (\hat{i} - 3\hat{k})$

(1)
$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -1 & 2 \\ 1 & 0 & -2 \end{vmatrix} = 2\hat{i} + 8\hat{j} + \hat{k}$$

$$\vec{c} = \vec{b} \times \hat{k} = 8\hat{i} - 2\hat{j}$$

$$\vec{c} - 2\hat{j} = 8\hat{i} - 4\hat{j}$$

Projection of $(\hat{i} - 2\hat{j})$ on $\vec{a}$

$$(\vec{c} - 2\hat{j}) \cdot \hat{a} = \frac{\langle 8, -4, 0 \rangle \cdot \langle 3, -1, 2 \rangle}{\sqrt{14}}$$

$$= \frac{28}{\sqrt{14}} = 2\sqrt{14}$$

Q2. let

(1) $\vec{a}_{11}$ = component of $\vec{a}$ along $\vec{b}$

$\vec{a}_1$ = component of $\vec{a}$ perpendicular to $\vec{b}$

$$\vec{a}_{11} = \frac{16}{11}(3\hat{i} + \hat{j} - \hat{k})$$

$$\vec{a}_1 = \frac{1}{11}(-4\hat{i} - 5\hat{j} - 17\hat{k})$$

$$\therefore \vec{a} = \vec{a}_{11} + \vec{a}_1$$

$$\therefore \vec{a} =$$

$$= \frac{16}{11}(3\hat{i} + \hat{j} - \hat{k}) + \frac{1}{11}(-4\hat{i} - 5\hat{j} - 17\hat{k})$$

$$\frac{11\hat{j} - 33\hat{k}}{11}$$

$$\alpha = 4\hat{i} + \hat{j} - 3\hat{k}$$

$$\alpha^2 + \beta^2 + \gamma^2 = 16 + 1 + 9 = 26$$

Q3. $\hat{a} \cdot \hat{b} = \frac{1}{2}$

(3) Now $(\lambda\hat{a} + 2\hat{b}) \cdot (3\hat{a} - \lambda\hat{b}) = 0$

$$3\lambda\hat{a} \cdot \hat{a} - \lambda^2\hat{a} \cdot \hat{b} + 6\hat{a} \cdot \hat{b} - 2\lambda\hat{b} \cdot \hat{b} = 0$$

$$3\lambda - \frac{\lambda^2}{2} + 3 - 2\lambda = 0$$


$$\lambda^2 - 2\lambda - 6 = 0$$

$$\lambda = 1 \pm \sqrt{7}$$

$$\Rightarrow \text{number of values} = 0$$

Q4.

(2)



$$\vec{c} = \alpha \vec{a} + \beta \vec{b} \dots (1)$$

$$\vec{a} \cdot \vec{c} = \alpha \vec{a} \cdot \vec{a} + \beta \vec{b} \cdot \vec{a}$$

$$0 = \alpha + \beta \cos 15^\circ \dots (2)$$

$$(1) \Rightarrow \vec{b} \cdot \vec{c} = \alpha \vec{a} \cdot \vec{b} + \beta \vec{b} \cdot \vec{b}$$

$$\Rightarrow \cos 75^\circ = \alpha \cos 15^\circ + \beta \dots (3)$$

$$(2) \& (3) \Rightarrow \cos 75^\circ = -\beta \cos^2 15^\circ + \beta$$

$$\beta = \frac{\cos 75^\circ}{\sin^2 15^\circ} = \frac{1}{\sin 15^\circ} = \frac{2\sqrt{2}}{\sqrt{3}-1}$$

$$(2) \Rightarrow \alpha = \frac{-\cos 15^\circ}{\sin 15^\circ} = \frac{-(\sqrt{3}+1)}{(\sqrt{3}-1)}$$

$$\therefore \vec{c} = \frac{-(\sqrt{3}+1)}{(\sqrt{3}-1)} \vec{a} + \left(\frac{2\sqrt{2}}{\sqrt{3}-1} \right) \vec{b}$$

Now

$$\alpha + \sqrt{2}(\sqrt{3}-1)\beta = \frac{-(\sqrt{3}+1)}{(\sqrt{3}-1)} + \frac{\sqrt{2}(\sqrt{3}-1) \cdot 2\sqrt{2}}{\sqrt{3}-1}$$

$$= \frac{-(\sqrt{3}+1)^2}{2} + 4$$

$$= \frac{-3-1-2\sqrt{3}+8}{2}$$

$$= 2 - \sqrt{3}$$

Q5. $\vec{c} = \lambda(\vec{b} \times (\vec{a} \times \vec{b}))$
(1) $= \lambda((\vec{b} \cdot \vec{b})\vec{a} - (\vec{a} \cdot \vec{b})\vec{b})$
 $= \lambda(11\vec{a} - 2\vec{b}) = \lambda(11\mathbf{i} + 22\mathbf{j} + 33\mathbf{k} - 6\mathbf{i} - 2\mathbf{j} + 2\mathbf{k})$
 $= \lambda(5\mathbf{i} + 20\mathbf{j} + 35\mathbf{k})$
 $= 5\lambda(5\mathbf{i} + 4\mathbf{j} + 7\mathbf{k})$
 $= \text{Given } \vec{c} \cdot \vec{a} = 5$
 $= 5\lambda(1 + 8 + 21) = 5 \Rightarrow \lambda = \frac{1}{30}$

$$\Rightarrow \vec{c} = \frac{1}{6}(\mathbf{i} + 4\mathbf{j} + 7\mathbf{k})$$

$$|\vec{c}| = \frac{\sqrt{1 + 16 + 49}}{6} = \sqrt{\frac{11}{6}}$$

Q6. Equation of line in the internal bisector of OA and OB is $(\sqrt{3} + 1)\hat{i} + (\sqrt{3} + 1)\hat{j}$

(3) $\Rightarrow$ line will be $y = x \Rightarrow x - y = 0$

$$D = \left| \frac{a - (1 - a)}{\sqrt{a^2 + (1 - a)^2}} \right| = \frac{9}{\sqrt{2}}$$

$$(2a - 1)^2 = \frac{81}{2}(a^2 + (1 - a)^2)$$

$$\Rightarrow 2(4a^2 - 4a + 1) = 81a^2 + 81a^2 - 162a - 81$$

$$\Rightarrow 162a^2 - 162a + 81 - 8a^2 + 8a - 2 = 0$$

$$\Rightarrow 154a^2 - 154a + 79 = 0$$

$$\text{Sum of values} = -\frac{(-154)}{154} = 1$$

Q7. $\vec{c} = \left(\frac{\vec{b} \cdot \vec{a}}{|\vec{b}|} \right) \frac{\vec{a}}{|\vec{a}|}$
(16) $= \left(\frac{\lambda + 8}{9} \right) (\hat{i} + 2\hat{j} + 2\hat{k})$

$$|\vec{a} + \vec{c}| = 7$$


$$\Rightarrow \left(\frac{\lambda + 8}{9} + 1 \right) \hat{i} + \left(\frac{2(\lambda + 8)}{9} + 2 \right) \hat{j} + \left(\frac{2(\lambda + 8)}{9} + 2 \right) \hat{k} = 7$$

$$\left(\frac{\lambda + 8}{9} + 1 \right)^2 + \left(\frac{2(\lambda + 8)}{9} + 2 \right)^2 + \left(\frac{2(\lambda + 8)}{9} + 2 \right)^2 = 49$$

$$\Rightarrow \lambda = 4 \Rightarrow \vec{c} = \frac{4}{3}\hat{i} + \frac{8}{3}\hat{j} + \frac{8}{3}\hat{k}$$

$$\text{Area of parallelogram} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{4}{3} & \frac{8}{3} & \frac{8}{3} \\ 4 & 0 & 4 \end{vmatrix} = 16$$

Q8. We know that



$$O \left(\text{orthocentre} \right) \frac{\vec{p} + \vec{q} + \vec{r}}{4}$$

$$C(\text{circum centre}) \alpha \vec{p} + \beta \vec{q} + \gamma \vec{r}$$

$$C(\text{centroid}) = \frac{\vec{p} + \vec{q} + \vec{r}}{3}$$

by relation

$$\Rightarrow 2(\alpha \vec{p} + \beta \vec{q} + \gamma \vec{r}) + \frac{\vec{p} + \vec{q} + \vec{r}}{4} = 3 \left(\frac{\vec{p} + \vec{q} + \vec{r}}{3} \right)$$

$$\Rightarrow 8(\alpha \vec{p} + \beta \vec{q} + \gamma \vec{r}) = 3(\vec{p} + \vec{q} + \vec{r})$$

$$\Rightarrow 8\alpha = 3, 8\beta = 3, 8\gamma = 3$$

$$\alpha = \frac{3}{8}, \beta = \frac{3}{8}, \gamma = \frac{3}{8}$$

$$\therefore \alpha + 2\beta + 3\gamma$$

$$\frac{3}{8} + \frac{6}{8} + \frac{15}{8} = \frac{24}{8} = 3$$

Q9. $\vec{a} = \hat{i} + \hat{j} + \hat{k}$

(6) $\vec{b} = 2\hat{i} + 2\hat{j} + \hat{k}$

$$\vec{d} = \vec{a} \times \vec{b}$$

$$= -\hat{i} + \hat{j}$$

$$|\vec{c} - 2\vec{a}|^2 = 8$$

$$|\vec{c}|^2 + 4|\vec{a}|^2 - 4\vec{a} \cdot \vec{c} = 8$$

$$|\vec{c}|^2 + 12 - 4|\vec{c}| = 8$$

$$|\vec{c}|^2 - 4|\vec{c}| + 4 = 0$$

$$|\vec{c}|^2 = 2$$

$$\vec{d} = \vec{a} \times \vec{b}$$

$$\vec{d} \times \vec{c} = (\vec{a} \times \vec{b}) \times \vec{c}$$

$$\left(|\vec{d}| \times |\vec{c}| \sin \frac{\pi}{4} \right)^2 = ((\vec{a} \cdot \vec{c})\vec{b} - (\vec{b} \cdot \vec{c})\vec{a})^2$$

$$4 = 4|\vec{b}|^2 + (\vec{b} \cdot \vec{c})^2 - 2(\vec{b} \cdot \vec{c})(\vec{a} \cdot \vec{b})$$

Let $\vec{b} \cdot \vec{c} = x$

$$4 = 36 + 3x^2 - 20x$$

$$3x^2 - 20x + 32 = 0$$


$$x = \frac{8}{3}, 4$$

$$\Rightarrow \vec{b} \cdot \vec{c} = \frac{8}{3}, 4$$

$$\Rightarrow \vec{b} \cdot \vec{c} = \frac{8}{3}$$

Now, $|10 - 3\vec{b} \cdot \vec{c}| + |\vec{d} \times \vec{c}|^2$

$$= |10 - 8| + (2)^2 = 6$$



Q10. $\vec{a} = 2\hat{i} - \hat{j} + 3\hat{k}$

(4) $\vec{b} = 3\hat{i} - 5\hat{j} + 3\hat{k}$

$$\vec{a} \times \vec{c} = \vec{c} \times \vec{b}$$

$$\vec{a} \times \vec{c} + \vec{b} \times \vec{c} = 0$$

$$(\vec{a} + \vec{b}) \times \vec{c} = 0$$

$$\Rightarrow \vec{c} = \lambda(\vec{a} + \vec{b})$$

$$\vec{c} = \lambda(5\hat{i} - 6\hat{j} + 4\hat{k}) \dots (1)$$

$$|\vec{c}|^2 = \lambda^2(25 + 36 + 16)$$

$$|\vec{c}|^2 = 77\lambda^2$$

$$(\vec{a} + \vec{c}) \cdot (\vec{b} + \vec{c}) = 168$$

$$\vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} + \vec{c} \cdot \vec{b} + |\vec{c}|^2 = 168$$

$$14 + \vec{c} \cdot (\vec{a} + \vec{b}) + 77\lambda^2 = 168$$

using equation (1)

$$\lambda|5\hat{i} - 6\hat{j} + 4\hat{k}|^2 + 77\lambda^2 = 154$$

$$77\lambda + 77\lambda^2 - 154 = 0$$

$$\lambda^2 + \lambda - 2 = 0$$

$$\lambda = -2, 1$$

$\therefore$ Maximum value of $|\vec{c}|^2$ occurs when $\lambda = -2$

$$|\vec{c}|^2 = 77\lambda^2$$

$$= 77 \times 4$$

$$= 308$$

Q11. (2) $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 3 \\ 2 & 3 & -1 \end{vmatrix} = \hat{i}(-7) + 7\hat{j} + 7\hat{k}$

$$\hat{a} = \pm \frac{(-7\hat{i} + 7\hat{j} + 7\hat{k})}{\sqrt{7^2 + 7^2 + 7^2}} = \pm \left(\frac{-\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}} \right)$$

$$\text{Now, } \cos \theta = \pm \frac{(-1 + 1 + 1)}{\sqrt{3} \cdot \sqrt{3}} = \pm \frac{1}{3}$$

$$\Rightarrow \cos^{-1} \left(\frac{-1}{3} \right) \Rightarrow \hat{a} = \frac{-(-\hat{i} + \hat{j} + \hat{k})}{\sqrt{3}}$$

$$\hat{a} = \frac{\hat{i} - \hat{j} - \hat{k}}{\sqrt{3}}$$

$$\cos \frac{\pi}{3} = \frac{1 - \alpha - 1}{\sqrt{3} \cdot \sqrt{\alpha^2 + 2}}$$

$$\frac{1}{2} = \frac{-\alpha}{\sqrt{3} \cdot \sqrt{\alpha^2 + 2}} \rightarrow \alpha < 0$$

$$3(\alpha^2 + 2) = 4\alpha^2$$


$$6 = \alpha^2$$

$$\alpha = \pm\sqrt{6}$$

$$\text{Clearly, } \alpha = -\sqrt{6}$$

Q12.

(4)



$$\text{Area of } \triangle ABC = \frac{1}{2} |\vec{AB} \times \vec{AC}|$$

$$= \frac{1}{2} |5\hat{i} + 3\hat{j} + \hat{k}| = \frac{1}{2} \sqrt{35}$$

volume of tetrahedron

$$= \frac{1}{3} \times \text{Base area} \times h = \frac{\sqrt{805}}{6\sqrt{2}}$$

$$\frac{1}{3} \times \frac{1}{2} \sqrt{35} \times h = \frac{\sqrt{805}}{6\sqrt{2}}$$

$$h = \sqrt{\frac{23}{2}}$$

Vector Algebra

JEE Main 2025 January

$$AE^2 = AD^2 - DE^2 = \frac{13}{18} \therefore AE = \sqrt{\frac{13}{18}}$$

$$\vec{AE} = |AE| \cdot \left(\frac{\hat{i} - 5\hat{k}}{\sqrt{26}} \right)$$

$$= \sqrt{\frac{13}{18}} \cdot \left(\frac{\hat{i} - 5\hat{k}}{\sqrt{26}} \right)$$

$$= \sqrt{\frac{13}{18}} \cdot \left(\frac{\hat{i} - 5\hat{k}}{\sqrt{26}} \right) = \frac{\hat{i} - 5\hat{k}}{6}$$

$$\text{P.V. of E} = \frac{\hat{i} - 5\hat{k}}{6} + \hat{i} + 2\hat{j} + \hat{k} = \frac{1}{6}(7\hat{i} + 12\hat{j} + \hat{k})$$



Three Dimensional Geometry

JEE Main 2025 January

Q1. Point B

(216) $(3\lambda + 1, -\lambda + 1, -1) \equiv (2\mu + 2, 0, \alpha\mu - 4)$

$$3\lambda + 1 = 2\mu + 2$$

$$-\lambda + 1 = 0$$

$$-1 = \alpha\mu - 4$$

$$\lambda = 1, \mu = 1, \alpha = 3$$

$$B(4, 0, -1)$$

Let Point 'P' is $(2\delta + 2, 0, 3\delta - 4)$

Dr's of AP $\propto 2\delta + 1, -1, 3\delta - 3$

$$AP \perp L_2 \Rightarrow \delta = \frac{7}{13}$$

$$P\left(\frac{40}{13}, 0, \frac{-31}{13}\right)$$

$$\therefore 26\alpha(PB)^2 = 26 \times 3 \times \left(\frac{144}{169} + \frac{324}{169}\right)$$

$$= 216$$

Q2. Line passing through $(1, 4, 0)$ and parallel to $\frac{x}{1} = \frac{y-2}{2} = \frac{z+3}{3}$ is $L : \frac{x-1}{1} = \frac{y-4}{2} = \frac{z}{3}$

(3) Any point on $L : (\lambda + 1, 2\lambda + 4, 3\lambda)$

Any point on $\frac{x-2}{2} = \frac{y-6}{3} = \frac{z-3}{4}$ is $(2\mu + 2, 3\mu + 6, 4\mu + 3)$

$$\left. \begin{array}{l} \lambda + 1 = 2\mu + 2 \\ 2\lambda + 4 = 3\mu + 6 \\ 3\lambda = 4\mu + 3 \end{array} \right\} \lambda = 1\mu = 0$$

Point: $(2, 6, 3)$

$$\text{Distance} = \sqrt{(2-1)^2 + (6-4)^2 + (3-0)^2}$$

$$= \sqrt{1+4+9} = \sqrt{14}$$

Q3. $\vec{a} = (2, 1, -3)$

(2) $\vec{b} = (-1, -3, -5)$

$$\vec{p} \times \vec{q} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -3 \\ 2 & 4 & -5 \end{vmatrix}$$

$$= 2\hat{i} - \hat{j}$$

$$\vec{b} - \vec{a} = -3\hat{i} - 4\hat{j} - 2\hat{k}$$

$$S_d = \frac{|(\vec{b} - \vec{a}) \cdot (\vec{p} \times \vec{q})|}{|\vec{p} \times \vec{q}|}$$

$$= \frac{2}{\sqrt{5}}$$

$$(S_d)^2 = \frac{4}{5}$$


$$m = 4, n = 5 \Rightarrow m + n = 9$$

Three Dimensional Geometry

JEE Main 2025 January

Q4.

(1)



Let coordinate of $R = (2r + 1, r + 2, 3r + 1)$

$\therefore PR$ is perpendicular to given line.

$$\therefore (2r - 3) \cdot 2 + (r - 2) \cdot 1 + (3r - 2) \cdot 3 = 0$$

$$\therefore r = 1$$

$\therefore$ Coordinate of $R = (3, 3, 4)$

$$\therefore (\alpha, \beta, \gamma) = (2, 2, 5)$$

$$\therefore \alpha + \beta + \gamma = 9$$

Q5.

(3)

$$\text{DR's of } L_3 = \vec{m} \times \vec{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ -1 & 2 & 1 \end{vmatrix}$$

$$= -5\hat{i} - 3\hat{j} + \hat{k}$$

$$L_3 : \frac{x - \alpha}{-5} = \frac{y - \beta}{-3} = \frac{z - \gamma}{1} = \lambda$$

$$A(\alpha - 5\lambda, \beta - 3\lambda, \gamma + \lambda)$$

$$L_1 : \frac{x - 1}{1} = \frac{y - 2}{-1} = \frac{z - 1}{2} = k$$

$$B(k + 1, -k + 2, 2k + 1)$$

Now

$$\alpha - 5\lambda = k + 1 \Rightarrow \alpha = 5\lambda + k + 1$$

$$\beta - 3\lambda = -k + 2 \Rightarrow \beta = 3\lambda - k + 2$$

$$\gamma + \lambda = 2k - 1 \Rightarrow \gamma = -\lambda + 2k + 1$$

$$|5\alpha - 11\beta - 8\gamma| = |-25|$$

$$= 25$$

Q6. Vector parallel to L

(4)

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -2 \\ 1 & 3 & 4 \end{vmatrix} = 10\hat{i} - 10\hat{j} + 5\hat{k}$$

$$= 5(2\hat{i} - 2\hat{j} + \hat{k})$$

Equation of ' L '

$$\frac{x-2}{2} = \frac{y+1}{-2} = \frac{z-3}{1} = \lambda(\text{say})$$

Three Dimensional Geometry

JEE Main 2025 January

$$\text{Let } Q(2\lambda + 2, -2\lambda - 1, \lambda + 3)$$

$$\Rightarrow 2\lambda + 2 = 0 \Rightarrow \lambda = -1$$

$$\Rightarrow Q(0, 1, 2)$$

$$d(P, Q) = 3$$

Q7. A(x, y, z) Let P(0, 3, 2), Q(2, 0, 3), R(0, 0, 1)

$$(3) \quad AP = AQ = AR$$

$$x^2 + (y - 3)^2 + (z - 2)^2 = (x - 2)^2 + y^2 + (z - 3)^2 = x^2 + y^2 + (z - 1)^2$$

In xy plane $z = 0$

$$\text{So, } x^2 - 4x + 4 + y^2 + 9 = x^2 + y^2 + 1$$

$$x = 3$$

$$9 + y^2 - 6y + 9 + 4 = x^2 + y^2 + 1$$

So, A(3, 2, 0) also B(1, 4, -1) & C(2, 0, -2)

$$\text{Now } AB = \sqrt{4 + 4 + 1} = 3$$

$$AC = \sqrt{1 + 4 + 4} = 3$$

$$BC = \sqrt{1 + 16 + 1} = \sqrt{18}$$

$$AB = AC$$

$$\text{isosceles } \Delta \& AB^2 + AC^2 = BC^2$$

right angle Δ


$$\text{Area of } \triangle ABC = \frac{1}{2} \times \text{base} \cdot \text{height}$$

$$\frac{1}{2} \times 3 \times 3 = \frac{9}{2}$$

So only S_1 is true

$$\text{Q8. } L_1 : \frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$$

$$(1) \quad L_2 : \frac{x-2}{3} = \frac{y-4}{4} = \frac{z-5}{5}$$



$$P(2\lambda + 1, 3\lambda + 2, 4\lambda + 3)$$

$$Q(3\mu + 2, 4\mu + 4, 5\mu + 5)$$

$$\text{Dir's of } PQ < 2\lambda - 3\mu - 1, 3\lambda - 4\mu - 2,$$

Three Dimensional Geometry

JEE Main 2025 January

$$4\lambda - 5\mu - 2 >$$

$$PQ = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix} = -\hat{i} + 2\hat{j} - \hat{k}$$

$$\Rightarrow \frac{2\lambda - 3\mu - 1}{-1} = \frac{3\lambda - 4\mu - 2}{2} = \frac{4\lambda - 5\mu - 2}{-1}$$

$$\Rightarrow \lambda = \frac{1}{3}\mu = \frac{-1}{6}$$

$$\Rightarrow P\left(\frac{5}{3}, 3, \frac{13}{3}\right) \quad Q\left(\frac{3}{2}, \frac{10}{3}, \frac{25}{6}\right)$$

Dr's $PQ \langle 1, -2, 1 \rangle$

$\therefore$ Line

$$\frac{y - \frac{5}{3}}{1} = \frac{y - 3}{-2} = \frac{y - \frac{13}{3}}{1}$$

Q9. Equation of line PQ is:

$$(2) \quad \frac{x+2}{3} = \frac{y+1}{2} = \frac{z-3}{2} = r \text{ (say)}$$

Let coordinate of $Q = (3r - 2, 2r - 1, 2r + 3)$


$$\therefore PR = 5$$

Then

$$(3r - 2 - 1)^2 + (2r - 1 - 3)^2 + (2r + 3 - 3)^2 = 25$$

$$\therefore r = 0 \text{ or } 2$$

$$\therefore \text{Coordinate of } Q = (4, 3, 7)$$



$$\therefore \text{square of area of } \triangle PQR = \left| \frac{1}{2} (\overrightarrow{PQ} \times \overrightarrow{PR}) \right|^2$$

$$= \left| \frac{1}{2} (6\hat{i} + 4\hat{j} + 4\hat{k}) \times (3\hat{i} + 4\hat{j}) \right|^2$$


$$= | -8\hat{i} + 6\hat{j} + 6\hat{k} |^2 = 136$$

Three Dimensional Geometry

JEE Main 2025 January

Q10.

(4)



$$\frac{x-1}{2} = \frac{y+2}{-1} = \frac{z+3}{2} = \lambda \text{ (let)}$$

$$(2\lambda + 1, -\lambda - 2, 2\lambda - 3)$$

$$\therefore \vec{PA} \cdot \vec{n} = 0$$

$$\Rightarrow (2\lambda - 1)2 + (-\lambda + 8)(-1) + (2\lambda - 4)2 = 0$$

$$\Rightarrow 4\lambda - 2 + \lambda - 8 + 4\lambda - 8 = 0$$


$$\Rightarrow 9\lambda - 18 = 0 \Rightarrow \lambda = 2$$

$$\therefore A(5, -4, 1)$$

$$\therefore AP = \sqrt{3^2 + 6^2 + 0^2} = \sqrt{45} = 3\sqrt{5}$$

Q11.

(28)



$$PR = \operatorname{cosec} \theta, PQ = 4 \sec(30^\circ + \theta)$$

For equilateral

$$d = PR = PQ$$

$$\Rightarrow \cos(\theta + 30^\circ) = 4 \sin \theta$$

$$\Rightarrow \frac{\sqrt{3}}{2} \cos \theta - \frac{1}{2} \sin \theta = 4 \sin \theta$$

$$\Rightarrow \tan \theta = \frac{1}{3\sqrt{3}}$$


$$QR^2 = d^2 = \operatorname{cosec}^2 \theta = 28$$

Three Dimensional Geometry

JEE Main 2025 January

Q12.

(4)



$$\frac{x-3}{7} = \frac{y-2}{-1} = \frac{z+1}{-2} = \lambda$$

$$\Rightarrow 7\lambda + 3, -\lambda + 2, -2\lambda - 1$$

dr's of QP $\Rightarrow$

$$7\lambda - 7, -\lambda + 5, -2\lambda$$

Now

$$(7\lambda - 7) \cdot 7 - (-\lambda + 5) + (2\lambda) \cdot 2 = 0$$

$$54\lambda - 54 = 0 \Rightarrow \lambda = 1$$

$$\therefore P = (10, 1, -3)$$


$$\vec{PQ} = -4\hat{j} + 2\hat{k}$$

$$\vec{PR} = -7\hat{i} - 3\hat{j} + 4\hat{k}$$

$$\text{Area} = \left| \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & -4 & 2 \\ -7 & -3 & 4 \end{vmatrix} \right| = 3\sqrt{30}$$

Q13.

(2)



Three Dimensional Geometry

JEE Main 2025 January

Let $M(3\lambda + 6, 2\lambda + 7, -2\lambda + 7)$

$$\overrightarrow{BM} = (3\lambda + 5)\hat{i} + (2\lambda + 5)\hat{j} + (-2\lambda + 4)\hat{k}$$

$$\overrightarrow{AC} \cdot \overrightarrow{BM} = 0 = 3(3\lambda + 5) + 2(2\lambda + 5) - 2(-2\lambda + 4)$$


$$\overrightarrow{BM} = 2\hat{i} + 3\hat{j} + 6\hat{k}$$

$$|\overrightarrow{BM}| = 7$$

$$\text{Area} = \frac{1}{2} \times 6 \times 7 = 21$$

Q14. Equation of line through point $(-1, 2, 1)$ is $\rightarrow$

(2)



$$\Rightarrow \frac{x+1}{2} = \frac{y-2}{3} = \frac{z-1}{4} = \lambda$$

$$\text{So, } \begin{cases} x = 2\lambda - 1 \\ y = 3\lambda + 2 \\ z = 4\lambda + 1 \end{cases}$$

$$\text{By (1)} \rightarrow \frac{x+2}{3} = \frac{y-3}{2} = \frac{z-4}{1} = \mu \text{ (Let)}$$

$$\text{So, } \begin{cases} x = 3\mu - 2 \\ y = 2\mu + 3 \\ z = \mu + 4 \end{cases}$$

For intersection point 'P'

$$x = 2\lambda - 1 = 3\mu - 2$$

$$y = 3\lambda + 2 = 2\mu + 3$$

$$z = 4\lambda + 1 = \mu + 4$$

So, point $P(x, y, z) = (1, 5, 5)$


& $Q(4, -5, 1)$

$$\therefore PQ = \sqrt{9 + 100 + 16}$$

$$= \sqrt{125} = 5\sqrt{5}$$

Q15.

(957)



Let $R(2\lambda + 1, 3\lambda - 1, 4\lambda)$

$$2\lambda + 1 = 5$$

$$\lambda = 2$$

$$R(5, 5, 8)$$

let $T(2\lambda + 1, 3\lambda - 1, 4\lambda)$

$$\vec{QT} = (2\lambda - 6)\hat{i} + (3\lambda + 1)\hat{j} + (4\lambda - 5)\hat{k}$$

$$\vec{b} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$\vec{QT} \cdot \vec{b} = 0$$

$$4\lambda - 12 + 9\lambda + 3 + 16\lambda - 20 = 0$$

$$\lambda = 1$$

$$T(3, 2, 4)$$


$$QT = \sqrt{33} \quad RT = \sqrt{29}$$

$$(\text{area of } \triangle PQR)^2 = \left(\frac{1}{2} \sqrt{29} \cdot 2\sqrt{33} \right)^2$$

$$= 957$$

Q16.

(4)



$$L = \frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7}$$

$$PQ = \frac{x - \frac{15}{7}}{1} = \frac{y - \frac{32}{7}}{4} = \frac{z - 7}{7} = \lambda$$

Three Dimensional Geometry

JEE Main 2025 January

$$\Rightarrow Q \left(\lambda + \frac{15}{7}, 4\lambda + \frac{32}{7}, 7\lambda + 7 \right)$$

Since Q lies on line L

$$\text{So, } \frac{\lambda + \frac{15}{7} + 1}{3} = \frac{7\lambda + 7 + 5}{7}$$

$$\Rightarrow 7\lambda + 22 = 21\lambda + 36$$

$$\Rightarrow \lambda = -1$$

$$\therefore \text{Point } Q \left(\frac{8}{7}, \frac{4}{7}, 0 \right)$$

$$PQ = \sqrt{\left(\frac{15}{7} - \frac{8}{7} \right)^2 + \left(\frac{32}{7} - \frac{4}{7} \right)^2 + (7 - 0)^2}$$

$$PQ = \sqrt{66}$$

$$\Rightarrow (PQ)^2 = 66$$

Q17. $L_1 : \vec{r} = (-\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} + 2\hat{j} + \hat{k})$

(3) $\Rightarrow \vec{r} = (\lambda - 1)\hat{i} + 2(\lambda + 1)\hat{j} + (\lambda + 1)\hat{k}$

$$L_2 : \vec{r} = (\hat{j} + \hat{k}) + \mu(2\hat{i} + 7\hat{j} + 3\hat{k})$$

$$\Rightarrow \vec{r} = 2\mu\hat{i} + (1 + 7\mu)\hat{j} + (1 + 3\mu)\hat{k}$$

For point of intersection equating respective components

$$\Rightarrow \lambda - 1 = 2\mu$$

$$2(\lambda + 1) = 1 + 7\mu$$

$$\lambda + 1 = 1 + 3\mu$$

We get

$$\Rightarrow \lambda = 3 \text{ and } \mu = 1$$

$$\Rightarrow \vec{a} + \vec{b} = 3\hat{i} + 9\hat{j} + 4\hat{k}$$

$$L_3 : \vec{r} = 2\hat{i} + 8\hat{j} + 4\hat{k} + \alpha(3\hat{i} + 9\hat{j} + 4\hat{k})$$

$$\text{For } \alpha = 2, \vec{r} = 8\hat{i} + 26\hat{j} + 12\hat{k}$$

Q18.

(4) 

Three Dimensional Geometry

JEE Main 2025 January

$$A = \left(\frac{5r-1}{r+1}, \frac{5r-1}{r+1}, \frac{10r+2}{r+1} \right)$$

$$(\vec{OQ} \cdot \vec{OA}) - \frac{1}{5} |\vec{OP} \times \vec{OA}|^2 = 10$$

$$\vec{OQ} = 5\hat{i} + 5\hat{j} + 10\hat{k}$$

$$\vec{OA} = \frac{5r-1}{r+1}\hat{i} + \frac{5r-1}{r+1}\hat{j} + \frac{10r+2}{r+1}\hat{k}$$

$$\vec{OP} = -\hat{i} - \hat{j} + 2\hat{k}$$

$$\vec{OP} \times \vec{OA} = \frac{1}{r+1} \begin{vmatrix} 5r-1 & 5r-1 & 10r+2 \\ 1 & 1 & 2 \end{vmatrix}$$

$$= \frac{1}{r+1} (\hat{i}(20r) - \hat{j}(20r))$$

$$= 5 \left(\frac{5r-1}{r+1} \right) + 5 \left(\frac{5r-1}{r+1} \right) + 10 \left(\frac{10r+2}{r+1} \right)$$

$$- \frac{1}{5} \left(\frac{2 \times 400r^2}{(r+1)^2} \right) = 10$$

$$\frac{150r+10}{r+1} - \frac{1}{5} \left(\frac{2 \times 400r^2}{(r+1)^2} \right) = 10$$

$$(150r+10)(r+1) - 160r^2 = 10(r+1)^2$$

$$(15r+1)(r+1) - 16r^2 = (r+1)^2$$


$$15r^2 + 16r + 1 - 16r^2 = r^2 + 2r + 1$$

$$-2r^2 + 14r = 0$$

$$r = 0, 7$$

Q19. General point on line L : $\frac{x-1}{1} = \frac{y+1}{-1} = \frac{z-2}{2}$

(4) is $(\lambda + 1, -\lambda - 1, 2\lambda + 2)$



DR's of PM are $(\lambda, -\lambda - 3, 2\lambda)$

$PM \perp L$

$$\Rightarrow \lambda + (-1)(-\lambda - 3) + 2(2\lambda) = 0 \Rightarrow 6\lambda + 3 = 0 \Rightarrow \boxed{\lambda = -\frac{1}{2}}$$

Three Dimensional Geometry

JEE Main 2025 January

$$P\left(\frac{1}{2}, \frac{-1}{2}, 1\right)$$

$$\text{Let another line } L' : \frac{x+1}{1} = \frac{y-1}{-1} = \frac{z+2}{1}$$

General point on line L' is $(\mu - 1, -\mu + 1, \mu - 2)$

Point of intersection of line L and L' is

$$\begin{array}{l} \lambda + 1 = \mu - 1 \\ \Rightarrow \mu - \lambda = 2 \dots (1) \end{array} \quad \left| \quad \begin{array}{l} 2\lambda + 2 = \mu - 2 \\ \Rightarrow 2\lambda = \mu - 4 \end{array} \right. \Rightarrow \boxed{\lambda = -2} \text{ and } \boxed{\mu = 0}$$

$$Q(-1, 1, -2)$$

$$2(PQ)^2 = 2 \left(\left(\frac{1}{2} + 1 \right)^2 + \left(\frac{-1}{2} - 1 \right)^2 + (1 + 2)^2 \right)$$

$$= 2 \left(\frac{9}{4} + \frac{9}{4} + 9 \right)$$

$$= 27$$

